



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, 15/62, C07K 14/705, 16/28, C12N 15/70	A2	(11) International Publication Number: WO 98/50547 (43) International Publication Date: 12 November 1998 (12.11.98)
---	-----------	---

(21) International Application Number: PCT/US98/08979

(22) International Filing Date: 7 May 1998 (07.05.98)

(30) Priority Data:

60/044,293	7 May 1997 (07.05.97)	US
60/072,212	22 January 1998 (22.01.98)	US
60/076,947	5 March 1998 (05.03.98)	US

(71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US).

(72) Inventors: HARDIMAN, Gerard, T.; 4 Howe Street, Watertown, MA 02172 (US). ROCK, Fernando, L.; 721 Shell Boulevard #203, Foster City, CA 94404 (US). BAZAN, J., Fernando; 775 University Drive, Menlo Park, CA 94025 (US). KASTELEIN, Robert, A.; 463 Summit Drive, Redwood City, CA 94062 (US).

(74) Agents: McLAUGHLIN, Jaye, P. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).


Published

Without international search report and to be republished upon receipt of that report.

(54) Title: HUMAN TOLL-LIKE RECEPTOR PROTEINS, RELATED REAGENTS AND METHODS

(57) Abstract

Nucleic acids encoding nine human receptors, designated DNAX Toll-like receptors 2-10 (DTLR2-10), homologous to the Drosophila Toll receptor and the human IL-1 receptor, purified DTLR proteins and fragments thereof, mono-/polyclonal antibodies against these receptors, and methods for diagnostic and therapeutic use.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HUMAN RECEPTOR PROTEINS; RELATED REAGENTS AND METHODS

This filing claims priority from U.S. Patent
5 Applications USSN 60/044,293, filed May 7, 1997; USSN
60/072,212, filed January 22, 1998; and USSN 60/076,947,
filed March 5, 1998, each of which is incorporated herein
by reference.

10 FIELD OF THE INVENTION

The present invention relates to compositions and
methods for affecting mammalian physiology, including
morphogenesis or immune system function. In particular,
it provides nucleic acids, proteins, and antibodies which
15 regulate development and/or the immune system.
Diagnostic and therapeutic uses of these materials are
also disclosed.

BACKGROUND OF THE INVENTION

20 Recombinant DNA technology refers generally to
techniques of integrating genetic information from a
donor source into vectors for subsequent processing, such
as through introduction into a host, whereby the
transferred genetic information is copied and/or
25 expressed in the new environment. Commonly, the genetic
information exists in the form of complementary DNA
(cDNA) derived from messenger RNA (mRNA) coding for a
desired protein product. The carrier is frequently a
plasmid having the capacity to incorporate cDNA for later
30 replication in a host and, in some cases, actually to
control expression of the cDNA and thereby direct
synthesis of the encoded product in the host.

For some time, it has been known that the mammalian
immune response is based on a series of complex cellular
35 interactions, called the "immune network". Recent
research has provided new insights into the inner
workings of this network. While it remains clear that

much of the immune response does, in fact, revolve around the network-like interactions of lymphocytes, macrophages, granulocytes, and other cells, immunologists now generally hold the opinion that soluble proteins, known as lymphokines, cytokines, or monokines, play critical roles in controlling these cellular interactions. Thus, there is considerable interest in the isolation, characterization, and mechanisms of action of cell modulatory factors, an understanding of which will lead to significant advancements in the diagnosis and therapy of numerous medical abnormalities, e.g., immune system disorders.

Lymphokines apparently mediate cellular activities in a variety of ways. They have been shown to support the proliferation, growth, and/or differentiation of pluripotential hematopoietic stem cells into vast numbers of progenitors comprising diverse cellular lineages which make up a complex immune system. Proper and balanced interactions between the cellular components are necessary for a healthy immune response. The different cellular lineages often respond in a different manner when lymphokines are administered in conjunction with other agents.

Cell lineages especially important to the immune response include two classes of lymphocytes: B-cells, which can produce and secrete immunoglobulins (proteins with the capability of recognizing and binding to foreign matter to effect its removal), and T-cells of various subsets that secrete lymphokines and induce or suppress the B-cells and various other cells (including other T-cells) making up the immune network. These lymphocytes interact with many other cell types.

Another important cell lineage is the mast cell (which has not been positively identified in all mammalian species), which is a granule-containing connective tissue cell located proximal to capillaries throughout the body. These cells are found in especially

high concentrations in the lungs, skin, and gastrointestinal and genitourinary tracts. Mast cells play a central role in allergy-related disorders, particularly anaphylaxis as follows: when selected
5 antigens crosslink one class of immunoglobulins bound to receptors on the mast cell surface, the mast cell degranulates and releases mediators, e.g., histamine, serotonin, heparin, and prostaglandins, which cause allergic reactions, e.g., anaphylaxis.

10 Research to better understand and treat various immune disorders has been hampered by the general inability to maintain cells of the immune system in vitro. Immunologists have discovered that culturing many of these cells can be accomplished through the use of T-
15 cell and other cell supernatants, which contain various growth factors, including many of the lymphokines.

The interleukin-1 family of proteins includes the IL-1 α , the IL-1 β , the IL-1RA, and recently the IL-1 γ (also designated Interferon-Gamma Inducing Factor, IGIF).
20 This related family of genes have been implicated in a broad range of biological functions. See Dinarello (1994) FASEB J. 8:1314-1325; Dinarello (1991) Blood 77:1627-1652; and Okamura, et al. (1995) Nature 378:88-91.

25 In addition, various growth and regulatory factors exist which modulate morphogenetic development. This includes, e.g., the Toll ligands, which signal through binding to receptors which share structural, and mechanistic, features characteristic of the IL-1
30 receptors. See, e.g., Lemaitre, et al. (1996) Cell 86:973-983; and Belvin and Anderson (1996) Ann. Rev. Cell & Devel. Biol. 12:393-416.

From the foregoing, it is evident that the discovery and development of new soluble proteins and their
35 receptors, including ones similar to lymphokines, should contribute to new therapies for a wide range of degenerative or abnormal conditions which directly or

indirectly involve development, differentiation, or function, e.g., of the immune system and/or hematopoietic cells. In particular, the discovery and understanding of novel receptors for lymphokine-like molecules which enhance or potentiate the beneficial activities of other lymphokines would be highly advantageous. The present invention provides new receptors for ligands exhibiting similarity to interleukin-1 like compositions and related compounds, and methods for their use.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a schematic comparison of the protein architectures of Drosophila and human DTLRs, and their relationship to vertebrate IL-1 receptors and plant disease resistance proteins. Three Drosophila (Dm) DTLRs (Toll, 18w, and the Mst ORF fragment) (Morisato and Anderson (1995) Ann. Rev. Genet. 29:371-399; Chiang and Beachy (1994) Mech. Develop. 47:225-239; Mitcham, et al. (1996) J. Biol. Chem. 271:5777-5783; and Eldon, et al. (1994) Develop. 120:885-899) are arrayed beside four complete (DTLRs 1-4) and one partial (DTLR5) human (Hu) receptors. Individual LRRs in the receptor ectodomains that are flagged by PRINTS (Attwood, et al. (1997) Nucleic Acids Res. 25:212-217) are explicitly noted by boxes; 'top' and 'bottom' Cys-rich clusters that flank the C- or N-terminal ends of LRR arrays are respectively drawn by apposed half-circles. The loss of the internal Cys-rich region in DTLRs 1-5 largely accounts for their smaller ectodomains (558, 570, 690, and 652 aa, respectively) when compared to the 784 and 977 aa extensions of Toll and 18w. The incomplete chains of DmMst and HuDTLR5 (519 and 153 aa ectodomains, respectively) are represented by dashed lines. The intracellular signaling module common to DTLRs, IL-1-type receptors (IL-1Rs), the intracellular protein Myd88, and the tobacco disease resistance gene N product (DRgN) is indicated below the membrane. See, e.g., Hardiman, et

al. (1996) Oncogene 13:2467-2475; and Rock, et al. (1998) Proc. Nat'l Acad. Sci. USA 95:588-. Additional domains include the trio of Ig-like modules in IL-1Rs (disulfide-linked loops); the DRgN protein features an NTPase domain (box) and Myd88 has a death domain (black oval).

Figures 2A-2B show conserved structural patterns in the signaling domains of Toll- and IL-1-like cytokine receptors, and two divergent modular proteins. Figure 2A shows a sequence alignment of the common TH domain.

DTLRs are labeled as in Figure 1; the human (Hu) or mouse (Mo) IL-1 family receptors (IL-1R1-6) are sequentially numbered as earlier proposed (Hardiman, et al. (1996) Oncogene 13:2467-2475); Myd88 and the sequences from tobacco (To) and flax, *L. usitatissimum* (Lu), represent C- and N-terminal domains, respectively, of larger, multidomain molecules. Ungapped blocks of sequence (numbered 1-10) are boxed. Triangles indicate deleterious mutations, while truncations N-terminal of the arrow eliminate bioactivity in human IL-1R1 (Heguy, et al. (1992) J. Biol. Chem. 267:2605-2609). PHD (Rost and Sander (1994) Proteins 19:55-72) and DSC (King and Sternberg (1996) Protein Sci. 5:2298-2310) secondary structure predictions of α -helix (H), β -strand (E), or coil (L) are marked. The amino acid shading scheme depicts chemically similar residues: hydrophobic, acidic, basic, Cys, aromatic, structure-breaking, and tiny. Diagnostic sequence patterns for IL-1Rs, DTLRs, and full alignment (ALL) were derived by Consensus at a stringency of 75%. Symbols for amino acid subsets are (see internet site for detail): o, alcohol; l, aliphatic; ., any amino acid; a, aromatic; c, charged; h, hydrophobic; -, negative; p, polar; +, positive; s, small; u, tiny; t, turnlike. Figure 2B shows a topology diagram of the proposed TH β/α domain fold. The parallel β -sheet (with β -strands A-E as yellow triangles) is seen at its C-terminal end; α -helices (circles labeled 1-5) link the β -strands; chain connections are to the front (visible) or

back (hidden). Conserved, charged residues at the C-end of the β -sheet are noted in gray (Asp) or as a lone black (Arg) residue (see text).

Figure 3 shows evolution of a signaling domain superfamily. The multiple TH module alignment of Figure 2A was used to derive a phylogenetic tree by the Neighbor-Joining method (Thompson, et al. (1994) Nucleic Acids Res. 22:4673-4680). Proteins labeled as in the alignment; the tree was rendered with TreeView.

Figures 4A-4D show FISH chromosomal mapping of human DTLR genes. Denatured chromosomes from synchronous cultures of human lymphocytes were hybridized to biotinylated DTLR cDNA probes for localization. The assignment of the FISH mapping data (left, Figures 4A, DTLR2; 4B, DTLR3; 4C, DTLR4; 4D, DTLR5) with chromosomal bands was achieved by superimposing FISH signals with DAPI banded chromosomes (center panels). Heng and Tsui (1994) Meth. Molec. Biol. 33:109-122. Analyses are summarized in the form of human chromosome ideograms (right panels).

Figures 5A-5F show mRNA blot analyses of Human DTLRs. Human multiple tissue blots (He, heart; Br, brain; Pl, placenta; Lu, lung; Li, liver; Mu, muscle; Ki, kidney; Pn, Pancreas; Sp, spleen; Th, thymus; Pr, prostate; Te, testis; Ov, ovary, SI, small intestine; Co, colon; PBL, peripheral blood lymphocytes) and cancer cell line (promyelocytic leukemia, HL60; cervical cancer, HELAS3; chronic myelogenous leukemia, K562; lymphoblastic leukemia, Molt4; colorectal adenocarcinoma, SW480; melanoma, G361; Burkitt's Lymphoma Raji, Burkitt's; colorectal adenocarcinoma, SW480; lung carcinoma, A549) containing approximately 2 μ g of poly(A)⁺ RNA per lane were probed with radiolabeled cDNAs encoding DTLR1 (Figures 5A-5C), DTLR2 (Figure 5D), DTLR3 (Figure 5E), and DTLR4 (Figure 5F) as described. Blots were exposed to X-ray film for 2 days (Figures 5A-5C) or one week (Figure 5D-5F) at -70° C with intensifying screens. An

anomalous 0.3 kB species appears in some lanes; hybridization experiments exclude a message encoding a DTLR cytoplasmic fragment.

SUMMARY OF THE INVENTION

5 The present invention is directed to nine novel related mammalian receptors, e.g., human, Toll receptor like molecular structures, designated DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9, and DTLR10, and their biological activities. It includes nucleic acids
10 coding for the polypeptides themselves and methods for their production and use. The nucleic acids of the invention are characterized, in part, by their homology to cloned complementary DNA (cDNA) sequences enclosed herein.

15 In certain embodiments, the invention provides a composition of matter selected from the group of: a substantially pure or recombinant DTLR2 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID
20 NO: 4; a natural sequence DTLR2 of SEQ ID NO: 4; a fusion protein comprising DTLR2 sequence; a substantially pure or recombinant DTLR3 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 6; a natural
25 sequence DTLR3 of SEQ ID NO: 6; a fusion protein comprising DTLR3 sequence; a substantially pure or recombinant DTLR4 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 26; a natural sequence
30 DTLR4 of SEQ ID NO: 26; a fusion protein comprising DTLR4 sequence; a substantially pure or recombinant DTLR5 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 10; a natural sequence DTLR5 of SEQ ID NO:
35 10; and a fusion protein comprising DTLR5 sequence.

In other embodiments, the invention provides a composition of matter selected from the group of: a

substantially pure or recombinant DTLR6 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 12; a natural sequence DTLR6 of SEQ ID NO: 12; a
5 fusion protein comprising DTLR6 sequence; a substantially pure or recombinant DTLR7 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 16 or 18 or; a natural sequence DTLR7 of SEQ ID NO: 16 or 18; a fusion
10 protein comprising DTLR7 sequence; a substantially pure or recombinant DTLR8 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 32; a natural sequence DTLR8 of SEQ ID NO: 32; a fusion protein
15 comprising DTLR8 sequence; a substantially pure or recombinant DTLR9 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 22; a natural sequence DTLR9 of SEQ ID NO: 22; and a fusion protein comprising
20 DTLR9 sequence; a substantially pure or recombinant DTLR10 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 34; a natural sequence DTLR10 of SEQ ID NO: 34; and a fusion protein comprising DTLR10
25 sequence.

Preferably, the substantially pure or isolated protein comprises a segment exhibiting sequence identity to a corresponding portion of a DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR 7, DTLR8, DTLR9, or DTLR10, wherein:
30 the homology is at least about 90% identity and the portion is at least about 9 amino acids; the homology is at least about 80% identity and the portion is at least about 17 amino acids; or the homology is at least about 70% identity and the portion is at least about 25 amino
35 acids. In specific embodiments, the composition of matter: is DTLR2, which comprises a mature sequence of SEQ ID NO: 4; or exhibits a post-translational+

modification pattern distinct from natural DTLR2; is DTLR3, which comprises a mature sequence of SEQ ID NO: 6; or exhibits a post-translational modification pattern distinct from natural DTLR3; is DTLR4, which: comprises a mature sequence of SEQ ID NO: 26; or exhibits a post-translational modification pattern distinct from natural DTLR4; or is DTLR5, which: comprises the complete sequence of SEQ ID NO: 10; or exhibits a post-translational modification pattern distinct from natural DTLR5; or is DTLR6, which comprises a mature sequence of SEQ ID NO: 12; or exhibits a post-translational modification pattern distinct from natural DTLR6; is DTLR7, which comprises a mature sequence of SEQ ID NO: 16 or 18; or exhibits a post-translational modification pattern distinct from natural DTLR7; is DTLR8, which: comprises a mature sequence of SEQ ID NO: 32; or exhibits a post-translational modification pattern distinct from natural DTLR8; or is DTLR9, which: comprises the complete sequence of SEQ ID NO: 22; or exhibits a post-translational modification pattern distinct from natural DTLR9; or is DTLR10, which: comprises the complete sequence of SEQ ID NO: 34; or exhibits a post-translational modification pattern distinct from natural DTLR10; or the composition of matter may be a protein or peptide which: is from a warm blooded animal selected from a mammal, including a primate, such as a human; comprises at least one polypeptide segment of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34; exhibits a plurality of portions exhibiting said identity; is a natural allelic variant of DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9, or DTLR10; has a length at least about 30 amino acids; exhibits at least two non-overlapping epitopes which are specific for a primate DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9, or DTLR10; exhibits a sequence identity at least about 90% over a length of at least about 20 amino acids to a primate DTLR2, DTLR3, DTLR4, DTLR5, DTLT6; exhibits at

least two non-overlapping epitopes which are specific for a primate DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9, or DTLR10; exhibits a sequence identity at least about 90% over a length of at least about 20 amino acids to a primate DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9, or DTLR10; is glycosylated; has a molecular weight of at least 100 kD with natural glycosylation; is a synthetic polypeptide; is attached to a solid substrate; is conjugated to another chemical moiety; is a 5-fold or less substitution from natural sequence; or is a deletion or insertion variant from a natural sequence.

Other embodiments include a composition comprising: a sterile DTLR2 protein or peptide; or the DTLR2 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration; a sterile DTLR3 protein or peptide; or the DTLR3 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration; a sterile DTLR4 protein or peptide; or the DTLR4 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration; a sterile DTLR5 protein or peptide; or the DTLR5 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration; a sterile DTLR6 protein or peptide; or the DTLR6 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration; a sterile DTLR7 protein or peptide; or the DTLR7 protein or peptide and a carrier,

wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration; a sterile DTLR8 protein or peptide; or the DTLR8 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration; a sterile DTLR9 protein or peptide; or the DTLR9 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration; a sterile DTLR10 protein or peptide; or the DTLR10 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

In certain fusion protein embodiments, the invention provides a fusion protein comprising: mature protein sequence of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34; a detection or purification tag, including a FLAG, His6, or Ig sequence; or sequence of another receptor protein.

Various kit embodiments include a kit comprising a DTLR protein or polypeptide, and: a compartment comprising the protein or polypeptide; and/or instructions for use or disposal of reagents in the kit. *

Binding compound embodiments include those comprising an antigen binding site from an antibody, which specifically binds to a natural DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9, or DTLR10 protein, wherein: the protein is a primate protein; the binding compound is an Fv, Fab, or Fab2 fragment; the binding compound is conjugated to another chemical moiety; or the antibody: is raised against a peptide sequence of a mature polypeptide of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34; is raised against a mature

DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9 or DTLR10; is raised to a purified human DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9 or DTLR10; is immunoselected; is a polyclonal antibody; binds to a
5 denatured DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9 or DTLR10; exhibits a K_d to antigen of at least 30 μM ; is attached to a solid substrate, including a bead or plastic membrane; is in a sterile composition; or is detectably labeled, including a radioactive or
10 fluorescent label. A binding composition kit often comprises the binding compound, and: a compartment comprising said binding compound; and/or instructions for use or disposal of reagents in the kit. Often the kit is capable of making a qualitative or quantitative analysis.

15 Other compositions include a composition comprising: a sterile binding compound, or the binding compound and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral
20 administration.

Nucleic acid embodiments include an isolated or recombinant nucleic acid encoding a DTLR2-10 protein or peptide or fusion protein, wherein: the DTLR is from a mammal; or the nucleic acid: encodes an antigenic peptide
25 sequence of of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34; encodes a plurality of antigenic peptide sequences of of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34; exhibits at least about 80% identity to a natural cDNA encoding said segment; is an expression
30 vector; further comprises an origin of replication; is from a natural source; comprises a detectable label; comprises synthetic nucleotide sequence; is less than 6 kb, preferably less than 3 kb; is from a mammal, including a primate; comprises a natural full length
35 coding sequence; is a hybridization probe for a gene encoding said DTLR; or is a PCR primer, PCR product, or mutagenesis primer. A cell, tissue, or organ comprising

such a recombinant nucleic acid is also provided.

Preferably, the cell is: a prokaryotic cell; a eukaryotic cell; a bacterial cell; a yeast cell; an insect cell; a mammalian cell; a mouse cell; a primate cell; or a human
5 cell. Kits are provided comprising such nucleic acids, and: a compartment comprising said nucleic acid; a compartment further comprising a primate DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9 or DTLR10 protein or polypeptide; and/or instructions for use or
10 disposal of reagents in the kit. Often, the kit is capable of making a qualitative or quantitative analysis.

Other embodiments include a nucleic acid which:
hybridizes under wash conditions of 30° C and less than 2M salt to SEQ ID NO: 3; hybridizes under wash conditions
15 of 30° C and less than 2 M salt to SEQ ID NO: 5; hybridizes under wash conditions of 30° C and less than 2M salt to SEQ ID NO: 25; hybridizes under wash conditions of 30° C and less than 2 M salt to SEQ ID NO: 9; hybridizes under wash conditions of 30° C and less
20 than 2M salt to SEQ ID NO: 11; hybridizes under wash conditions of 30° C and less than 2 M salt to SEQ ID NO: 15 or 17; hybridizes under wash conditions of 30° C and less than 2M salt to SEQ ID NO: 31; hybridizes under wash conditions of 30° C and less than 2 M salt to SEQ ID NO:
25 21; hybridizes under wash conditions of 30° C and less than 2 M salt to SEQ ID NO: 33; exhibits at least about 85% identity over a stretch of at least about 30 nucleotides to a primate DTLR2 DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9 or DTLR10.

30 Preferably, such nucleic acid will have such properties, wherein: wash conditions are at 45° C and/or 500 mM salt; or the identity is at least 90% and/or the stretch is at least 55 nucleotides. More preferably, the wash conditions are at 55° C and/or 150 mM salt; or the
35 identity is at least 95% and/or the stretch is at least 75 nucleotides.

The invention also provides a method of modulating physiology or development of a cell or tissue culture cells comprising contacting the cell with an agonist or antagonist of a mammalian DTLR2, DTLR3, DTLR4, DTLR5,
5 DTLR6, DTLR7, DTLR8, DTLR9, or DTLR10.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. General

10 The present invention provides the amino acid sequence and DNA sequence of mammalian, herein primate DNAX Toll like receptor molecules (DTLR) having particular defined properties, both structural and biological. These have been designated herein as DTLR2,
15 DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9, and DTLR10, respectively, and increase the number of members of the human Toll like receptor family from 1 to 10. Various cDNAs encoding these molecules were obtained from primate, e.g., human, cDNA sequence libraries. Other
20 primate or other mammalian counterparts would also be desired.

Some of the standard methods applicable are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring
25 Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and periodic supplements) Current Protocols
30 in Molecular Biology, Greene/Wiley, New York; each of which is incorporated herein by reference.

A complete nucleotide and corresponding amino acid sequence of a human DTLR1 coding segment is shown in SEQ ID NO: 1 and 2. See also Nomura, et al. (1994) DNA Res
35 1:27-35. A complete nucleotide and corresponding amino acid sequence of a human DTLR2 coding segment is shown in SEQ ID NO: 3 and 4. A complete nucleotide and

corresponding amino acid sequence of a human DTLR3 coding segment is shown in SEQ ID NO: 5 and 6. A complete nucleotide and corresponding amino acid sequence of a human DTLR4 coding segment is shown in SEQ ID NO: 7 and 8. An alternate nucleic acid and corresponding amino acid sequence of a human DTLR4 coding segment is provided in SEQ ID NO: 25 and 26. A partial nucleotide and corresponding amino acid sequence of a human DTLR5 coding segment is shown in SEQ ID NO: 9 and 10. A complete nucleotide and corresponding amino acid sequence of a human DTLR6 coding segment is shown in SEQ ID NO: 11 and 12 and a partial sequence of a mouse DTLR6 is provided in SEQ ID NO: 13 and 14. Additional mouse DTLR6 sequence is provided in SEQ ID NO: 27 and 29 (nucleotide sequence) and SEQ ID NO: 28 and 30 (amino acid sequence). Partial nucleotide (SEQ ID NO: 15 and 17) and corresponding amino acid sequence (SEQ ID NO: 16 and 18) of a human DTLR7 coding segment is also provided. Partial nucleotide and corresponding amino acid sequence of a human DTLR8 coding segment is shown in SEQ ID NO: 19 and 20. A more complete nucleotide and corresponding amino acid sequence of a human DTLR coding segment is shown in SEQ ID NO: 31 and 32. Partial nucleotide and corresponding amino acid sequence of a human DTLR9 coding segment is shown in SEQ ID NO: 21 and 22. Partial nucleotide and corresponding amino acid sequence of a human DTLR10 coding segment is shown in SEQ ID NO: 23 and 24. More complete nucleotide and corresponding amino acid sequence of a human DTLR10 coding segment is shown in SEQ ID NO: 33 and 34. A partial nucleotide sequence for a mouse DTLR10 coding segment is provided in SEQ ID NO: 35.

Table 1: Comparison of intracellular domains of human DTLRs.
 DTLR1 is SEQ ID NO: 2; DTLR2 is SEQ ID NO: 4; DTLR3 is SEQ ID NO: 6; DTLR4 is SEQ ID NO: 8; DTLR5 is SEQ ID NO: 10; and DTLR6 is SEQ ID NO: 12. Particularly important and conserved, e.g.,
 5 characteristic, residues correspond, across the DTLRs, to SEQ ID NO: 18 residues tyr10-tyr13; trp26; cys46; trp52; pro54-gly55; ser69; lys71; trp134-pro135; and phe144-trp145.

10	DTLR1	QRNLQFHAFISYSGHD---SFWVKNELLPNLEKEG-----MQICLHERNF
	DTLR9	KENLQFHAFISYSEHD---SAWVKSELVPYLEKED-----IQICLHERNF
	DTLR8	-----NELIPNLEKEDGS---ILICLYESYF
	DTLR2	SRNICYDAFVSYSERD---AYWVENLMVQELFNFP---FKLCLHKRDF
	DTLR6	SPDCCYDAFIVYDTKDPVTEWVLAELVAKLEDPREK--HFNLCLLEERDW
	DTLR7	TSQTFYDAYISYDTKDASVTDWVINELRYHLEESRDK--NVLLCLEERDW
15	DTLR10	EDALPYDAFVVFDTKTXSAVADWVYNELRGQLEECRGRW-ALRLCLEERDW
	DTLR4	RGENIYDAFVIYSSQD---EDWVRNELVKNLEEGVPP---FQLCLHYRDF
	DTLR5	PDMYKYDAYLCFSSKD---FTWVQNALLKHLDTQYSDQNRFNLCFEERDF
	DTLR3	TEQFEYAAAYIIHAYKD---KDWVWEHFSSMEKEDQS---LKFCLEERDF
		: . . : * : :
20	DTLR1	VPGKSIVENIITC-IEKSYKSIFVLSPNFVQSEWCH-YELYFAHHNLFHE
	DTLR9	VPGKSIVENIINC-IEKSYKSIFVLSPNFVQSEWCH-YELYFAHHNLFHE
	DTLR8	DPGKSISENIVSF-IEKSYKSIFVLSPNFVQSEWCH-YEFYFAHHNLFHE
	DTLR2	IPGKWIIDNIIDS-IEKSHKTVEVLSENFVKSEWCK-YELDFSHFRLFEE
25	DTLR6	LPGQPVLENLSQS-IQLSKKTVFVMTDKYAKTENFK-IAFYLSHQRLMDE
	DTLR7	DPLGAIIDNLMQS-INQSKKTVFVLTKKYAKSWNFK-TAFYLLXLQRLMGE
	DTLR10	LPGKTLFENLWAS-VYGSRKTLFVLAHTDRVSGLLR-AIFLLAQQRLL-
	DTLR4	IPGVAIAANIIHEGFHKSARKVIVVVSQHFIQSRWCI-FEYEAQTWQFLS
	DTLR5	VPGENRIANIQDA-IWNSRKIVCLVSRHFLRDGWCL-EAFSYAQGRCLSD
30	DTLR3	EAGVFELEAIVNS-IKRSRKIFVITHLLKDPCKRFKVVHHAQQAEQ
		. * : . * * : : : :
	DTLR1	GSNSLILILLEPIPQYSIPSSYHKLKLSLMARRTYLEWPKEKSKRGLFWAN
	DTLR9	GSNNLILILLEPIPQNSIPNKYHKLKALMTQRTYLQWPKEKSKRGLFWA-
35	DTLR8	NSDHIILILLEPIPFYCIPTRYHKLKLEALLEKKAYLEWPKDRRCGLFWAN
	DTLR2	NNDAAILILLEPIEKKAIPQRFCKLRKIMNTKTYLEWPMDEAQRGFWVN
	DTLR6	KVDVILIFLEKPFQK---SKFLQLRKRLCGSSVLEWPTNPQAHFYFWQC
	DTLR7	NMDVIIIFILLEPVLQH---SPYLRLRQRICKSSILQWPDNPKAERLFWQT
	DTLR10	-----
40	DTLR4	SRAGIIFIVLQKVEKT-LLRQQVELYRLLSRNTYLEWEDSVLGRHIFWRR
	DTLR5	LNSALIMVVVGSLSQY-QLMKHQSIRGFVQKQQYLWPEDLQDVGWFLHK
	DTLR3	NLDSIILVFLEEIPDYKLNHALCLRRGMFKSHCILNWPVQKERIGAFRHK
45	DTLR1	LRAAINIKLTEQAKK-----
	DTLR9	-----
	DTLR8	LRAAVNVNVLATREMYELQTFTELNEESRGSTISLMRTDCL
	DTLR2	LRAAIKS-----
	DTLR6	LKNALATDNHVAYSQVPKETV-----
50	DTLR7	LXNVVLTENDSRYNMYVDSIKQY-----
	DTLR10	-----
	DTLR4	LRKALLDGKSWNPEGTVGTGCNWQEATSI-----
	DTLR5	LSQQILKKEKEKKKDNNIPLQTVATIS-----
	DTLR3	LQVALGSKNSVH-----
55		

As used herein, the term DNAX Toll like receptor 2 (DTLR2) shall be used to describe a protein comprising a protein or peptide segment having or sharing the amino acid sequence shown in SEQ ID NO: 4, or a substantial fragment thereof. Similarly, with a DTLR3 and SEQ ID NO: 6; DTLR4 and SEQ ID NO: 26; DTLR5 and SEQ ID NO: 10; DTLR6 and SEQ ID NO: 12; DTLR7 and SEQ ID NO: 16 and 18; DTLR8 and SEQ ID NO: 32; DTLR9 and SEQ ID NO: 22; and DTLR10 and SEQ ID NO: 34.

The invention also includes a protein variations of the respective DTLR allele whose sequence is provided, e.g., a mutein agonist or antagonist. Typically, such agonists or antagonists will exhibit less than about 10% sequence differences, and thus will often have between 1- and 11-fold substitutions, e.g., 2-, 3-, 5-, 7-fold, and others. It also encompasses allelic and other variants, e.g., natural polymorphic, of the protein described. Typically, it will bind to its corresponding biological receptor with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. The term shall also be used herein to refer to related naturally occurring forms, e.g., alleles, polymorphic variants, and metabolic variants of the mammalian protein.

This invention also encompasses proteins or peptides having substantial amino acid sequence identity with the amino acid sequence in SEQ ID NO: 4. It will include sequence variants with relatively few substitutions, e.g., preferably less than about 3-5. Similar features apply to the other DTLR sequences provided in SEQ ID NO: 6, 26, 10, 12, 16, 18, 32, 22 and 34.

A substantial polypeptide "fragment", or "segment", is a stretch of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14

amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids. Sequences of segments of different proteins can be compared to one another over appropriate length stretches.

Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches, if necessary, by introducing gaps as required. See, e.g., Needleham, et al., (1970) J. Mol. Biol. 48:443-453; Sankoff, et al., (1983) chapter one in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison, Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group (GCG), Madison, WI; each of which is incorporated herein by reference. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are intended to include natural allelic and interspecies variations in the cytokine sequence. Typical homologous proteins or peptides will have from 50-100% homology (if gaps can be introduced), to 60-100% homology (if conservative substitutions are included) with an amino acid sequence segment of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34. Homology measures will be at least about 70%, generally at least 76%, more generally at least 81%, often at least 85%, more often at least 88%, typically at least 90%, more typically at least 92%, usually at least 94%, more usually at least 95%,

preferably at least 96%, and more preferably at least 97%, and in particularly preferred embodiments, at least 98% or more. The degree of homology will vary with the length of the compared segments. Homologous proteins or peptides, such as the allelic variants, will share most biological activities with the embodiments described in SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34. Particularly interesting regions of comparison, at the amino acid or nucleotide levels, correspond to those within each of the blocks 1-10, or intrablock regions, corresponding to those indicated in Figure 2A.

As used herein, the term "biological activity" is used to describe, without limitation, effects on inflammatory responses, innate immunity, and/or morphogenic development by respective ligands. For example, these receptors should, like IL-1 receptors, mediate phosphatase or phosphorylase activities, which activities are easily measured by standard procedures. See, e.g., Hardie, et al. (eds. 1995) The Protein Kinase FactBook vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) Meth. Enzymol. 200:38-62; Hunter, et al. (1992) Cell 70:375-388; Lewin (1990) Cell 61:743-752; Pines, et al. (1991) Cold Spring Harbor Symp. Quant. Biol. 56:449-463; and Parker, et al. (1993) Nature 363:736-738. The receptors exhibit biological activities much like regulatable enzymes, regulated by ligand binding. However, the enzyme turnover number is more close to an enzyme than a receptor complex. Moreover, the numbers of occupied receptors necessary to induce such enzymatic activity is less than most receptor systems, and may number closer to dozens per cell, in contrast to most receptors which will trigger at numbers in the thousands per cell. The receptors, or portions thereof, may be useful as phosphate labeling enzymes to label general or specific substrates.

The terms ligand, agonist, antagonist, and analog of, e.g., a DTLR, include molecules that modulate the

characteristic cellular responses to Toll ligand like proteins, as well as molecules possessing the more standard structural binding competition features of ligand-receptor interactions, e.g., where the receptor is a natural receptor or an antibody. The cellular responses likely are mediated through binding of various Toll ligands to cellular receptors related to, but possibly distinct from, the type I or type II IL-1 receptors. See, e.g., Belvin and Anderson (1996) Ann. Rev. Cell Dev. Biol. 12:393-416; Morisato and Anderson (1995) Ann. Rev. Genetics 29:371-3991 and Hultmark (1994) Nature 367:116-117.

Also, a ligand is a molecule which serves either as a natural ligand to which said receptor, or an analog thereof, binds, or a molecule which is a functional analog of the natural ligand. The functional analog may be a ligand with structural modifications, or may be a wholly unrelated molecule which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists, see, e.g., Goodman, et al. (eds) (1990) Goodman & Gilman's: The Pharmacological Bases of Therapeutics, Pergamon Press, New York.

Rational drug design may also be based upon structural studies of the molecular shapes of a receptor or antibody and other effectors or ligands. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein

Crystallography, Academic Press, New York, which is hereby incorporated herein by reference.

II. Activities

5 The Toll like receptor proteins will have a number of different biological activities, e.g., in phosphate metabolism, being added to or removed from specific substrates, typically proteins. Such will generally result in modulation of an inflammatory function, other
10 innate immunity response, or a morphological effect. The DTLR2, 3, 4, 5, 6, 7, 8, 9, or 10 proteins are homologous to other Toll like receptor proteins, but each have structural differences. For example, a human DTLR2 gene coding sequence probably has about 70% identity with the
15 nucleotide coding sequence of mouse DTLR2. At the amino acid level, there is also likely to be reasonable identity.

 The biological activities of the DTLRs will be related to addition or removal of phosphate moieties to
20 substrates, typically in a specific manner, but occasionally in a non specific manner. Substrates may be identified, or conditions for enzymatic activity may be assayed by standard methods, e.g., as described in Hardie, et al. (eds. 1995) The Protein Kinase FactBook
25 vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) Meth. Enzymol. 200:38-62; Hunter, et al. (1992) Cell 70:375-388; Lewin (1990) Cell 61:743-752; Pines, et al. (1991) Cold Spring Harbor Symp. Quant. Biol. 56:449-463; and Parker, et al. (1993) Nature
30 363:736-738.

III. Nucleic Acids

 This invention contemplates use of isolated nucleic acid or fragments, e.g., which encode these or closely
35 related proteins, or fragments thereof, e.g., to encode a corresponding polypeptide, preferably one which is biologically active. In addition, this invention covers

isolated or recombinant DNA which encodes such proteins or polypeptides having characteristic sequences of the respective DTLRs, individually or as a group. Typically, the nucleic acid is capable of hybridizing, under
5 appropriate conditions, with a nucleic acid sequence segment shown in SEQ ID NOs: 3, 5, 25, 9, 11, 15, 17, 31, 21, or 33, but preferably not with a corresponding segment of SEQ ID NO: 1. Said biologically active protein or polypeptide can be a full length protein, or
10 fragment, and will typically have a segment of amino acid sequence highly homologous to one shown in SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34. Further, this invention covers the use of isolated or recombinant nucleic acid, or fragments thereof, which encode proteins
15 having fragments which are equivalent to the DTLR2-10 proteins. The isolated nucleic acids can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others from the natural gene.

20 An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially pure, e.g., separated from other components which naturally accompany a native sequence, such as ribosomes, polymerases, and flanking genomic sequences from the
25 originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates, which are thereby distinguishable from naturally occurring compositions, and chemically
30 synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule, either completely or substantially pure.

An isolated nucleic acid will generally be a
35 homogeneous composition of molecules, but will, in some embodiments, contain heterogeneity, preferably minor. This heterogeneity is typically found at the polymer ends

or portions not critical to a desired biological function or activity.

A "recombinant" nucleic acid is typically defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence. Typically this intervention involves in vitro manipulation, although under certain circumstances it may involve more classical animal breeding techniques. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants as found in their natural state. Thus, for example, products made by transforming cells with any unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such a process is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a restriction enzyme sequence recognition site. Alternatively, the process is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms, e.g., encoding a fusion protein. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. This will include a dimeric repeat. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode equivalent

polypeptides to fragments of DTLR2-10 and fusions of sequences from various different related molecules, e.g., other IL-1 receptor family members.

A "fragment" in a nucleic acid context is a
5 contiguous segment of at least about 17 nucleotides,
generally at least 21 nucleotides, more generally at
least 25 nucleotides, ordinarily at least 30 nucleotides,
more ordinarily at least 35 nucleotides, often at least
39 nucleotides, more often at least 45 nucleotides,
10 typically at least 50 nucleotides, more typically at
least 55 nucleotides, usually at least 60 nucleotides,
more usually at least 66 nucleotides, preferably at least
72 nucleotides, more preferably at least 79 nucleotides,
and in particularly preferred embodiments will be at
15 least 85 or more nucleotides. Typically, fragments of
different genetic sequences can be compared to one
another over appropriate length stretches, particularly
defined segments such as the domains described below.

A nucleic acid which codes for a DTLR2-10 will be
20 particularly useful to identify genes, mRNA, and cDNA
species which code for itself or closely related
proteins, as well as DNAs which code for polymorphic,
allelic, or other genetic variants, e.g., from different
individuals or related species. Preferred probes for
25 such screens are those regions of the interleukin which
are conserved between different polymorphic variants or
which contain nucleotides which lack specificity, and
will preferably be full length or nearly so. In other
situations, polymorphic variant specific sequences will
30 be more useful.

This invention further covers recombinant nucleic
acid molecules and fragments having a nucleic acid
sequence identical to or highly homologous to the
isolated DNA set forth herein. In particular, the
35 sequences will often be operably linked to DNA segments
which control transcription, translation, and DNA

replication. These additional segments typically assist in expression of the desired nucleic acid segment.

Homologous, or highly identical, nucleic acid sequences, when compared to one another or the sequences shown in SEQ ID NO: 3, 5, 25, 9, 11, 15, 17, 31, 21, or 33 exhibit significant similarity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. Comparative hybridization conditions are described in greater detail below.

Substantial identity in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 60% of the nucleotides, generally at least 66%, ordinarily at least 71%, often at least 76%, more often at least 80%, usually at least 84%, more usually at least 88%, typically at least 91%, more typically at least about 93%, preferably at least about 95%, more preferably at least about 96 to 98% or more, and in particular embodiments, as high at about 99% or more of the nucleotides, including, e.g., segments encoding structural domains such as the segments described below. Alternatively, substantial identity will exist when the segments will hybridize under selective hybridization conditions, to a strand or its complement, typically using a sequence derived from SEQ ID NO: 3, 5, 25, 9, 11, 15, 17, 31, 21, or 33.

Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 14 nucleotides, more typically at least about 65%, preferably at least about 75%, and more preferably at least about 90%. See, Kanehisa (1984) Nuc. Acids Res. 12:203-213, which is incorporated herein by reference. The length of homology comparison, as described, may be over longer stretches, and in certain embodiments will be

over a stretch of at least about 17 nucleotides,
generally at least about 20 nucleotides, ordinarily at
least about 24 nucleotides, usually at least about 28
nucleotides, typically at least about 32 nucleotides,
5 more typically at least about 40 nucleotides, preferably
at least about 50 nucleotides, and more preferably at
least about 75 to 100 or more nucleotides.

Stringent conditions, in referring to homology in
the hybridization context, will be stringent combined
10 conditions of salt, temperature, organic solvents, and
other parameters typically controlled in hybridization
reactions. Stringent temperature conditions will usually
include temperatures in excess of about 30° C, more
usually in excess of about 37° C, typically in excess of
15 about 45° C, more typically in excess of about 55° C,
preferably in excess of about 65° C, and more preferably
in excess of about 70° C. Stringent salt conditions will
ordinarily be less than about 500 mM, usually less than
about 400 mM, more usually less than about 300 mM,
20 typically less than about 200 mM, preferably less than
about 100 mM, and more preferably less than about 80 mM,
even down to less than about 20 mM. However, the
combination of parameters is much more important than the
measure of any single parameter. See, e.g., Wetmur and
25 Davidson (1968) J. Mol. Biol. 31:349-370, which is hereby
incorporated herein by reference.

Alternatively, for sequence comparison, typically
one sequence acts as a reference sequence, to which test
sequences are compared. When using a sequence comparison
30 algorithm, test and reference sequences are input into a
computer, subsequence coordinates are designated, if
necessary, and sequence algorithm program parameters are
designated. The sequence comparison algorithm then
calculates the percent sequence identity for the test
35 sequence(s) relative to the reference sequence, based on
the designated program parameters.

Optical alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math. 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443, by the search for similarity method of Pearson and Lipman (1988) Proc. Nat'l Acad. Sci. USA 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally Ausubel et al., supra).

One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng and Doolittle (1987) J. Mol. Evol. 35:351-360. The method used is similar to the method described by Higgins and Sharp (1989) CABIOS 5:151-153. The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. For example, a reference sequence can be compared to other test sequences to determine the percent sequence identity relationship using the following

parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps.

Another example of algorithm that is suitable for determining percent sequence identity and sequence
5 similarity is the BLAST algorithm, which is described Altschul, et al. (1990) J. Mol. Biol. 215:403-410.
Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm
10 involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is
15 referred to as the neighborhood word score threshold (Altschul, et al., supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as
20 far as the cumulative alignment score can be increased. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the
25 accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a wordlength (W) of 11, the
30 BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) Proc. Nat'l Acad. Sci. USA 89:10915) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

In addition to calculating percent sequence
35 identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin and Altschul (1993) Proc. Nat'l Acad. Sci.

USA 90:5873-5787). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

A further indication that two nucleic acid sequences of polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, e.g., where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions, as described below.

The isolated DNA can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode this protein or its derivatives. These modified sequences can be used to produce mutant proteins (muteins) or to enhance the expression of variant species. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant DTLR-like derivatives include predetermined or site-specific mutations of the protein or its fragments, including silent mutations using genetic code degeneracy. "Mutant DTLR" as used herein encompasses a polypeptide otherwise falling within the homology definition of the DTLR as set

forth above, but having an amino acid sequence which differs from that of other DTLR-like proteins as found in nature, whether by way of deletion, substitution, or insertion. In particular, "site specific mutant DTLR" encompasses a protein having substantial homology with a protein of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34, and typically shares most of the biological activities or effects of the forms disclosed herein.

Although site specific mutation sites are predetermined, mutants need not be site specific. Mammalian DTLR mutagenesis can be achieved by making amino acid insertions or deletions in the gene, coupled with expression. Substitutions, deletions, insertions, or any combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy-terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mammalian DTLR mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis. See also Sambrook, et al. (1989) and Ausubel, et al. (1987 and periodic Supplements).

The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Polymerase chain reaction (PCR) techniques can often be applied in mutagenesis. Alternatively, mutagenesis primers are commonly used methods for generating defined mutations at predetermined sites. See, e.g., Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach and Dveksler (1995; eds.) PCR Primer: A Laboratory Manual Cold Spring Harbor Press, CSH, NY.

10 IV. Proteins, Peptides

As described above, the present invention encompasses primate DTLR2-10, e.g., whose sequences are disclosed in SEQ ID NOS: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34, and described above. Allelic and other variants
15 are also contemplated, including, e.g., fusion proteins combining portions of such sequences with others, including epitope tags and functional domains.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using
20 segments from these rodent proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. Thus, the fusion product of a DTLR with an IL-1 receptor is a continuous protein molecule having sequences fused
25 in a typical peptide linkage, typically made as a single translation product and exhibiting properties, e.g., sequence or antigenicity, derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences.

30 In addition, new constructs may be made from combining similar functional or structural domains from other related proteins, e.g., IL-1 receptors or other DTLRs, including species variants. For example, ligand-binding or other segments may be "swapped" between
35 different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992,

each of which is incorporated herein by reference. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of receptor-binding specificities. For example, the ligand binding domains from other related receptor molecules may be added or substituted for other domains of this or related proteins. The resulting protein will often have hybrid function and properties. For example, a fusion protein may include a targetting domain which may serve to provide sequestering of the fusion protein to a particular subcellular organelle.

Candidate fusion partners and sequences can be selected from various sequence data bases, e.g., GenBank, c/o IntelliGenetics, Mountain View, CA; and BCG, University of Wisconsin Biotechnology Computing Group, Madison, WI, which are each incorporated herein by reference.

The present invention particularly provides muteins which bind Toll ligands, and/or which are affected in signal transduction. Structural alignment of human DTLR1-10 with other members of the IL-1 family show conserved features/residues. See, e.g., Figure 3A. Alignment of the human DTLR sequences with other members of the IL-1 family indicates various structural and functionally shared features. See also, Bazan, et al. (1996) Nature 379:591; Lodi, et al. (1994) Science 263:1762-1766; Sayle and Milner-White (1995) TIBS 20:374-376; and Gronenberg, et al. (1991) Protein Engineering 4:263-269.

The IL-1 α and IL-1 β ligands bind an IL-1 receptor type I as the primary receptor and this complex then forms a high affinity receptor complex with the IL-1 receptor type III. Such receptor subunits are probably shared with the new IL-1 family members.

Similar variations in other species counterparts of DTLR2-10 sequences, e.g., in the corresponding regions, should provide similar interactions with ligand or

substrate. Substitutions with either mouse sequences or human sequences are particularly preferred. Conversely, conservative substitutions away from the ligand binding interaction regions will probably preserve most signaling activities.

"Derivatives" of the primate DTLR2-10 include amino acid sequence mutants, glycosylation variants, metabolic derivatives and covalent or aggregative conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in the DTLR amino acid side chains or at the N- or C- termini, e.g., by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl-moieties including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the receptors or fragments thereof with other proteins of polypeptides. These derivatives can be synthesized in

recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred derivatization sites with
5 cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between the receptors and other homologous or heterologous proteins are also provided. Homologous polypeptides may be fusions between different
10 receptors, resulting in, for instance, a hybrid protein exhibiting binding specificity for multiple different Toll ligands, or a receptor which may have broadened or weakened specificity of substrate effect. Likewise, heterologous fusions may be constructed which would
15 exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a receptor, e.g., a ligand-binding segment, so that the presence or location of a desired ligand may be
20 easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609, which is hereby incorporated herein by reference. Other gene fusion partners include glutathione-S-transferase (GST), bacterial β -galactosidase, trpE, Protein A, β -lactamase, alpha
25 amylase, alcohol dehydrogenase, and yeast alpha mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will
30 produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an
35 appropriate primer sequence.

Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation,

sulfonation, biotinylation, or the addition or removal of other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments, the modifications will be useful labeling reagents, or
5 serve as purification targets, e.g., affinity ligands.

Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, for
10 example, in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1-3, Cold Spring Harbor Laboratory, and Ausubel, et al. (eds. 1987 and periodic supplements) Current Protocols in Molecular Biology,
15 Greene/Wiley, New York, which are each incorporated herein by reference. Techniques for synthesis of polypeptides are described, for example, in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL
20 Press, Oxford; each of which is incorporated herein by reference. See also Dawson, et al. (1994) Science 266:776-779 for methods to make larger polypeptides.

This invention also contemplates the use of derivatives of a DTLR2-10 other than variations in amino
25 acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, for
30 example with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of a receptor or other binding molecule, e.g., an antibody. For example, a Toll
35 ligand can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated Sepharose, by methods which are well known in the art, or adsorbed onto

polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of a DTLR receptor, antibodies, or other similar molecules. The ligand can also be labeled with a detectable group, for example radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays.

A DTLR of this invention can be used as an immunogen for the production of antisera or antibodies specific, e.g., capable of distinguishing between other IL-1 receptor family members, for the DTLR or various fragments thereof. The purified DTLR can be used to screen monoclonal antibodies or antigen-binding fragments prepared by immunization with various forms of impure preparations containing the protein. In particular, the term "antibodies" also encompasses antigen binding fragments of natural antibodies; e.g., Fab, Fab2, Fv, etc. The purified DTLR can also be used as a reagent to detect antibodies generated in response to the presence of elevated levels of expression, or immunological disorders which lead to antibody production to the endogenous receptor. Additionally, DTLR fragments may also serve as immunogens to produce the antibodies of the present invention, as described immediately below. For example, this invention contemplates antibodies having binding affinity to or being raised against the amino acid sequences shown in SEQ ID NOS: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34, fragments thereof, or various homologous peptides. In particular, this invention contemplates antibodies having binding affinity to, or having been raised against, specific fragments which are predicted to be, or actually are, exposed at the exterior protein surface of the native DTLR.

The blocking of physiological response to the receptor ligands may result from the inhibition of binding of the ligand to the receptor, likely through competitive inhibition. Thus, in vitro assays of the

present invention will often use antibodies or antigen binding segments of these antibodies, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either ligand binding region mutations and modifications, or other mutations and modifications, e.g., which affect signaling or enzymatic function.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to the receptor or fragments compete with a test compound for binding to a ligand or other antibody. In this manner, the neutralizing antibodies or fragments can be used to detect the presence of a polypeptide which shares one or more binding sites to a receptor and can also be used to occupy binding sites on a receptor that might otherwise bind a ligand.

V. Making Nucleic Acids and Protein

DNA which encodes the protein or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples. Natural sequences can be isolated using standard methods and the sequences provided herein. Other species counterparts can be identified by hybridization techniques, or by various PCR techniques, combined with or by searching in sequence databases, e.g., GenBank.

This DNA can be expressed in a wide variety of host cells for the synthesis of a full-length receptor or fragments which can in turn, for example, be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified ligand binding or kinase/phosphatase domains; and for structure/function studies. Variants or fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These

molecules can be substantially free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The protein, or portions thereof, may be expressed as fusions with other proteins.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired receptor gene or its fragments, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control elements are capable of effecting expression within a suitable host. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

The vectors of this invention include those which contain DNA which encodes a protein, as described, or a fragment thereof encoding a biologically active equivalent polypeptide. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNA coding for such a protein in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNA coding for the receptor is inserted into the vector such

that growth of the host containing the vector expresses the cDNA in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of
5 copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the protein or its fragments in various hosts using vectors that do not contain a replication
10 origin that is recognized by the host cell. It is also possible to use vectors that cause integration of the protein encoding portion or its fragments into the host DNA by recombination.

Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other
15 vehicles which enable the integration of DNA fragments into the genome of the host. Expression vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids are the most commonly used form of vector but
20 all other forms of vectors which serve an equivalent function and which are, or become, known in the art are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory
25 Manual, Elsevier, N.Y., and Rodriguez, et al. (eds) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, 1988, which are incorporated herein by reference.

Transformed cells are cells, preferably mammalian,
30 that have been transformed or transfected with receptor vectors constructed using recombinant DNA techniques. Transformed host cells usually express the desired protein or its fragments, but for purposes of cloning, amplifying, and manipulating its DNA, do not need to
35 express the subject protein. This invention further contemplates culturing transformed cells in a nutrient medium, thus permitting the receptor to accumulate in the

cell membrane. The protein can be recovered, either from the culture or, in certain instances, from the culture medium.

For purposes of this invention, nucleic sequences
5 are operably linked when they are functionally related to each other. For example, DNA for a presequence or secretory leader is operably linked to a polypeptide if it is expressed as a preprotein or participates in directing the polypeptide to the cell membrane or in
10 secretion of the polypeptide. A promoter is operably linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to permit translation. Usually, operably
15 linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression.

Suitable host cells include prokaryotes, lower
20 eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., E. coli and B. subtilis. Lower eukaryotes include yeasts, e.g., S. cerevisiae and Pichia, and species of the genus Dictyostelium. Higher eukaryotes include established
25 tissue culture cell lines from animal cells, both of non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used
30 herein, E. coli and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express the receptor or its fragments include, but are
35 not limited to, such vectors as those containing the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters

(pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-, and Ipp-derived Promoters", in Vectors: A Survey of Molecular Cloning Vectors and Their
5 Uses, (eds. Rodriguez and Denhardt), Buttersworth, Boston, Chapter 10, pp. 205-236, which is incorporated herein by reference.

Lower eukaryotes, e.g., yeasts and Dictyostelium, may be transformed with DTLR sequence containing vectors.
10 For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, Saccharomyces cerevisiae. It will be used to generically represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically
15 consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the receptor or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors
20 for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Suitable vectors include derivatives of the
25 following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YEp-series); integrating types (such as the YIp-series), or mini-chromosomes (such as the YCp-series).

30 Higher eukaryotic tissue culture cells are normally the preferred host cells for expression of the functionally active interleukin protein. In principle, any higher eukaryotic tissue culture cell line is workable, e.g., insect baculovirus expression systems,
35 whether from an invertebrate or vertebrate source. However, mammalian cells are preferred. Transformation or transfection and propagation of such cells has become

a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines.

5 Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a

10 selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of

15 suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo PolyA, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

For secreted proteins, an open reading frame usually

20 encodes a polypeptide that consists of a mature or secreted product covalently linked at its N-terminus to a signal peptide. The signal peptide is cleaved prior to secretion of the mature, or active, polypeptide. The cleavage site can be predicted with a high degree of

25 accuracy from empirical rules, e.g., von-Heijne (1986) Nucleic Acids Research 14:4683-4690, and the precise amino acid composition of the signal peptide does not appear to be critical to its function, e.g., Randall, et al. (1989) Science 243:1156-1159; Kaiser et al. (1987)

30 Science 235:312-317.

It will often be desired to express these polypeptides in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression

35 system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a

heterologous expression system. For example, the receptor gene may be co-transformed with one or more genes encoding mammalian or other glycosylating enzymes. Using this approach, certain mammalian glycosylation
5 patterns will be achievable in prokaryote or other cells.

The source of DTLR can be a eukaryotic or prokaryotic host expressing recombinant DTLR, such as is described above. The source can also be a cell line such as mouse Swiss 3T3 fibroblasts, but other mammalian cell
10 lines are also contemplated by this invention, with the preferred cell line being from the human species.

Now that the sequences are known, the primate DTLRs, fragments, or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These
15 include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis, Springer-Verlag, New York; and Bodanszky (1984) The
20 Principles of Peptide Synthesis, Springer-Verlag, New York; all of each which are incorporated herein by reference. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (e.g.,
25 p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimide process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both
30 applicable to the foregoing processes. Similar techniques can be used with partial DTLR sequences.

The DTLR proteins, fragments, or derivatives are suitably prepared in accordance with the above processes as typically employed in peptide synthesis, generally
35 either by a so-called stepwise process which comprises condensing an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to

the terminal amino acid. Amino groups that are not being used in the coupling reaction typically must be protected to prevent coupling at an incorrect location.

If a solid phase synthesis is adopted, the
5 C-terminal amino acid is bound to an insoluble carrier or support through its carboxyl group. The insoluble carrier is not particularly limited as long as it has a binding capability to a reactive carboxyl group. Examples of such insoluble carriers include halomethyl
10 resins, such as chloromethyl resin or bromomethyl resin, hydroxymethyl resins, phenol resins, tert-alkyloxycarbonylhydrazidated resins, and the like.

An amino group-protected amino acid is bound in sequence through condensation of its activated carboxyl
15 group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After synthesizing the complete sequence, the peptide is split off from the insoluble carrier to produce the peptide. This solid-phase approach is
20 generally described by Merrifield, et al. (1963) in J. Am. Chem. Soc. 85:2149-2156, which is incorporated herein by reference.

The prepared protein and fragments thereof can be isolated and purified from the reaction mixture by means
25 of peptide separation, for example, by extraction, precipitation, electrophoresis, various forms of chromatography, and the like. The receptors of this invention can be obtained in varying degrees of purity depending upon desired uses. Purification can be
30 accomplished by use of the protein purification techniques disclosed herein, see below, or by the use of the antibodies herein described in methods of immunoabsorbant affinity chromatography. This immunoabsorbant affinity chromatography is carried out by
35 first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate cells, lysates of other cells expressing

the receptor, or lysates or supernatants of cells producing the protein as a result of DNA techniques, see below.

Generally, the purified protein will be at least
5 about 40% pure, ordinarily at least about 50% pure, usually at least about 60% pure, typically at least about 70% pure, more typically at least about 80% pure, preferable at least about 90% pure and more preferably at least about 95% pure, and in particular embodiments, 97%-
10 99% or more. Purity will usually be on a weight basis, but can also be on a molar basis. Different assays will be applied as appropriate.

VI. Antibodies

15 Antibodies can be raised to the various mammalian, e.g., primate DTLR proteins and fragments thereof, both in naturally occurring native forms and in their recombinant forms, the difference being that antibodies to the active receptor are more likely to recognize
20 epitopes which are only present in the native conformations. Denatured antigen detection can also be useful in, e.g., Western analysis. Anti-idiotypic antibodies are also contemplated, which would be useful as agonists or antagonists of a natural receptor or an
25 antibody.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of the protein can be raised by immunization of animals with conjugates of the fragments with immunogenic proteins.
30 Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective protein, or screened for agonistic or antagonistic activity. These monoclonal antibodies will usually bind with at least a K_D of about
35 1 mM, more usually at least about 300 μ M, typically at least about 100 μ M, more typically at least about 30 μ M,

preferably at least about 10 μ M, and more preferably at least about 3 μ M or better.

The antibodies, including antigen binding fragments, of this invention can have significant diagnostic or therapeutic value. They can be potent antagonists that bind to the receptor and inhibit binding to ligand or inhibit the ability of the receptor to elicit a biological response, e.g., act on its substrate. They also can be useful as non-neutralizing antibodies and can be coupled to toxins or radionuclides to bind producing cells, or cells localized to the source of the interleukin. Further, these antibodies can be conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker.

The antibodies of this invention can also be useful in diagnostic applications. As capture or non-neutralizing antibodies, they might bind to the receptor without inhibiting ligand or substrate binding. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying ligand. They may be used as reagents for Western blot analysis, or for immunoprecipitation or immunopurification of the respective protein.

Protein fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. Mammalian DTLR and its fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York; and Williams, et al. (1967) Methods in Immunology and Immunochemistry, Vol. 1, Academic Press, New York; each of which are incorporated herein by reference, for descriptions of methods of preparing polyclonal antisera. A typical

method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin is isolated.

5 In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds) Basic and
10 Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed) Academic Press, New York;
15 and particularly in Kohler and Milstein (1975) in Nature 256: 495-497, which discusses one method of generating monoclonal antibodies. Each of these references is incorporated herein by reference. Summarized briefly, this method involves injecting an animal with an
20 immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones,
25 each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic
30 substance.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989)
35 "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-

546, each of which is hereby incorporated herein by reference. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. 5 Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific 10 and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 15 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant or chimeric immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; or made in transgenic mice, see Mendez, et al. (1997) Nature Genetics 15:146-156. These references 20 are incorporated herein by reference.

The antibodies of this invention can also be used for affinity chromatography in isolating the DTLRs. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, 25 Sephadex, or the like, where a cell lysate may be passed through the column, the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified protein will be released. The protein may be used to purify antibody.

30 The antibodies may also be used to screen expression libraries for particular expression products. Usually the antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

35 Antibodies raised against a DTLR will also be used to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological

conditions related to expression of the protein or cells which express the protein. They also will be useful as agonists or antagonists of the ligand, which may be competitive inhibitors or substitutes for naturally occurring ligands.

A DTLR protein that specifically binds to or that is specifically immunoreactive with an antibody generated against a defined immunogen, such as an immunogen consisting of the amino acid sequence of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34, is typically determined in an immunoassay. The immunoassay typically uses a polyclonal antiserum which was raised, e.g., to a protein of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34. This antiserum is selected to have low crossreactivity against other IL-1R family members, e.g., DTLR1, preferably from the same species, and any such crossreactivity is removed by immunoabsorption prior to use in the immunoassay.

In order to produce antisera for use in an immunoassay, the protein of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34, or a combination thereof, is isolated as described herein. For example, recombinant protein may be produced in a mammalian cell line. An appropriate host, e.g., an inbred strain of mice such as balb/c, is immunized with the selected protein, typically using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane, supra). Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen. Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, e.g., a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of 10^4 or greater are selected and tested for their cross reactivity against other IL-1R family members, e.g., mouse DTLRs or human DTLR1, using a competitive binding

immunoassay such as the one described in Harlow and Lane, supra, at pages 570-573. Preferably at least two DTLR family members are used in this determination in conjunction with either or some of the human DTLR2-10.

- 5 These IL-1R family members can be produced as recombinant proteins and isolated using standard molecular biology and protein chemistry techniques as described herein.

Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For
10 example, the proteins of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34, or various fragments thereof, can be immobilized to a solid support. Proteins added to the assay compete with the binding of the antisera to the immobilized antigen. The ability of the above proteins
15 to compete with the binding of the antisera to the immobilized protein is compared to the protein of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 and/or 34. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera
20 with less than 10% crossreactivity with each of the proteins listed above are selected and pooled. The cross-reacting antibodies are then removed from the pooled antisera by immunoabsorbtion with the above-listed proteins.

- 25 The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein to the immunogen protein (e.g., the IL-1R like protein of SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 and/or 34). In order to make this
30 comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required is less than twice the amount
35 of the protein of the selected protein or proteins that is required, then the second protein is said to

specifically bind to an antibody generated to the immunogen.

It is understood that these DTLR proteins are members of a family of homologous proteins that comprise at least 10 so far identified genes. For a particular gene product, such as the DTLR2-10, the term refers not only to the amino acid sequences disclosed herein, but also to other proteins that are allelic, non-allelic or species variants. It also understood that the terms include nonnatural mutations introduced by deliberate mutation using conventional recombinant technology such as single site mutation, or by excising short sections of DNA encoding the respective proteins, or by substituting new amino acids, or adding new amino acids. Such minor alterations must substantially maintain the immunoidentity of the original molecule and/or its biological activity. Thus, these alterations include proteins that are specifically immunoreactive with a designated naturally occurring IL-1R related protein, for example, the DTLR proteins shown in SEQ ID NO: 4, 6, 26, 10, 12, 16, 18, 32, 22 or 34. The biological properties of the altered proteins can be determined by expressing the protein in an appropriate cell line and measuring the appropriate effect upon lymphocytes. Particular protein modifications considered minor would include conservative substitution of amino acids with similar chemical properties, as described above for the IL-1R family as a whole. By aligning a protein optimally with the protein of DTLR2-10 and by using the conventional immunoassays described herein to determine immunoidentity, one can determine the protein compositions of the invention.

VII. Kits and quantitation

Both naturally occurring and recombinant forms of the IL-1R like molecules of this invention are particularly useful in kits and assay methods. For example, these methods would also be applied to screening

for binding activity, e.g., ligands for these proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds per year. See, e.g., a BIOMEK
5 automated workstation, Beckman Instruments, Palo Alto, California, and Fodor, et al. (1991) Science 251:767-773, which is incorporated herein by reference. The latter describes means for testing binding by a plurality of defined polymers synthesized on a solid substrate. The
10 development of suitable assays to screen for a ligand or agonist/antagonist homologous proteins can be greatly facilitated by the availability of large amounts of purified, soluble DTLRs in an active state such as is provided by this invention.

15 Purified DTLR can be coated directly onto plates for use in the aforementioned ligand screening techniques. However, non-neutralizing antibodies to these proteins can be used as capture antibodies to immobilize the respective receptor on the solid phase, useful, e.g., in
20 diagnostic uses.

This invention also contemplates use of DTLR2-10, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of the protein or its ligand.

25 Alternatively, or additionally, antibodies against the molecules may be incorporated into the kits and methods. Typically the kit will have a compartment containing either a defined DTLR peptide or gene segment or a reagent which recognizes one or the other. Typically,
30 recognition reagents, in the case of peptide, would be a receptor or antibody, or in the case of a gene segment, would usually be a hybridization probe.

A preferred kit for determining the concentration of, e.g., DTLR4, a sample would typically comprise a
35 labeled compound, e.g., ligand or antibody, having known binding affinity for DTLR4, a source of DTLR4 (naturally occurring or recombinant) as a positive control, and a

means for separating the bound from free labeled compound, for example a solid phase for immobilizing the DTLR4 in the test sample. Compartments containing reagents, and instructions, will normally be provided.

5 Antibodies, including antigen binding fragments, specific for mammalian DTLR or a peptide fragment, or receptor fragments are useful in diagnostic applications to detect the presence of elevated levels of ligand and/or its fragments. Diagnostic assays may be
10 homogeneous (without a separation step between free reagent and antibody-antigen complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA),
15 enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA) and the like. For example, unlabeled antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to DTLR4 or to a particular
20 fragment thereof. These assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH., and Coligan (Ed.) (1991) and periodic supplements, Current Protocols In Immunology Greene/Wiley, New York.

25 Anti-idiotypic antibodies may have similar use to serve as agonists or antagonists of DTLR4. These should be useful as therapeutic reagents under appropriate circumstances.

 Frequently, the reagents for diagnostic assays are
30 supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody, or labeled ligand is provided. This is usually in conjunction with other
35 additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also

contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent, and will contain instructions for proper use and disposal of reagents. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium having appropriate concentrations for performing the assay.

The aforementioned constituents of the diagnostic assays may be used without modification or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, a test compound, DTLR, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as ^{125}I , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Both of the patents are incorporated herein by reference. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound from the free ligand, or alternatively the bound from the free test compound. The DTLR can be immobilized on various matrixes followed by washing. Suitable matrices include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the receptor to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of antibody/antigen complex by any of several methods including those utilizing, e.g.,

an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in
5 Rattle, et al. (1984) Clin. Chem. 30(9):1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678, each of which is incorporated herein by reference.

The methods for linking protein or fragments to
10 various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of
15 thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves
20 use of oligonucleotide or polynucleotide sequences taken from the sequence of a DTLR. These sequences can be used as probes for detecting levels of the respective DTLR in patients suspected of having an immunological disorder. The preparation of both RNA and DNA nucleotide sequences,
25 the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least about 18 nucleotides, and
30 the polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly ^{32}P . However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a
35 polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides,

fluorescers, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies
5 in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected. The use of probes to the novel anti-sense RNA may be carried out in
10 any conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART). This also includes amplification techniques such as polymerase chain
15 reaction (PCR).

Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers.
20 Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97.

VIII. Therapeutic Utility

25 This invention provides reagents with significant therapeutic value. The DTLRs (naturally occurring or recombinant), fragments thereof, mutein receptors, and antibodies, along with compounds identified as having binding affinity to the receptors or antibodies, should
30 be useful in the treatment of conditions exhibiting abnormal expression of the receptors of their ligands. Such abnormality will typically be manifested by immunological disorders. Additionally, this invention should provide therapeutic value in various diseases or
35 disorders associated with abnormal expression or abnormal triggering of response to the ligand. The Toll ligands have been suggested to be involved in morphologic

development, e.g., dorso-ventral polarity determination, and immune responses, particularly the primitive innate responses. See, e.g., Sun, et al. (1991) Eur. J. Biochem. 196:247-254; Hultmark (1994) Nature 367:116-117.

5 Recombinant DTLRs, muteins, agonist or antagonist antibodies thereto, or antibodies can be purified and then administered to a patient. These reagents can be combined for therapeutic use with additional active ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, along with
10 physiologically innocuous stabilizers and excipients. These combinations can be sterile, e.g., filtered, and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations.
15 This invention also contemplates use of antibodies or binding fragments thereof which are not complement binding.

 Ligand screening using DTLR or fragments thereof can be performed to identify molecules having binding
20 affinity to the receptors. Subsequent biological assays can then be utilized to determine if a putative ligand can provide competitive binding, which can block intrinsic stimulating activity. Receptor fragments can be used as a blocker or antagonist in that it blocks the
25 activity of ligand. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of ligand, e.g., inducing signaling. This invention further contemplates the therapeutic use of antibodies to
30 DTLRs as antagonists.

 The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medicants
35 administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts

useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds) (1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, (current edition), Mack Publishing Co., Easton, Penn.; each of which is hereby incorporated herein by reference. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Because of the likely high affinity binding, or turnover numbers, between a putative ligand and its receptors, low dosages of these reagents would be initially expected to be effective. And the signaling pathway suggests extremely low amounts of ligand may have effect. Thus, dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10 μ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or slow release apparatus will often be utilized for continuous administration.

DTLRs, fragments thereof, and antibodies or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in any conventional dosage formulation. While it is possible for the active

ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier must be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds) (1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences (current edition), Mack Publishing Co., Easton, Penn.; Avis, et al. (eds. 1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, NY; Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Tablets Dekker, NY; and Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, NY. The therapy of this invention may be combined with or used in association with other therapeutic agents, particularly agonists or antagonists of other IL-1 family members.

IX. Ligands

The description of the Toll receptors herein provide means to identify ligands, as described above. Such ligand should bind specifically to the respective receptor with reasonably high affinity. Various constructs are made available which allow either labeling of the receptor to detect its ligand. For example, directly labeling DTLR, fusing onto it markers for secondary labeling, e.g., FLAG or other epitope tags, etc., will allow detection of receptor. This can be histological, as an affinity method for biochemical

purification, or labeling or selection in an expression cloning approach. A two-hybrid selection system may also be applied making appropriate constructs with the available DTLR sequences. See, e.g., Fields and Song
5 (1989) Nature 340:245-246.

Generally, descriptions of DTLRs will be analogously applicable to individual specific embodiments directed to DTLR2, DTLR3, DTLR4, DTLR5, DTLR6, DTLR7, DTLR8, DTLR9, and/or DTLR10 reagents and compositions.

10 The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the inventions to the specific embodiments.

15 EXAMPLES

I. General Methods

Some of the standard methods are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor
20 Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and Supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York. Methods for
25 protein purification include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and
30 others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Coligan, et al. (ed. 1996) and periodic supplements, Current Protocols In Protein Science Greene/Wiley, New York; Deutscher (1990) "Guide to Protein Purification" in Methods in Enzymology, vol. 182,
35 and other volumes in this series; and manufacturer's literature on use of protein purification products, e.g., Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA.

Combination with recombinant techniques allow fusion to appropriate segments, e.g., to a FLAG sequence or an equivalent which can be fused via a protease-removable sequence. See, e.g., Hochuli (1989) Chemische Industrie 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering, Principle and Methods 12:87-98, Plenum Press, N.Y.; and Crowe, et al. (1992) OIAexpress: The High Level Expression & Protein Purification System QUIAGEN, Inc., Chatsworth, CA.

Standard immunological techniques and assays are described, e.g., in Hertzzenberg, et al. (eds. 1996) Weir's Handbook of Experimental Immunology vols. 1-4, Blackwell Science; Coligan (1991) Current Protocols in Immunology Wiley/Greene, NY; and Methods in Enzymology volumes. 70, 73, 74, 84, 92, 93, 108, 116, 121, 132, 150, 162, and 163.

Assays for vascular biological activities are well known in the art. They will cover angiogenic and angiostatic activities in tumor, or other tissues, e.g., arterial smooth muscle proliferation (see, e.g., Koyoma, et al. (1996) Cell 87:1069-1078), monocyte adhesion to vascular epithelium (see McEvoy, et al. (1997) J. Exp. Med. 185:2069-2077), etc. See also Ross (1993) Nature 362:801-809; Rekhater and Gordon (1995) Am. J. Pathol. 147:668-677; Thyberg, et al. (1990) Atherosclerosis 10:966-990; and Gumbiner (1996) Cell 84:345-357.

Assays for neural cell biological activities are described, e.g., in Wouterlood (ed. 1995) Neuroscience Protocols modules 10, Elsevier; Methods in Neurosciences Academic Press; and Neuromethods Humana Press, Totowa, NJ. Methodology of developmental systems is described, e.g., in Meisami (ed.) Handbook of Human Growth and Developmental Biology CRC Press; and Chrispeels (ed.) Molecular Techniques and Approaches in Developmental Biology Interscience.

Computer sequence analysis is performed, e.g., using available software programs, including those from the GCG (U. Wisconsin) and GenBank sources. Public sequence databases were also used, e.g., from GenBank, NCBI, EMBO, and others.

Many techniques applicable to IL-10 receptors may be applied to DTLRs, as described, e.g., in USSN 08/110,683 (IL-10 receptor), which is incorporated herein by reference for all purposes.

II. Novel Family of Human Receptors

Abbreviations: DTLR, Toll-like receptor; IL-1R, interleukin-1 receptor; TH, Toll homology; LRR, leucine-rich repeat; EST, expressed sequence tag; STS, sequence tagged site; FISH, fluorescence in situ hybridization.

The discovery of sequence homology between the cytoplasmic domains of *Drosophila* Toll and human interleukin-1 (IL-1) receptors has sown the conviction that both molecules trigger related signaling pathways tied to the nuclear translocation of Rel-type transcription factors. This conserved signaling scheme governs an evolutionarily ancient immune response in both insects and vertebrates. We report the molecular cloning of a novel class of putative human receptors with a protein architecture that is closely similar to *Drosophila* Toll in both intra- and extra-cellular segments. Five human Toll-like receptors, designated DTLRs 1-5, are likely the direct homologs of the fly molecule, and as such could constitute an important and unrecognized component of innate immunity in humans; intriguingly, the evolutionary retention of DTLRs in vertebrates may indicate another role, akin to Toll in the dorso-ventralization of the *Drosophila* embryo, as regulators of early morphogenetic patterning. Multiple tissue mRNA blots indicate markedly different patterns of

expression for the human DTLRs. Using fluorescence in situ hybridization and Sequence-Tagged Site database analyses, we also show that the cognate DTLR genes reside on chromosomes 4 (DTLRs 1, 2, and 3), 9 (DTLR4), and 1
5 (DTLR5). Structure prediction of the aligned Toll-homology (TH) domains from varied insect and human DTLRs, vertebrate IL-1 receptors, and MyD88 factors, and plant disease resistance proteins, recognizes a parallel β/α fold with an acidic active site; a similar structure
10 notably recurs in a class of response regulators broadly involved in transducing sensory information in bacteria.

The seeds of the morphogenetic gulf that so dramatically separates flies from humans are planted in
15 familiar embryonic shapes and patterns, but give rise to very different cell complexities. DeRobertis and Sasai (1996) Nature 380:37-40; and Arendt and Nübler-Jung (1997) Mech. Develop. 61:7-21. This divergence of developmental plans between insects and vertebrates is
20 choreographed by remarkably similar signaling pathways, underscoring a greater conservation of protein networks and biochemical mechanisms from unequal gene repertoires. Miklos and Rubin (1996) Cell 86:521-529; and Chothia (1994) Develop. 1994 Suppl., 27-33. A powerful way to
25 chart the evolutionary design of these regulatory pathways is by inferring their likely molecular components (and biological functions) through interspecies comparisons of protein sequences and structures. Miklos and Rubin (1996) Cell 86:521-529;
30 Chothia (1994) Develop. 1994 Suppl., 27-33 (3-5); and Banfi, et al. (1996) Nature Genet. 13:167-174.

A universally critical step in embryonic development is the specification of body axes, either born from innate asymmetries or triggered by external cues.
35 DeRobertis and Sasai (1996) Nature 380:37-40; and Arendt and Nübler-Jung (1997) Mech. Develop. 61:7-21. As a model system, particular attention has been focused on

the phylogenetic basis and cellular mechanisms of dorsoventral polarization . DeRobertis and Sasai (1996) Nature 380:37-40; and Arendt and Nübler-Jung (1997) Mech. Develop. 61:7-21. A prototype molecular strategy for
5 this transformation has emerged from the Drosophila embryo, where the sequential action of a small number of genes results in a ventralizing gradient of the transcription factor Dorsal. St. Johnston and Nüsslein-Volhard (1992) Cell 68:201-219; and Morisato and Anderson
10 (1995) Ann. Rev. Genet. 29:371-399.

This signaling pathway centers on Toll, a transmembrane receptor that transduces the binding of a maternally-secreted ventral factor, Spätzle, into the cytoplasmic engagement of Tube, an accessory molecule,
15 and the activation of Pelle, a Ser/Thr kinase that catalyzes the dissociation of Dorsal from the inhibitor Cactus and allows migration of Dorsal to ventral nuclei (Morisato and Anderson (1995) Ann. Rev. Genet. 29:371-399; and Belvin and Anderson (1996) Ann. Rev. Cell
20 Develop. Biol. 12:393-416. The Toll pathway also controls the induction of potent antimicrobial factors in the adult fly (Lemaitre, et al. (1996) Cell 86:973-983); this role in Drosophila immune defense strengthens mechanistic parallels to IL-1 pathways that govern a host
25 of immune and inflammatory responses in vertebrates. Belvin and Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416; and Wasserman (1993) Molec. Biol. Cell 4:767-771. A Toll-related cytoplasmic domain in IL-1 receptors directs the binding of a Pelle-like kinase, IRAK, and the
30 activation of a latent NF- κ B/I- κ B complex that mirrors the embrace of Dorsal and Cactus. Belvin and Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416; and Wasserman (1993) Molec. Biol. Cell 4:767-771.

We describe the cloning and molecular
35 characterization of four new Toll-like molecules in humans, designated DTLRs 2-5 (following Chiang & Beachy (1994) Mech. Develop. 47:225-239), that reveal a receptor

family more closely tied to *Drosophila* Toll homologs than to vertebrate IL-1 receptors. The DTLR sequences are derived from human ESTs; these partial cDNAs were used to draw complete expression profiles in human tissues for the five DTLRs, map the chromosomal locations of cognate genes, and narrow the choice of cDNA libraries for full-length cDNA retrievals. Spurred by other efforts (Banfi, et al. (1996) Nature Genet. 13:167-174; and Wang, et al. (1996) J. Biol. Chem. 271:4468-4476), we are assembling, by structural conservation and molecular parsimony, a biological system in humans that is the counterpart of a compelling regulatory scheme in *Drosophila*. In addition, a biochemical mechanism driving Toll signaling is suggested by the proposed tertiary fold of the Toll-homology (TH) domain, a core module shared by DTLRs, a broad family of IL-1 receptors, mammalian MyD88 factors and plant disease resistance proteins. Mitcham, et al. (1996) J. Biol. Chem. 271:5777-5783; and Hardiman, et al. (1996) Oncogene 13:2467-2475. We propose that a signaling route coupling morphogenesis and primitive immunity in insects, plants, and animals (Belvin and Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416; and Wilson, et al. (1997) Curr. Biol. 7:175-178) may have roots in bacterial two-component pathways.

25

Computational Analysis.

Human sequences related to insect DTLRs were identified from the EST database (dbEST) at the National Center for Biotechnology Information (NCBI) using the BLAST server (Altschul, et al. (1994) Nature Genet. 6:119-129). More sensitive pattern- and profile-based methods (Bork and Gibson (1996) Meth. Enzymol. 266:162-184) were used to isolate the signaling domains of the DTLR family that are shared with vertebrate and plant proteins present in nonredundant databases. The progressive alignment of DTLR intra- or extracellular domain sequences was carried out by ClustalW (Thompson,

et al. (1994) Nucleic Acids Res. 22:4673-4680); this program also calculated the branching order of aligned sequences by the Neighbor-Joining algorithm (5000 bootstrap replications provided confidence values for the tree groupings).

Conserved alignment patterns, discerned at several degrees of stringency, were drawn by the Consensus program (internet URL <http://www.bork.embl-heidelberg.de/Alignment/consensus.html>). The PRINTS library of protein fingerprints (<http://www.biochem.ucl.ac.uk/bsm/dbbrowser/PRINTS/PRINTS.html>) (Attwood, et al. (1997) Nucleic Acids Res. 25:212-217) reliably identified the myriad leucine-rich repeats (LRRs) present in the extracellular segments of DTLRs with a compound motif (PRINTS code Leurichrpt) that flexibly matches N- and C-terminal features of divergent LRRs. Two prediction algorithms whose three-state accuracy is above 72% were used to derive a consensus secondary structure for the intracellular domain alignment, as a bridge to fold recognition efforts (Fischer, et al. (1996) FASEB J. 10:126-136). Both the neural network program PHD (Rost and Sander (1994) Proteins 19:55-72) and the statistical prediction method DSC (King and Sternberg (1996) Protein Sci. 5:2298-2310) have internet servers (URLs http://www.embl-heidelberg.de/predictprotein/phd_pred.html and http://bonsai.lif.icnet.uk/bmm/dsc/dsc_read_align.html, respectively). The intracellular region encodes the THD region discussed, e.g., in Hardiman, et al. (1996) Oncogene 13:2467-2475; and Rock, et al. (1998) Proc. Nat'l Acad. Sci. USA 95:588-593, each of which is incorporated herein by reference. This domain is very important in the mechanism of signaling by the receptors, which transfers a phosphate group to a substrate.

35

Cloning of full-length human DTLR cDNAs.

PCR primers derived from the Toll-like Humrsc786 sequence (Genbank accession code D13637) (Nomura, et al. (1994) DNA Res 1:27-35) were used to probe a human erythroleukemic, TF-1 cell line-derived cDNA library (Kitamura, et al. (1989) Blood 73:375-380) to yield the DTLR1 cDNA sequence. The remaining DTLR sequences were flagged from dbEST, and the relevant EST clones obtained from the I.M.A.G.E. consortium (Lennon, et al. (1996) Genomics 33:151-152) via Research Genetics (Huntsville, AL): CloneID#'s 80633 and 117262 (DTLR2), 144675 (DTLR3), 202057 (DTLR4) and 277229 (DTLR5). Full length cDNAs for human DTLRs 2-4 were cloned by DNA hybridization screening of λ gt10 phage, human adult lung, placenta, and fetal liver 5'-Stretch Plus cDNA libraries (Clontech), respectively; the DTLR5 sequence is derived from a human multiple-sclerosis plaque EST. All positive clones were sequenced and aligned to identify individual DTLR ORFs: DTLR1 (2366 bp clone, 786 aa ORF), DTLR2 (2600 bp, 784 aa), DTLR3 (3029 bp, 904 aa), DTLR4 (3811 bp, 879 aa) and DTLR5 (1275 bp, 370 aa). Probes for DTLR3 and DTLR4 hybridizations were generated by PCR using human placenta (Stratagene) and adult liver (Clontech) cDNA libraries as templates, respectively; primer pairs were derived from the respective EST sequences. PCR reactions were conducted using T. aquaticus Tagplus DNA polymerase (Stratagene) under the following conditions: 1 x (94° C, 2 min) 30 x (55° C, 20 sec; 72° C 30 sec; 94° C 20 sec), 1 x (72° C, 8 min). For DTLR2 full-length cDNA screening, a 900 bp fragment generated by EcoRI/XbaI digestion of the first EST clone (ID# 80633) was used as a probe.

mRNA blots and chromosomal localization.

Human multiple tissue (Cat# 1, 2) and cancer cell line blots (Cat# 7757-1), containing approximately 2 μ g of poly(A)⁺ RNA per lane, were purchased from Clontech (Palo Alto, CA). For DTLRs 1-4, the isolated full-length

cDNAs served as probes, for DTLR5 the EST clone (ID #277229) plasmid insert was used. Briefly, the probes were radiolabeled with [α -³²P] dATP using the Amersham Rediprime random primer labeling kit (RPN1633).

- 5 Prehybridization and hybridizations were performed at 65° C in 0.5 M Na₂HPO₄, 7% SDS, 0.5 M EDTA (pH 8.0). All stringency washes were conducted at 65° C with two initial washes in 2 x SSC, 0.1% SDS for 40 min followed by a subsequent wash in 0.1 x SSC, 0.1% SDS for 20 min.
- 10 Membranes were then exposed at -70° C to X-Ray film (Kodak) in the presence of intensifying screens. More detailed studies by cDNA library Southern (14) were performed with selected human DTLR clones to examine their expression in hemopoietic cell subsets.
- 15 Human chromosomal mapping was conducted by the method of fluorescence in situ hybridization (FISH) as described in Heng and Tsui (1994) Meth. Molec. Biol. 33:109-122, using the various full-length (DTLRs 2-4) or partial (DTLR5) cDNA clones as probes. These analyses
- 20 were performed as a service by SeeDNA Biotech Inc. (Ontario, Canada). A search for human syndromes (or mouse defects in syntenic loci) associated with the mapped DTLR genes was conducted in the Dysmorphic Human-Mouse Homology Database by internet server
- 25 (http://www.hgmp.mrc.ac.uk/DHMH/hum_chrom1.html).

Conserved architecture of insect and human DTLR ectodomains.

- 30 The Toll family in *Drosophila* comprises at least four distinct gene products: Toll, the prototype receptor involved in dorsoventral patterning of the fly embryo (Morisato and Anderson (1995) Ann. Rev. Genet. 29:371-399) and a second named '18 Wheeler' (18w) that may also be involved in early embryonic development (Chiang and
- 35 Beachy (1994) Mech. Develop. 47:225-239; Eldon, et al. (1994) Develop. 120:885-899); two additional receptors are predicted by incomplete, Toll-like ORFs downstream of

the male-specific-transcript (Mst) locus (Genbank code X67703) or encoded by the 'sequence-tagged-site' (STS) Dm2245 (Genbank code G01378) (Mitcham, et al. (1996) J. Biol. Chem. 271:5777-5783). The extracellular segments
5 of Toll and 18w are distinctively composed of imperfect, ~24 amino acid LRR motifs (Chiang and Beachy (1994) Mech. Develop. 47:225-239; and Eldon, et al. (1994) Develop. 120:885-899). Similar tandem arrays of LRRs commonly form the adhesive antennae of varied cell surface
10 molecules and their generic tertiary structure is presumed to mimic the horseshoe-shaped cradle of a ribonuclease inhibitor fold, where seventeen LRRs show a repeating β/α -hairpin, 28 residue motif (Buchanan and Gay (1996) Prog. Biophys. Molec. Biol. 65:1-44). The
15 specific recognition of Spätzle by Toll may follow a model proposed for the binding of cystine-knot fold glycoprotein hormones by the multi-LRR ectodomains of serpentine receptors, using the concave side of the curved β -sheet (Kajava, et al. (1995) Structure 3:867-
20 877); intriguingly, the pattern of cysteines in Spätzle, and an orphan Drosophila ligand, Trunk, predict a similar cystine-knot tertiary structure (Belvin and Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416; and Casanova, et al. (1995) Genes Develop. 9:2539-2544).
25 The 22 and 31 LRR ectodomains of Toll and 18w, * respectively (the Mst ORF fragment displays 16 LRRs), are most closely related to the comparable 18, 19, 24, and 22 LRR arrays of DTLRs 1-4 (the incomplete DTLR5 chain presently includes four membrane-proximal LRRs) by
30 sequence and pattern analysis (Altschul, et al. (1994) Nature Genet. 6:119-129; and Bork and Gibson (1996) Meth. Enzymol. 266:162-184) (Fig. 1). However, a striking difference in the human DTLR chains is the common loss of
35 a ~90 residue cysteine-rich region that is variably embedded in the ectodomains of Toll, 18w and the Mst ORF (distanced four, six and two LRRs, respectively, from the membrane boundary). These cysteine clusters are

bipartite, with distinct 'top' (ending an LRR) and 'bottom' (stacked atop an LRR) halves (Chiang and Beachy (1994) Mech. Develop. 47:225-239; Eldon, et al. (1994) Develop. 120:885-899; and Buchanan and Gay (1996) Prog. Biophys. Molec. Biol. 65:1-44); the 'top' module recurs in both Drosophila and human DTLRs as a conserved juxtamembrane spacer (Fig. 1). We suggest that the flexibly located cysteine clusters in Drosophila receptors (and other LRR proteins), when mated 'top' to 'bottom', form a compact module with paired termini that can be inserted between any pair of LRRs without altering the overall fold of DTLR ectodomains; analogous 'extruded' domains decorate the structures of other proteins (Russell (1994) Protein Engin. 7:1407-1410).

15 Molecular design of the TH signaling domain.

Sequence comparison of Toll and IL-1 type-I (IL-1R1) receptors has disclosed a distant resemblance of a ~200 amino acid cytoplasmic domain that presumably mediates signaling by similar Rel-type transcription factors. Belvin and Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416; and (Belvin and Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416; and Wasserman (1993) Molec. Biol. Cell 4:767-771). More recent additions to this functional paradigm include a pair of plant disease resistance proteins from tobacco and flax that feature an N-terminal TH module followed by nucleotide-binding (NTPase) and LRR segments (Wilson, et al. (1997) Curr. Biol. 7:175-178); by contrast, a 'death domain' preceeds the TH chain of MyD88, an intracellular myeloid differentiation marker (Mitcham, et al. (1996) J. Biol. Chem. 271:5777-5783; and Hardiman, et al. (1996) Oncogene 13:2467-2475) (Fig. 1). New IL-1-type receptors include IL-1R3, an accessory signaling molecule, and orphan receptors IL-1R4 (also called ST2/Fit-1/T1), IL-1R5 (IL-1R-related protein), and IL-1R6 (IL-1R-related protein-2) (Mitcham, et al. (1996) J. Biol. Chem. 271:5777-

5783;Hardiman, et al. (1996) Oncogene 13:2467-2475).

With the new human DTLR sequences, we have sought a structural definition of this evolutionary thread by analyzing the conformation of the common TH module: ten
5 blocks of conserved sequence comprising 128 amino acids form the minimal TH domain fold; gaps in the alignment mark the likely location of sequence and length-variable loops (Fig. 2a).

Two prediction algorithms that take advantage of the
10 patterns of conservation and variation in multiply aligned sequences, PHD (Rost and Sander (1994) Proteins 19:55-72) and DSC (King and Sternberg (1996) Protein Sci. 5:2298-2310), produced strong, concordant results for the TH signaling module (Fig. 2a). Each block contains a
15 discrete secondary structural element: the imprint of alternating β -strands (labeled A-E) and α -helices (numbered 1-5) is diagnostic of an β/α -class fold with α -helices on both faces of a parallel β -sheet. Hydrophobic β -strands A, C and D are predicted to form 'interior'
20 staves in the β -sheet, while the shorter, amphipathic β -strands B and E resemble typical 'edge' units (Fig. 2a). This assignment is consistent with a strand order of B-A-C-D-E in the core β -sheet (Fig. 2b); fold comparison ('mapping') and recognition ('threading') programs
25 (Fischer, et al. (1996) FASEB J. 10:126-136) strongly return this doubly wound β/α topology. A surprising, functional prediction of this outline structure for the TH domain is that many of the conserved, charged residues in the multiple alignment map to the C-terminal end of
30 the β -sheet: residue Asp16 (block numbering scheme - Fig. 2a) at the end of β A, Arg39 and Asp40 following β B, Glu75 in the first turn of α 3, and the more loosely conserved Glu/Asp residues in the β D- α 4 loop, or after β E (Fig. 2a). The location of four other conserved residues
35 (Asp7, Glu28, and the Arg57-Arg/Lys58 pair) is compatible with a salt bridge network at the opposite, N-terminal end of the β -sheet (Fig. 2a).

Signaling function depends on the structural integrity of the TH domain. Inactivating mutations or deletions within the module boundaries (Fig. 2a) have been catalogued for IL-1R1 and Toll. Heguy, et al. (1992) J. Biol. Chem. 267:2605-2609; Croston, et al. (1995) J. Biol. Chem. 270:16514-16517; Schneider, et al. (1991) Genes Develop. 5:797-807; Norris and Manley (1992) Genes Develop. 6:1654-1667; Norris and Manley (1995) Genes Develop. 9:358-369; and Norris and Manley (1996) Genes Develop. 10:862-872. The human DTLR1-5 chains extending past the minimal TH domain (8, 0, 6, 22 and 18 residue lengths, respectively) are most closely similar to the stubby, 4 aa 'tail' of the Mst ORF. Toll and 18w display unrelated 102 and 207 residue tails (Fig. 2a) that may negatively regulate the signaling of the fused TH domains. Norris and Manley (1995) Genes Develop. 9:358-369; and Norris and Manley (1996) Genes Develop. 10:862-872.

The evolutionary relationship between the disparate proteins that carry the TH domain can best be discerned by a phylogenetic tree derived from the multiple alignment (Fig. 3). Four principal branches segregate the plant proteins, the MyD88 factors, IL-1 receptors and Toll-like molecules; the latter branch clusters the *Drosophila* and human DTLRs.

Chromosomal dispersal of human DTLR genes.

In order to investigate the genetic linkage of the nascent human DTLR gene family, we mapped the chromosomal loci of four of the five genes by FISH (Fig. 4). The DTLR1 gene has previously been charted by the human genome project: an STS database locus (dbSTS accession number G06709, corresponding to STS WI-7804 or SHGC-12827) exists for the Humrsc786 cDNA (Nomura, et al. (1994) DNA Res 1:27-35) and fixes the gene to chromosome 4 marker interval D4S1587-D42405 (50-56 cM) circa 4p14. This assignment has recently been corroborated by FISH

analysis. Taguchi, et al. (1996) Genomics 32:486-488.
In the present work, we reliably assign the remaining
DTLR genes to loci on chromosome 4q32 (DTLR2), 4q35
(DTLR3), 9q32-33 (DTLR4) and 1q33.3 (DTLR5). During the
5 course of this work, an STS for the parent DTLR2 EST
(cloneID # 80633) has been generated (dbSTS accession
number T57791 for STS SHGC-33147) and maps to the
chromosome 4 marker interval D4S424-D4S1548 (143-153 cM)
at 4q32 -in accord with our findings. There is a ~50 cM
10 gap between DTLR2 and DTLR3 genes on the long arm of
chromosome 4.

DTLR genes are differentially expressed.

Both Toll and 18w have complex spatial and temporal
15 patterns of expression in Drosophila that may point to
functions beyond embryonic patterning. St. Johnston and
Nüsslein-Volhard (1992) Cell 68:201-219; Morisato and
Anderson (1995) Ann. Rev. Genet. 29:371-399; Belvin and
Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416;
20 Lemaitre, et al. (1996) Cell 86:973-983; Chiang and
Beachy (1994) Mech. Develop. 47:225-239; and Eldon, et
al. (1994) Develop. 120:885-899. We have examined the
spatial distribution of DTLR transcripts by mRNA blot
analysis with varied human tissue and cancer cell lines
25 using radioabeled DTLR cDNAs (Fig. 5). DTLR1 is found to
be ubiquitously expressed, and at higher levels than the
other receptors. Presumably reflecting alternative
splicing, 'short' 3.0 kB and 'long' 8.0 kB DTLR1
transcript forms are present in ovary and spleen,
30 respectively (Fig. 5, panels A & B). A cancer cell mRNA
panel also shows the prominent overexpression of DTLR1 in
a Burkitt's Lymphoma Raji cell line (Fig. 5, panel C).
DTLR2 mRNA is less widely expressed than DTLR1, with a
4.0 kB species detected in lung and a 4.4 kB transcript
35 evident in heart, brain and muscle. The tissue
distribution pattern of DTLR3 echoes that of DTLR2 (Fig.
5, panel E). DTLR3 is also present as two major

transcripts of approximately 4.0 and 6.0 kB in size, and the highest levels of expression are observed in placenta and pancreas. By contrast, DTLR4 and DTLR5 messages appear to be extremely tissue-specific. DTLR4 was
5 detected only in placenta as a single transcript of ~7.0 kB in size. A faint 4.0 kB signal was observed for DTLR5 in ovary and peripheral blood monocytes.

10 Components of an evolutionarily ancient regulatory system.

The original molecular blueprints and divergent fates of signaling pathways can be reconstructed by comparative genomic approaches. Miklos and Rubin (1996) Cell 86:521-529; Chothia (1994) Develop. 1994 Suppl., 27-
15 33; Banfi, et al. (1996) Nature Genet. 13:167-174; and Wang, et al. (1996) J. Biol. Chem. 271:4468-4476. We have used this logic to identify an emergent gene family in humans, encoding five receptor paralogs at present, DTLRs 1-5, that are the direct evolutionary counterparts
20 of a *Drosophila* gene family headed by Toll (Figs. 1-3). The conserved architecture of human and fly DTLRs, conserved LRR ectodomains and intracellular TH modules (Fig. 1), intimates that the robust pathway coupled to Toll in *Drosophila* (6, 7) survives in vertebrates. The
25 best evidence borrows from a reiterated pathway: the manifold IL-1 system and its repertoire of receptor-fused TH domains, IRAK, NF- κ B and I- κ B homologs (Belvin and Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416; Wasserman (1993) Molec. Biol. Cell 4:767-771; Hardiman,
30 et al. (1996) Oncogene 13:2467-2475; and Cao, et al. (1996) Science 271:1128-1131); a Tube-like factor has also been characterized. It is not known whether DTLRs can productively couple to the IL-1R signaling machinery, or instead, a parallel set of proteins is used.
35 Differently from IL-1 receptors, the LRR cradle of human DTLRs is predicted to retain an affinity for Spätzle/Trunk-related cystine-knot factors; candidate

DTLR ligands (called PENs) that fit this mold have been isolated.

Biochemical mechanisms of signal transduction can be gauged by the conservation of interacting protein folds in a pathway. Miklos and Rubin (1996) Cell 86:521-529; Chothia (1994) Develop. 1994 Suppl., 27-33. At present, the Toll signaling paradigm involves some molecules whose roles are narrowly defined by their structures, actions or fates: Pelle is a Ser/Thr kinase (phosphorylation), Dorsal is an NF-KB-like transcription factor (DNA-binding) and Cactus is an ankyrin-repeat inhibitor (Dorsal binding, degradation). Belvin and Anderson (1996) Ann. Rev. Cell Develop. Biol. 12:393-416. By contrast, the functions of the Toll TH domain and Tube remain enigmatic. Like other cytokine receptors (Heldin (1995) Cell 80:213-223), ligand-mediated dimerization of Toll appears to be the triggering event: free cysteines in the juxtamembrane region of Toll create constitutively active receptor pairs (Schneider, et al. (1991) Genes Develop. 5:797-807), and chimeric Torso-Toll receptors signal as dimers (Galindo, et al. (1995) Develop. 121:2209-2218); yet, severe truncations or wholesale loss of the Toll ectodomain results in promiscuous intracellular signaling (Norris and Manley (1995) Genes Develop. 9:358-369; and Winans and Hashimoto (1995) Molec. Biol. Cell 6:587-596), reminiscent of oncogenic receptors with catalytic domains (Heldin (1995) Cell 80:213-223). Tube is membrane-localized, engages the N-terminal (death) domain of Pelle and is phosphorylated, but neither Toll-Tube or Toll-Pelle interactions are registered by two-hybrid analysis (Galindo, et al. (1995) Develop. 121:2209-2218; and Großhans, et al. (1994) Nature 372:563-566); this latter result suggests that the conformational 'state' of the Toll TH domain somehow affects factor recruitment. Norris and Manley (1996) Genes Develop. 10:862-872; and Galindo, et al. (1995) Develop. 121:2209-2218.

At the heart of these vexing issues is the structural nature of the Toll TH module. To address this question, we have taken advantage of the evolutionary diversity of TH sequences from insects, plants and
5 vertebrates, incorporating the human DTLR chains, and extracted the minimal, conserved protein core for structure prediction and fold recognition (Fig. 2). The strongly predicted (β/α)₅ TH domain fold with its asymmetric cluster of acidic residues is topologically
10 identical to the structures of response regulators in bacterial two-component signaling pathways (Volz (1993) Biochemistry 32:11741-11753; and Parkinson (1993) Cell 73:857-871) (Fig. 2). The prototype chemotaxis regulator CheY transiently binds a divalent cation in an 'aspartate
15 pocket' at the C-end of the core β -sheet; this cation provides electrostatic stability and facilitates the activating phosphorylation of an invariant Asp. Volz (1993) Biochemistry 32:11741-11753. Likewise, the TH domain may capture cations in its acidic nest, but
20 activation, and downstream signaling, could depend on the specific binding of a negatively charged moiety: anionic ligands can overcome intensely negative binding-site potentials by locking into precise hydrogen-bond networks. Ledvina, et al. (1996) Proc. Natl. Acad. Sci.
25 USA 93:6786-6791. Intriguingly, the TH domain may not simply act as a passive scaffold for the assembly of a Tube/Pelle complex for Toll, or homologous systems in plants and vertebrates, but instead actively participate as a true conformational trigger in the signal
30 transducing machinery. Perhaps explaining the conditional binding of a Tube/Pelle complex, Toll dimerization could promote unmasking, by regulatory receptor tails (Norris and Manley (1995) Genes Develop. 9:358-369; Norris and Manley (1996) Genes Develop.
35 10:862-872), or binding by small molecule activators of the TH pocket. However, 'free' TH modules inside the cell (Norris and Manley (1995) Genes Develop. 9:358-369;

Winans and Hashimoto (1995) Molec. Biol. Cell 6:587-596) could act as catalytic, CheY-like triggers by activating and docking with errant Tube/Pelle complexes.

5 Morphogenetic receptors and immune defense.

The evolutionary link between insect and vertebrate immune systems is stamped in DNA: genes encoding antimicrobial factors in insects display upstream motifs similar to acute phase response elements known to bind
10 NF- κ B transcription factors in mammals. Hultmark (1993) Trends Genet. 9:178-183. Dorsal, and two Dorsal-related factors, Dif and Relish, help induce these defense proteins after bacterial challenge (Reichhart, et al. (1993) C. R. Acad. Sci. Paris 316:1218-1224; Ip, et al.
15 (1993) Cell 75:753-763; and Dushay, et al. (1996) Proc. Natl. Acad. Sci. USA 93:10343-10347); Toll, or other DTLRs, likely modulate these rapid immune responses in adult *Drosophila* (Lemaitre, et al. (1996) Cell 86:973-983; and Rosetto, et al. (1995) Biochem. Biophys. Res.
20 Commun. 209:111-116). These mechanistic parallels to the IL-1 inflammatory response in vertebrates are evidence of the functional versatility of the Toll signaling pathway, and suggest an ancient synergy between embryonic patterning and innate immunity (Belvin and Anderson
25 (1996) Ann. Rev. Cell Develop. Biol. 12:393-416; Lemaitre, et al. (1996) Cell 86:973-983; Wasserman (1993) Molec. Biol. Cell 4:767-771; Wilson, et al. (1997) Curr. Biol. 7:175-178; Hultmark (1993) Trends Genet. 9:178-183; Reichhart, et al. (1993) C. R. Acad. Sci. Paris 316:1218-
30 1224; Ip, et al. (1993) Cell 75:753-763; Dushay, et al. (1996) Proc. Natl. Acad. Sci. USA 93:10343-10347; Rosetto, et al. (1995) Biochem. Biophys. Res. Commun. 209:111-116; Medzhitov and Janeway (1997) Curr. Opin. Immunol. 9:4-9; and Medzhitov and Janeway (1997) Curr.
35 Opin. Immunol. 9:4-9). The closer homology of insect and human DTLR proteins invites an even stronger overlap of biological functions that supersedes the purely immune

parallels to IL-1 systems, and lends potential molecular regulators to dorso-ventral and other transformations of vertebrate embryos. DeRobertis and Sasai (1996) Nature 380:37-40; and Arendt and Nübler-Jung (1997) Mech. Develop. 61:7-21.

The present description of an emergent, robust receptor family in humans mirrors the recent discovery of the vertebrate Frizzled receptors for Wnt patterning factors. Wang, et al. (1996) J. Biol. Chem. 271:4468-4476. As numerous other cytokine-receptor systems have roles in early development (Lemaire and Kodjabachian (1996) Trends Genet. 12:525-531), perhaps the distinct cellular contexts of compact embryos and gangly adults simply result in familiar signaling pathways and their diffusible triggers having different biological outcomes at different times, e.g., morphogenesis versus immune defense for DTLRs. For insect, plant, and human Toll-related systems (Hardiman, et al. (1996) Oncogene 13:2467-2475; Wilson, et al. (1997) Curr. Biol. 7:175-178), these signals course through a regulatory TH domain that intriguingly resembles a bacterial transducing engine (Parkinson (1993) Cell 73:857-871).

In particular, the DTLR6 exhibits structural features which establish its membership in the family. Moreover, members of the family have been implicated in a number of significant developmental disease conditions and with function of the innate immune system. In particular, the DTLR6 has been mapped to the X chromosome to a location which is a hot spot for major developmental abnormalities. See, e.g., The Sanger Center: human X chromosome website <http://www.sanger.ac.uk/HGP/ChrX/index.shtml>; and the Baylor College of Medicine Human Genome Sequencing website <http://gc.bcm.tmc.edu:8088/cgi-bin/seq/home>.

The accession number for the deposited PAC is AC003046. This accession number contains sequence from two PACs: RPC-164K3 and RPC-263P4. These two PAC

sequences mapped on human chromosome Xp22 at the Baylor web site between STS markers DXS704 and DXS7166. This region is a "hot spot" for severe developmental abnormalities.

5

III. Amplification of DTLR fragment by PCR

Two appropriate primer sequences are selected (see Tables 1 through 10). RT-PCR is used on an appropriate mRNA sample selected for the presence of message to
10 produce a partial or full length cDNA, e.g., a sample which expresses the gene. See, e.g., Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach and Dveksler (1995; eds.) PCR Primer: A Laboratory Manual
15 Cold Spring Harbor Press, CSH, NY. Such will allow determination of a useful sequence to probe for a full length gene in a cDNA library. The TLR6 is a contiguous sequence in the genome, which may suggest that the other TLRs are also. Thus, PCR on genomic DNA may yield full
20 length contiguous sequence, and chromosome walking methodology would then be applicable. Alternatively, sequence databases will contain sequence corresponding to portions of the described embodiments, or closely related forms, e.g., alternative splicing, etc. Expression
25 cloning techniques also may be applied on cDNA libraries.

IV. Tissue distribution of DTLRs

Message for each gene encoding these DTLRs has been detected. See Figures 5A-5F. Other cells and tissues
30 will be assayed by appropriate technology, e.g., PCR, immunoassay, hybridization, or otherwise. Tissue and organ cDNA preparations are available, e.g., from Clontech, Mountain View, CA. Identification of sources of natural expression are useful, as described.

35 Southern Analysis: DNA (5 µg) from a primary amplified cDNA library is digested with appropriate restriction enzymes to release the inserts, run on a 1% agarose gel and

transferred to a nylon membrane (Schleicher and Schuell, Keene, NH).

Samples for human mRNA isolation would typically include, e.g.: peripheral blood mononuclear cells
5 (monocytes, T cells, NK cells, granulocytes, B cells), resting (T100); peripheral blood mononuclear cells, activated with anti-CD3 for 2, 6, 12 h pooled (T101); T cell, TH0 clone Mot 72, resting (T102); T cell, TH0 clone Mot 72, activated with anti-CD28 and anti-CD3 for 3, 6,
10 12 h pooled (T103); T cell, TH0 clone Mot 72, anergic treated with specific peptide for 2, 7, 12 h pooled (T104); T cell, TH1 clone HY06, resting (T107); T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T108); T cell, TH1 clone HY06, anergic
15 treated with specific peptide for 2, 6, 12 h pooled (T109); T cell, TH2 clone HY935, resting (T110); T cell, TH2 clone HY935, activated with anti-CD28 and anti-CD3 for 2, 7, 12 h pooled (T111); T cells CD4+CD45RO- T cells polarized 27 days in anti-CD28, IL-4, and anti IFN- γ , TH2
20 polarized, activated with anti-CD3 and anti-CD28 4 h (T116); T cell tumor lines Jurkat and Hut78, resting (T117); T cell clones, pooled AD130.2, Tc783.12, Tc783.13, Tc783.58, Tc782.69, resting (T118); T cell random $\gamma\delta$ T cell clones, resting (T119); Splenocytes,
25 resting (B100); Splenocytes, activated with anti-CD40 and IL-4 (B101); B cell EBV lines pooled WT49, RSB, JY, CVIR, 721.221, RM3, HSY, resting (B102); B cell line JY, activated with PMA and ionomycin for 1, 6 h pooled (B103); NK 20 clones pooled, resting (K100); NK 20 clones
30 pooled, activated with PMA and ionomycin for 6 h (K101); NK1 clone, derived from peripheral blood of LGL leukemia patient, IL-2 treated (K106); NK cytotoxic clone 640-A30-1, resting (K107); hematopoietic precursor line TF1, activated with PMA and ionomycin for 1, 6 h pooled
35 (C100); U937 premonocytic line, resting (M100); U937 premonocytic line, activated with PMA and ionomycin for 1, 6 h pooled (M101); elutriated monocytes, activated

- with LPS, IFN γ , anti-IL-10 for 1, 2, 6, 12, 24 h pooled (M102); elutriated monocytes, activated with LPS, IFN γ , IL-10 for 1, 2, 6, 12, 24 h pooled (M103); elutriated monocytes, activated with LPS, IFN γ , anti-IL-10 for 4, 16 h pooled (M106); elutriated monocytes, activated with LPS, IFN γ , IL-10 for 4, 16 h pooled (M107); elutriated monocytes, activated LPS for 1 h (M108); elutriated monocytes, activated LPS for 6 h (M109); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, resting (D101); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, activated with PMA and ionomycin for 1 hr (D102); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, activated with PMA and ionomycin for 6 hr (D103); DC 95% CD1a+, from CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D104); DC 95% CD14+, ex CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin 1, 6 hr pooled (D105); DC CD1a+ CD86+, from CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D106); DC from monocytes GM-CSF, IL-4 5 days, resting (D107); DC from monocytes GM-CSF, IL-4 5 days, resting (D108); DC from monocytes GM-CSF, IL-4 5 days, activated LPS 4, 16 h pooled (D109); DC from monocytes GM-CSF, IL-4 5 days, activated TNF α , monocyte supe for 4, 16 h pooled (D110); leiomyoma L11 benign tumor (X101); normal myometrium M5 (O115); malignant leiomyosarcoma GS1 (X103); lung fibroblast sarcoma line MRC5, activated with PMA and ionomycin for 1, 6 h pooled (C101); kidney epithelial carcinoma cell line CHA, activated with PMA and ionomycin for 1, 6 h pooled (C102); kidney fetal 28 wk male (O100); lung fetal 28 wk male (O101); liver fetal 28 wk male (O102); heart fetal 28 wk male (O103); brain fetal 28 wk male (O104); gallbladder fetal 28 wk male (O106); small intestine fetal 28 wk male (O107); adipose tissue fetal 28 wk male (O108); ovary fetal 25 wk female (O109); uterus fetal 25 wk female (O110); testes fetal 28 wk male

(O111); spleen fetal 28 wk male (O112); adult placenta 28 wk (O113); and tonsil inflamed, from 12 year old (X100).

Samples for mouse mRNA isolation can include, e.g.:
resting mouse fibroblastic L cell line (C200); Braf:ER
5 (Braf fusion to estrogen receptor) transfected cells,
control (C201); T cells, TH1 polarized (Mel14 bright,
CD4+ cells from spleen, polarized for 7 days with IFN- γ
and anti IL-4; T200); T cells, TH2 polarized (Mel14
bright, CD4+ cells from spleen, polarized for 7 days with
10 IL-4 and anti-IFN- γ ; T201); T cells, highly TH1 polarized
(see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367;
activated with anti-CD3 for 2, 6, 16 h pooled; T202); T
cells, highly TH2 polarized (see Openshaw, et al. (1995)
J. Exp. Med. 182:1357-1367; activated with anti-CD3 for
15 2, 6, 16 h pooled; T203); CD44- CD25+ pre T cells, sorted
from thymus (T204); TH1 T cell clone D1.1, resting for 3
weeks after last stimulation with antigen (T205); TH1 T
cell clone D1.1, 10 μ g/ml ConA stimulated 15 h (T206);
TH2 T cell clone CDC35, resting for 3 weeks after last
20 stimulation with antigen (T207); TH2 T cell clone CDC35,
10 μ g/ml ConA stimulated 15 h (T208); Mel14+ naive T
cells from spleen, resting (T209); Mel14+ T cells,
polarized to Th1 with IFN- γ /IL-12/anti-IL-4 for 6, 12, 24
h pooled (T210); Mel14+ T cells, polarized to Th2 with
25 IL-4/anti-IFN- γ for 6, 13, 24 h pooled (T211);
unstimulated mature B cell leukemia cell line A20 (B200);
unstimulated B cell line CH12 (B201); unstimulated large
B cells from spleen (B202); B cells from total spleen,
LPS activated (B203); metrizamide enriched dendritic
30 cells from spleen, resting (D200); dendritic cells from
bone marrow, resting (D201); monocyte cell line RAW 264.7
activated with LPS 4 h (M200); bone-marrow macrophages
derived with GM and M-CSF (M201); macrophage cell line
J774, resting (M202); macrophage cell line J774 + LPS +
35 anti-IL-10 at 0.5, 1, 3, 6, 12 h pooled (M203);
macrophage cell line J774 + LPS + IL-10 at 0.5, 1, 3, 5,
12 h pooled (M204); aerosol challenged mouse lung tissue,

Th2 primers, aerosol OVA challenge 7, 14, 23 h pooled
(see Garlisi, et al. (1995) Clinical Immunology and
Immunopathology 75:75-83; X206); Nippostrongylus-infected
lung tissue (see Coffman, et al. (1989) Science 245:308-
5 310; X200); total adult lung, normal (O200); total lung,
rag-1 (see Schwarz, et al. (1993) Immunodeficiency 4:249-
252; O205); IL-10 K.O. spleen (see Kuhn, et al. (1991)
Cell 75:263-274; X201); total adult spleen, normal
(O201); total spleen, rag-1 (O207); IL-10 K.O. Peyer's
10 patches (O202); total Peyer's patches, normal (O210); IL-
10 K.O. mesenteric lymph nodes (X203); total mesenteric
lymph nodes, normal (O211); IL-10 K.O. colon (X203);
total colon, normal (O212); NOD mouse pancreas (see
Makino, et al. (1980) Jikken Dobutsu 29:1-13; X205);
15 total thymus, rag-1 (O208); total kidney, rag-1 (O209);
total heart, rag-1 (O202); total brain, rag-1 (O203);
total testes, rag-1 (O204); total liver, rag-1 (O206);
rat normal joint tissue (O300); and rat arthritic joint
tissue (X300).

20

V. Cloning of species counterparts of DTLRs

Various strategies are used to obtain species
counterparts of these DTLRs, preferably from other
25 primates. One method is by cross hybridization using
closely related species DNA probes. It may be useful to
go into evolutionarily similar species as intermediate
steps. Another method is by using specific PCR primers
based on the identification of blocks of similarity or
30 difference between particular species, e.g., human,
genes, e.g., areas of highly conserved or nonconserved
polypeptide or nucleotide sequence. Alternatively,
antibodies may be used for expression cloning.

35 VI. Production of mammalian DTLR protein

An appropriate, e.g., GST, fusion construct is
engineered for expression, e.g., in E. coli. For

example, a mouse IGIF pGex plasmid is constructed and transformed into *E. coli*. Freshly transformed cells are grown in LB medium containing 50 µg/ml ampicillin and induced with IPTG (Sigma, St. Louis, MO). After
5 overnight induction, the bacteria are harvested and the pellets containing the DTLR protein are isolated. The pellets are homogenized in TE buffer (50 mM Tris-base pH 8.0, 10 mM EDTA and 2 mM pefabloc) in 2 liters. This material is passed through a microfluidizer
10 (Microfluidics, Newton, MA) three times. The fluidized supernatant is spun down on a Sorvall GS-3 rotor for 1 h at 13,000 rpm. The resulting supernatant containing the DTLR protein is filtered and passed over a glutathione-SEPHAROSE column equilibrated in 50 mM Tris-base pH 8.0.
15 The fractions containing the DTLR-GST fusion protein are pooled and cleaved with thrombin (Enzyme Research Laboratories, Inc., South Bend, IN). The cleaved pool is then passed over a Q-SEPHAROSE column equilibrated in 50 mM Tris-base. Fractions containing DTLR are pooled and
20 diluted in cold distilled H₂O, to lower the conductivity, and passed back over a fresh Q-Sepharose column, alone or in succession with an immunoaffinity antibody column.. Fractions containing the DTLR protein are pooled, aliquoted, and stored in the -70° C freezer.
25 Comparision of the CD spectrum with DTLR1 protein may suggest that the protein is correctly folded. See Hazuda, et al. (1969) J. Biol. Chem. 264:1689-1693.

VII. Biological Assays with DTLRs

30 Biological assays will generally be directed to the ligand binding feature of the protein or to the kinase/phosphatase activity of the receptor. The activity will typically be reversible, as are many other enzyme actions. mediate phosphatase or phosphorylase
35 activities, which activities are easily measured by standard procedures. See, e.g., Hardie, et al. (eds. 1995) The Protein Kinase FactBook vols. I and II,

Academic Press, San Diego, CA; Hanks, et al. (1991) Meth. Enzymol. 200:38-62; Hunter, et al. (1992) Cell 70:375-388; Lewin (1990) Cell 61:743-752; Pines, et al. (1991) Cold Spring Harbor Symp. Quant. Biol. 56:449-463; and
5 Parker, et al. (1993) Nature 363:736-738.

The family of interleukins 1 contains molecules, each of which is an important mediator of inflammatory disease. For a comprehensive review, see Dinarello (1996) "Biologic basis for interleukin-1 in disease"
10 Blood 87:2095-2147. There are suggestions that the various Toll ligands may play important roles in the initiation of disease, particularly inflammatory responses. The finding of novel proteins related to the IL-1 family furthers the identification of molecules that
15 provide the molecular basis for initiation of disease and allow for the development of therapeutic strategies of increased range and efficacy.

VIII. Preparation of antibodies specific for, e.g.,
20 DTLR4

Inbred Balb/c mice are immunized intraperitoneally with recombinant forms of the protein, e.g., purified DTLR4 or stable transfected NIH-3T3 cells. Animals are boosted at appropriate time points with protein, with or
25 without additional adjuvant, to further stimulate antibody production. Serum is collected, or hybridomas produced with harvested spleens.

Alternatively, Balb/c mice are immunized with cells transformed with the gene or fragments thereof, either
30 endogenous or exogenous cells, or with isolated membranes enriched for expression of the antigen. Serum is collected at the appropriate time, typically after numerous further administrations. Various gene therapy techniques may be useful, e.g., in producing protein in
35 situ, for generating an immune response.

Monoclonal antibodies may be made. For example, splenocytes are fused with an appropriate fusion partner

and hybridomas are selected in growth medium by standard procedures. Hybridoma supernatants are screened for the presence of antibodies which bind to the desired DTLR, e.g., by ELISA or other assay. Antibodies which
5 specifically recognize specific DTLR embodiments may also be selected or prepared.

In another method, synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan
10 (1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press. In appropriate situations, the binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a
15 substrate for panning methods. Nucleic acids may also be introduced into cells in an animal to produce the antigen, which serves to elicit an immune response. See, e.g., Wang, et al. (1993) Proc. Nat'l. Acad. Sci. 90:4156-4160; Barry, et al. (1994) BioTechniques 16:616-
20 619; and Xiang, et al. (1995) Immunity 2: 129-135.

IX. Production of fusion proteins with, e.g., DTLR5

Various fusion constructs are made with DTLR5. This portion of the gene is fused to an epitope tag, e.g., a
25 FLAG tag, or to a two hybrid system construct. See, e.g., Fields and Song (1989) Nature 340:245-246.

The epitope tag may be used in an expression cloning procedure with detection with anti-FLAG antibodies to detect a binding partner, e.g., ligand for the respective
30 DTLR5. The two hybrid system may also be used to isolate proteins which specifically bind to DTLR5.

X. Chromosomal mapping of DTLRs

Chromosome spreads are prepared. In situ
35 hybridization is performed on chromosome preparations obtained from phytohemagglutinin-stimulated lymphocytes cultured for 72 h. 5-bromodeoxyuridine is added for the

final seven hours of culture (60 µg/ml of medium), to ensure a posthybridization chromosomal banding of good quality.

An appropriate fragment, e.g., a PCR fragment, amplified with the help of primers on total B cell cDNA template, is cloned into an appropriate vector. The vector is labeled by nick-translation with ³H. The radiolabeled probe is hybridized to metaphase spreads as described in Mattei, et al. (1985) Hum. Genet. 69:327-331.

After coating with nuclear track emulsion (KODAK NTB₂), slides are exposed, e.g., for 18 days at 4° C. To avoid any slipping of silver grains during the banding procedure, chromosome spreads are first stained with buffered Giemsa solution and metaphase photographed. R-banding is then performed by the fluorochrome-photolysis-Giemsa (FPG) method and metaphases rephotographed before analysis.

Alternatively, FISH can be performed, as described above. The DTLR genes are located on different chromosomes. DTLR2 and DTLR3 are localized to human chromosome 4; DTLR4 is localized to human chromosome 9, and DTLR5 is localized to human chromosome 1. See Figures 4A-4D.

25

XI. Structure activity relationship

Information on the criticality of particular residues is determined using standard procedures and analysis. Standard mutagenesis analysis is performed, e.g., by generating many different variants at determined positions, e.g., at the positions identified above, and evaluating biological activities of the variants. This may be performed to the extent of determining positions which modify activity, or to focus on specific positions to determine the residues which can be substituted to either retain, block, or modulate biological activity.

Alternatively, analysis of natural variants can indicate what positions tolerate natural mutations. This may result from populational analysis of variation among individuals, or across strains or species. Samples from
5 selected individuals are analysed, e.g., by PCR analysis and sequencing. This allows evaluation of population polymorphisms.

XI. Isolation of a ligand for a DTLR

10 A DTLR can be used as a specific binding reagent to identify its binding partner, by taking advantage of its specificity of binding, much like an antibody would be used. A binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to
15 a substrate for panning methods.

The binding composition is used to screen an expression library made from a cell line which expresses a binding partner, i.e., ligand, preferably membrane associated. Standard staining techniques are used to
20 detect or sort surface expressed ligand, or surface expressing transformed cells are screened by panning. Screening of intracellular expression is performed by various staining or immunofluorescence procedures. See also McMahan, et al. (1991) EMBO J. 10:2821-2832.

25 For example, on day 0, precoat 2-chamber permanox slides with 1 ml per chamber of fibronectin, 10 ng/ml in PBS, for 30 min at room temperature. Rinse once with PBS. Then plate COS cells at $2-3 \times 10^5$ cells per chamber in 1.5 ml of growth media. Incubate overnight at 37° C.

30 On day 1 for each sample, prepare 0.5 ml of a solution of 66 µg/ml DEAE-dextran, 66 µM chloroquine, and 4 µg DNA in serum free DME. For each set, a positive control is prepared, e.g., of DTLR-FLAG cDNA at 1 and 1/200 dilution, and a negative mock. Rinse cells with
35 serum free DME. Add the DNA solution and incubate 5 hr at 37° C. Remove the medium and add 0.5 ml 10% DMSO in

DME for 2.5 min. Remove and wash once with DME. Add 1.5 ml growth medium and incubate overnight.

On day 2, change the medium. On days 3 or 4, the cells are fixed and stained. Rinse the cells twice with
5 Hank's Buffered Saline Solution (HBSS) and fix in 4% paraformaldehyde (PFA)/glucose for 5 min. Wash 3X with HBSS. The slides may be stored at -80° C after all liquid is removed. For each chamber, 0.5 ml incubations are performed as follows. Add HBSS/saponin (0.1%) with
10 32 µl/ml of 1 M NaN₃ for 20 min. Cells are then washed with HBSS/saponin 1X. Add appropriate DTLR or DTLR/antibody complex to cells and incubate for 30 min. Wash cells twice with HBSS/saponin. If appropriate, add first antibody for 30 min. Add second antibody, e.g.,
15 Vector anti-mouse antibody, at 1/200 dilution, and incubate for 30 min. Prepare ELISA solution, e.g., Vector Elite ABC horseradish peroxidase solution, and preincubate for 30 min. Use, e.g., 1 drop of solution A (avidin) and 1 drop solution B (biotin) per 2.5 ml
20 HBSS/saponin. Wash cells twice with HBSS/saponin. Add ABC HRP solution and incubate for 30 min. Wash cells twice with HBSS, second wash for 2 min, which closes cells. Then add Vector diaminobenzoic acid (DAB) for 5 to 10 min. Use 2 drops of buffer plus 4 drops DAB plus 2
25 drops of H₂O₂ per 5 ml of glass distilled water.

Carefully remove chamber and rinse slide in water. Air dry for a few minutes, then add 1 drop of Crystal Mount and a cover slip. Bake for 5 min at 85-90° C.

Evaluate positive staining of pools and
30 progressively subclone to isolation of single genes responsible for the binding.

Alternatively, DTLR reagents are used to affinity purify or sort out cells expressing a putative ligand. See, e.g., Sambrook, et al. or Ausubel, et al.

35 Another strategy is to screen for a membrane bound receptor by panning. The receptor cDNA is constructed as described above. The ligand can be immobilized and used

to immobilize expressing cells. Immobilization may be achieved by use of appropriate antibodies which recognize, e.g., a FLAG sequence of a DTLR fusion construct, or by use of antibodies raised against the
5 first antibodies. Recursive cycles of selection and amplification lead to enrichment of appropriate clones and eventual isolation of receptor expressing clones.

Phage expression libraries can be screened by mammalian DTLRs. Appropriate label techniques, e.g.,
10 anti-FLAG antibodies, will allow specific labeling of appropriate clones.

All citations herein are incorporated herein by reference to the same extent as if each individual publication or patent
15 application was specifically and individually indicated to be incorporated by reference.

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be
20 apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled; and the invention is not to be limited by the
25 specific embodiments that have been presented herein by way of example.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5 (i) APPLICANT: (A) NAME: Schering Corporation
(B) STREET: 2000 Galloping Hill Road
(C) CITY: Kenilworth
(D) STATE: New Jersey
10 (E) COUNTRY: USA
(F) POSTAL CODE: 07033
(G) TELEPHONE: (908) 298-4000
(H) TELEFAX: (908) 298-5388

15 (ii) TITLE OF INVENTION: HUMAN RECEPTOR PROTEINS; RELATED
REAGENTS AND METHODS

(iii) NUMBER OF SEQUENCES: 35

20 (iv) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: Macintosh Power PC
(C) OPERATING SYSTEM: 8.0
25 (D) SOFTWARE: Microsoft Word 6.0

(v) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
30 (C) CLASSIFICATION:

(vi) PRIOR APPLICATION DATA:
(A) APPLICATION NO.: USSN 60/044,293
(B) FILING DATE: 07-MAY-1997

35 (A) APPLICATION NO.: USSN 60/072,212
(B) FILING DATE: 22-JAN-1998

(A) APPLICATION NO.: USSN 60/076,947
(B) FILING DATE: 05-MAR-1998

40 (2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2367 base pairs
45 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..2358

55 (ix) FEATURE:
(A) NAME/KEY: mat_peptide
(B) LOCATION: 67..2358

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

5	ATG ACT AGC ATC TTC CAT TTT GCC ATT ATC TTC ATG TTA ATA CTT CAG Met Thr Ser Ile Phe His Phe Ala Ile Ile Phe Met Leu Ile Leu Gln -22 -20 -15 -10	48
10	ATC AGA ATA CAA TTA TCT GAA GAA AGT GAA TTT TTA GTT GAT AGG TCA Ile Arg Ile Gln Leu Ser Glu Glu Ser Glu Phe Leu Val Asp Arg Ser -5 1 5 10.	96
15	AAA AAC GGT CTC ATC CAC GTT CCT AAA GAC CTA TCC CAG AAA ACA ACA Lys Asn Gly Leu Ile His Val Pro Lys Asp Leu Ser Gln Lys Thr Thr 15 20 25	144
20	ATC TTA AAT ATA TCG CAA AAT TAT ATA TCT GAG CTT TGG ACT TCT GAC Ile Leu Asn Ile Ser Gln Asn Tyr Ile Ser Glu Leu Trp Thr Ser Asp 30 35 40	192
25	ATC TTA TCA CTG TCA AAA CTG AGG ATT TTG ATA ATT TCT CAT AAT AGA Ile Leu Ser Leu Ser Lys Leu Arg Ile Leu Ile Ile Ser His Asn Arg 45 50 55	240
30	ATC CAG TAT CTT GAT ATC AGT GTT TTC AAA TTC AAC CAG GAA TTG GAA Ile Gln Tyr Leu Asp Ile Ser Val Phe Lys Phe Asn Gln Glu Leu Glu 60 65 70	288
35	TAC TTG GAT TTG TCC CAC AAC AAG TTG GTG AAG ATT TCT TGC CAC CCT Tyr Leu Asp Leu Ser His Asn Lys Leu Val Lys Ile Ser Cys His Pro 75 80 85 90	336
40	ACT GTG AAC CTC AAG CAC TTG GAC CTG TCA TTT AAT GCA TTT GAT GCC Thr Val Asn Leu Lys His Leu Asp Leu Ser Phe Asn Ala Phe Asp Ala 95 100 105	384
45	CTG CCT ATA TGC AAA GAG TTT GGC AAT ATG TCT CAA CTA AAA TTT CTG Leu Pro Ile Cys Lys Glu Phe Gly Asn Met Ser Gln Leu Lys Phe Leu 110 115 120	432
50	GGG TTG AGC ACC ACA CAC TTA GAA AAA TCT AGT GTG CTG CCA ATT GCT Gly Leu Ser Thr Thr His Leu Glu Lys Ser Ser Val Leu Pro Ile Ala 125 130 135	480
55	CAT TTG AAT ATC AGC AAG GTC TTG CTG GTC TTA GGA GAG ACT TAT GGG His Leu Asn Ile Ser Lys Val Leu Leu Val Leu Gly Glu Thr Tyr Gly 140 145 150	528
60	GAA AAA GAA GAC CCT GAG GGC CTT CAA GAC TTT AAC ACT GAG AGT CTG Glu Lys Glu Asp Pro Glu Gly Leu Gln Asp Phe Asn Thr Glu Ser Leu 155 160 165 170	576
65	CAC ATT GTG TTC CCC ACA AAC AAA GAA TTC CAT TTT ATT TTG GAT GTG His Ile Val Phe Pro Thr Asn Lys Glu Phe His Phe Ile Leu Asp Val 175 180 185	624
70	TCA GTC AAG ACT GTA GCA AAT CTG GAA CTA TCT AAT ATC AAA TGT GTG Ser Val Lys Thr Val Ala Asn Leu Glu Leu Ser Asn Ile Lys Cys Val 190 195 200	672
75	CTA GAA GAT AAC AAA TGT TCT TAC TTC CTA AGT ATT CTG GCG AAA CTT Leu Glu Asp Asn Lys Cys Ser Tyr Phe Leu Ser Ile Leu Ala Lys Leu	720

	205	210	215	
5	CAA ACA AAT CCA AAG TTA TCA AGT CTT ACC TTA AAC AAC ATT GAA ACA Gln Thr Asn Pro Lys Leu Ser Ser Leu Thr Leu Asn Asn Ile Glu Thr	768		
	220	225	230	
10	ACT TGG AAT TCT TTC ATT AGG ATC CTC CAA CTA GTT TGG CAT ACA ACT Thr Trp Asn Ser Phe Ile Arg Ile Leu Gln Leu Val Trp His Thr Thr	816		
	235	240	245	250
15	GTA TGG TAT TTC TCA ATT TCA AAC GTG AAG CTA CAG GGT CAG CTG GAC Val Trp Tyr Phe Ser Ile Ser Asn Val Lys Leu Gln Gly Gln Leu Asp	864		
	255	260	265	
20	TTC AGA GAT TTT GAT TAT TCT GGC ACT TCC TTG AAG GCC TTG TCT ATA Phe Arg Asp Phe Asp Tyr Ser Gly Thr Ser Leu Lys Ala Leu Ser Ile	912		
	270	275	280	
25	CAC CAA GTT GTC AGC GAT GTG TTC GGT TTT CCG CAA AGT TAT ATC TAT His Gln Val Val Ser Asp Val Phe Gly Phe Pro Gln Ser Tyr Ile Tyr	960		
	285	290	295	
30	GAA ATC TTT TCG AAT ATG AAC ATC AAA AAT TTC ACA GTG TCT GGT ACA Glu Ile Phe Ser Asn Met Asn Ile Lys Asn Phe Thr Val Ser Gly Thr	1008		
	300	305	310	
35	CGC ATG GTC CAC ATG CTT TGC CCA TCC AAA ATT AGC CCG TTC CTG CAT Arg Met Val His Met Leu Cys Pro Ser Lys Ile Ser Pro Phe Leu His	1056		
	315	320	325	330
40	TTG GAT TTT TCC AAT AAT CTC TTA ACA GAC ACG GTT TTT GAA AAT TGT Leu Asp Phe Ser Asn Asn Leu Leu Thr Asp Thr Val Phe Glu Asn Cys	1104		
	335	340	345	
45	GGG CAC CTT ACT GAG TTG GAG ACA CTT ATT TTA CAA ATG AAT CAA TTA Gly His Leu Thr Glu Leu Glu Thr Leu Ile Leu Gln Met Asn Gln Leu	1152		
	350	355	360	
50	AAA GAA CTT TCA AAA ATA GCT GAA ATG ACT ACA CAG ATG AAG TCT CTG Lys Glu Leu Ser Lys Ile Ala Glu Met Thr Thr Gln Met Lys Ser Leu	1200		
	365	370	375	
55	CAA CAA TTG GAT ATT AGC CAG AAT TCT GTA AGC TAT GAT GAA AAG AAA Gln Gln Leu Asp Ile Ser Gln Asn Ser Val Ser Tyr Asp Glu Lys Lys	1248		
	380	385	390	
60	GGA GAC TGT TCT TGG ACT AAA AGT TTA TTA AGT TTA AAT ATG TCT TCA Gly Asp Cys Ser Trp Thr Lys Ser Leu Leu Ser Leu Asn Met Ser Ser	1296		
	395	400	405	410
65	AAT ATA CTT ACT GAC ACT ATT TTC AGA TGT TTA CCT CCC AGG ATC AAG Asn Ile Leu Thr Asp Thr Ile Phe Arg Cys Leu Pro Pro Arg Ile Lys	1344		
	415	420	425	
70	GTA CTT GAT CTT CAC AGC AAT AAA ATA AAG AGC ATT CCT AAA CAA GTC Val Leu Asp Leu His Ser Asn Lys Ile Lys Ser Ile Pro Lys Gln Val	1392		
	430	435	440	
75	GTA AAA CTG GAA GCT TTG CAA GAA CTC AAT GTT GCT TTC AAT TCT TTA Val Lys Leu Glu Ala Leu Gln Glu Leu Asn Val Ala Phe Asn Ser Leu	1440		
	445	450	455	

5	ACT Thr	GAC Asp	CTT Leu	CCT Pro	GGA Gly	TGT Cys	GGC Gly	AGC Ser	TTT Phe	AGC Ser	AGC Ser	CTT Leu	TCT Ser	GTA Val	TTG Leu	ATC Ile	1488
	460 465 470																
10	ATT Ile	GAT Asp	CAC His	AAT Asn	TCA Ser	GTT Val	TCC Ser	CAC His	CCA Pro	TCA Ser	GCT Ala	GAT Asp	TTC Phe	TTC Phe	CAG Gln	AGC Ser	1536
	475 480 485 490																
15	TGC Cys	CAG Gln	AAG Lys	ATG Met	AGG Arg	TCA Ser	ATA Ile	AAA Lys	GCA Ala	GGG Gly	GAC Asp	AAT Asn	CCA Pro	TTC Phe	CAA Gln	TGT Cys	1584
	495 500 505																
20	ACC Thr	TGT Cys	GAG Glu	CTC Leu	GGA Gly	GAA Glu	TTT Phe	GTC Val	AAA Lys	AAT Asn	ATA Ile	GAC Asp	CAA Gln	GTA Val	TCA Ser	AGT Ser	1632
	510 515 520																
25	GAA Glu	GTG Val	TTA Leu	GAG Glu	GGC Gly	TGG Trp	CCT Pro	GAT Asp	TCT Ser	TAT Tyr	AAG Lys	TGT Cys	GAC Asp	TAC Tyr	CCG Pro	GAA Glu	1680
	525 530 535																
30	AGT Ser	TAT Tyr	AGA Arg	GGA Gly	ACC Thr	CTA Leu	CTA Leu	AAG Lys	GAC Asp	TTT Phe	CAC His	ATG Met	TCT Ser	GAA Glu	TTA Leu	TCC Ser	1728
	540 545 550																
35	TGC Cys	AAC Asn	ATA Ile	ACT Thr	CTG Leu	CTG Leu	ATC Ile	GTC Val	ACC Thr	ATC Ile	GTT Val	GCC Ala	ACC Thr	ATG Met	CTG Leu	GTG Val	1776
	555 560 565 570																
40	TTG Leu	GCT Ala	GTG Val	ACT Thr	GTG Val	ACC Thr	TCC Ser	CTC Leu	TGC Cys	ATC Ile	TAC Tyr	TTG Leu	GAT Asp	CTG Leu	CCC Pro	TGG Trp	1824
	575 580 585																
45	TAT Tyr	CTC Leu	AGG Arg	ATG Met	GTG Val	TGC Cys	CAG Gln	TGG Trp	ACC Thr	CAG Gln	ACC Thr	CGG Arg	CGC Arg	AGG Arg	GCC Ala	AGG Arg	1872
	590 595 600																
50	AAC Asn	ATA Ile	CCC Pro	TTA Leu	GAA Glu	GAA Glu	CTC Leu	CAA Gln	AGA Arg	AAT Asn	CTC Leu	CAG Gln	TTT Phe	CAT His	GCA Ala	TTT Phe	1920
	605 610 615																
55	ATT Ile	TCA Ser	TAT Tyr	AGT Ser	GGG Gly	CAC His	GAT Asp	TCT Ser	TTC Phe	TGG Trp	GTG Val	AAG Lys	AAT Asn	GAA Glu	TTA Leu	TTG Leu	1968
	620 625 630																
60	CCA Pro	AAC Asn	CTA Leu	GAG Glu	AAA Lys	GAA Glu	GGT Gly	ATG Met	CAG Gln	ATT Ile	TGC Cys	CTT Leu	CAT His	GAG Glu	AGA Arg	AAC Asn	2016
	635 640 645 650																
65	TTT Phe	GTT Val	CCT Pro	GGC Gly	AAG Lys	AGC Ser	ATT Ile	GTG Val	GAA Glu	AAT Asn	ATC Ile	ATC Ile	ACC Thr	TGC Cys	ATT Ile	GAG Glu	2064
	655 660 665																
70	AAG Lys	AGT Ser	TAC Tyr	AAG Lys	TCC Ser	ATC Ile	TTT Phe	GTT Val	TTG Leu	TCT Ser	CCC Pro	AAC Asn	TTT Phe	GTC Val	CAG Gln	AGT Ser	2112
	670 675 680																
75	GAA Glu	TGG Trp	TGC Cys	CAT His	TAT Tyr	GAA Glu	CTC Leu	TAC Tyr	TTT Phe	GCC Ala	CAT His	CAC His	AAT Asn	CTC Leu	TTT Phe	CAT His	2160
	685 690 695																

	GAA GGA TCT AAT AGC TTA ATC CTG ATC TTG CTG GAA CCC ATT CCG CAG	2208
	Glu Gly Ser Asn Ser Leu Ile Leu Ile Leu Leu Glu Pro Ile Pro Gln	
	700 705 710	
5	TAC TCC ATT CCT AGC AGT TAT CAC AAG CTC AAA AGT CTC ATG GCC AGG	2256
	Tyr Ser Ile Pro Ser Ser Tyr His Lys Leu Lys Ser Leu Met Ala Arg	
	715 720 725 730	
10	AGG ACT TAT TTG GAA TGG CCC AAG GAA AAG AGC AAA CGT GGC CTT TTT	2304
	Arg Thr Tyr Leu Glu Trp Pro Lys Glu Lys Ser Lys Arg Gly Leu Phe	
	735 740 745	
	TGG GCT AAC TTA AGG GCA GCC ATT AAT ATT AAG CTG ACA GAG CAA GCA	2352
15	Trp Ala Asn Leu Arg Ala Ala Ile Asn Ile Lys Leu Thr Glu Gln Ala	
	750 755 760	
	AAG AAA TAGTCTAGA	2367
	Lys Lys	
20		
	(2) INFORMATION FOR SEQ ID NO:2:	
	(i) SEQUENCE CHARACTERISTICS:	
25	(A) LENGTH: 786 amino acids	
	(B) TYPE: amino acid	
	(D) TOPOLOGY: linear	
	(ii) MOLECULE TYPE: protein	
30	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:	
	Met Thr Ser Ile Phe His Phe Ala Ile Ile Phe Met Leu Ile Leu Gln	
	-22 -20 -15 -10	
35	Ile Arg Ile Gln Leu Ser Glu Glu Ser Glu Phe Leu Val Asp Arg Ser	
	-5 1 5 10	
40	Lys Asn Gly Leu Ile His Val Pro Lys Asp Leu Ser Gln Lys Thr Thr	
	15 20 25	
	Ile Leu Asn Ile Ser Gln Asn Tyr Ile Ser Glu Leu Trp Thr Ser Asp	
	30 35 40	
45	Ile Leu Ser Leu Ser Lys Leu Arg Ile Leu Ile Ile Ser His Asn Arg	
	45 50 55	
	Ile Gln Tyr Leu Asp Ile Ser Val Phe Lys Phe Asn Gln Glu Leu Glu	
	60 65 70	
50	Tyr Leu Asp Leu Ser His Asn Lys Leu Val Lys Ile Ser Cys His Pro	
	75 80 85 90	
55	Thr Val Asn Leu Lys His Leu Asp Leu Ser Phe Asn Ala Phe Asp Ala	
	95 100 105	
	Leu Pro Ile Cys Lys Glu Phe Gly Asn Met Ser Gln Leu Lys Phe Leu	
	110 115 120	
60	Gly Leu Ser Thr Thr His Leu Glu Lys Ser Ser Val Leu Pro Ile Ala	
	125 130 135	

His Leu Asn Ile Ser Lys Val Leu Leu Val Leu Gly Glu Thr Tyr Gly
 140 145 150
 5 Glu Lys Glu Asp Pro Glu Gly Leu Gln Asp Phe Asn Thr Glu Ser Leu
 155 160 165 170
 His Ile Val Phe Pro Thr Asn Lys Glu Phe His Phe Ile Leu Asp Val
 175 180 185
 10 Ser Val Lys Thr Val Ala Asn Leu Glu Leu Ser Asn Ile Lys Cys Val
 190 195 200
 Leu Glu Asp Asn Lys Cys Ser Tyr Phe Leu Ser Ile Leu Ala Lys Leu
 205 210 215
 15 Gln Thr Asn Pro Lys Leu Ser Ser Leu Thr Leu Asn Asn Ile Glu Thr
 220 225 230
 20 Thr Trp Asn Ser Phe Ile Arg Ile Leu Gln Leu Val Trp His Thr Thr
 235 240 245 250
 Val Trp Tyr Phe Ser Ile Ser Asn Val Lys Leu Gln Gly Gln Leu Asp
 255 260 265
 25 Phe Arg Asp Phe Asp Tyr Ser Gly Thr Ser Leu Lys Ala Leu Ser Ile
 270 275 280
 His Gln Val Val Ser Asp Val Phe Gly Phe Pro Gln Ser Tyr Ile Tyr
 285 290 295
 30 Glu Ile Phe Ser Asn Met Asn Ile Lys Asn Phe Thr Val Ser Gly Thr
 300 305 310
 35 Arg Met Val His Met Leu Cys Pro Ser Lys Ile Ser Pro Phe Leu His
 315 320 325 330
 Leu Asp Phe Ser Asn Asn Leu Leu Thr Asp Thr Val Phe Glu Asn Cys
 335 340 345
 40 Gly His Leu Thr Glu Leu Glu Thr Leu Ile Leu Gln Met Asn Gln Leu
 350 355 360
 Lys Glu Leu Ser Lys Ile Ala Glu Met Thr Thr Gln Met Lys Ser Leu
 365 370 375
 45 Gln Gln Leu Asp Ile Ser Gln Asn Ser Val Ser Tyr Asp Glu Lys Lys
 380 385 390
 50 Gly Asp Cys Ser Trp Thr Lys Ser Leu Leu Ser Leu Asn Met Ser Ser
 395 400 405 410
 Asn Ile Leu Thr Asp Thr Ile Phe Arg Cys Leu Pro Pro Arg Ile Lys
 415 420 425
 55 Val Leu Asp Leu His Ser Asn Lys Ile Lys Ser Ile Pro Lys Gln Val
 430 435 440
 60 Val Lys Leu Glu Ala Leu Gln Glu Leu Asn Val Ala Phe Asn Ser Leu
 445 450 455

Thr Asp Leu Pro Gly Cys Gly Ser Phe Ser Ser Leu Ser Val Leu Ile
 460 465 470
 5 Ile Asp His Asn Ser Val Ser His Pro Ser Ala Asp Phe Phe Gln Ser
 475 480 485 490
 Cys Gln Lys Met Arg Ser Ile Lys Ala Gly Asp Asn Pro Phe Gln Cys
 495 500 505
 10 Thr Cys Glu Leu Gly Glu Phe Val Lys Asn Ile Asp Gln Val Ser Ser.
 510 515 520
 Glu Val Leu Glu Gly Trp Pro Asp Ser Tyr Lys Cys Asp Tyr Pro Glu
 525 530 535
 15 Ser Tyr Arg Gly Thr Leu Leu Lys Asp Phe His Met Ser Glu Leu Ser
 540 545 550
 Cys Asn Ile Thr Leu Leu Ile Val Thr Ile Val Ala Thr Met Leu Val
 555 560 565 570
 Leu Ala Val Thr Val Thr Ser Leu Cys Ile Tyr Leu Asp Leu Pro Trp
 575 580 585
 25 Tyr Leu Arg Met Val Cys Gln Trp Thr Gln Thr Arg Arg Arg Ala Arg
 590 595 600
 Asn Ile Pro Leu Glu Glu Leu Gln Arg Asn Leu Gln Phe His Ala Phe
 605 610 615
 30 Ile Ser Tyr Ser Gly His Asp Ser Phe Trp Val Lys Asn Glu Leu Leu
 620 625 630
 Pro Asn Leu Glu Lys Glu Gly Met Gln Ile Cys Leu His Glu Arg Asn
 635 640 645 650
 Phe Val Pro Gly Lys Ser Ile Val Glu Asn Ile Ile Thr Cys Ile Glu
 655 660 665
 40 Lys Ser Tyr Lys Ser Ile Phe Val Leu Ser Pro Asn Phe Val Gln Ser
 670 675 680
 Glu Trp Cys His Tyr Glu Leu Tyr Phe Ala His His Asn Leu Phe His
 685 690 695
 45 Glu Gly Ser Asn Ser Leu Ile Leu Ile Leu Leu Glu Pro Ile Pro Gln
 700 705 710
 Tyr Ser Ile Pro Ser Ser Tyr His Lys Leu Lys Ser Leu Met Ala Arg
 715 720 725 730
 Arg Thr Tyr Leu Glu Trp Pro Lys Glu Lys Ser Lys Arg Gly Leu Phe
 735 740 745
 55 Trp Ala Asn Leu Arg Ala Ala Ile Asn Ile Lys Leu Thr Glu Gln Ala
 750 755 760
 Lys Lys

60

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2355 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..2352

(ix) FEATURE:
 (A) NAME/KEY: mat_peptide
 (B) LOCATION: 67..2352

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATG	CCA	CAT	ACT	TTG	TGG	ATG	GTG	TGG	GTC	TTG	GGG	GTC	ATC	ATC	AGC	48
Met	Pro	His	Thr	Leu	Trp	Met	Val	Trp	Val	Leu	Gly	Val	Ile	Ile	Ser	
-22		-20				-15					-10					
CTC	TCC	AAG	GAA	GAA	TCC	TCC	AAT	CAG	GCT	TCT	CTG	TCT	TGT	GAC	CGC	96
Leu	Ser	Lys	Glu	Glu	Ser	Ser	Asn	Gln	Ala	Ser	Leu	Ser	Cys	Asp	Arg	
-5					1					5					10	
AAT	GGT	ATC	TGC	AAG	GGC	AGC	TCA	GGA	TCT	TTA	AAC	TCC	ATT	CCC	TCA	144
Asn	Gly	Ile	Cys	Lys	Gly	Ser	Ser	Gly	Ser	Leu	Asn	Ser	Ile	Pro	Ser	
				15				20						25		
GGG	CTC	ACA	GAA	GCT	GTA	AAA	AGC	CTT	GAC	CTG	TCC	AAC	AAC	AGG	ATC	192
Gly	Leu	Thr	Glu	Ala	Val	Lys	Ser	Leu	Asp	Leu	Ser	Asn	Asn	Arg	Ile	
			30					35						40		
ACC	TAC	ATT	AGC	AAC	AGT	GAC	CTA	CAG	AGG	TGT	GTG	AAC	CTC	CAG	GCT	240
Thr	Tyr	Ile	Ser	Asn	Ser	Asp	Leu	Gln	Arg	Cys	Val	Asn	Leu	Gln	Ala	
		45				50						55				
CTG	GTG	CTG	ACA	TCC	AAT	GGA	ATT	AAC	ACA	ATA	GAG	GAA	GAT	TCT	TTT	288
Leu	Val	Leu	Thr	Ser	Asn	Gly	Ile	Asn	Thr	Ile	Glu	Glu	Asp	Ser	Phe	
	60					65					70					
TCT	TCC	CTG	GGC	AGT	CTT	GAA	CAT	TTA	GAC	TTA	TCC	TAT	AAT	TAC	TTA	336
Ser	Ser	Leu	Gly	Ser	Leu	Glu	His	Leu	Asp	Leu	Ser	Tyr	Asn	Tyr	Leu	
	75				80					85					90	
TCT	AAT	TTA	TCG	TCT	TCC	TGG	TTC	AAG	CCC	CTT	TCT	TCT	TTA	ACA	TTC	384
Ser	Asn	Leu	Ser	Ser	Ser	Trp	Phe	Lys	Pro	Leu	Ser	Ser	Leu	Thr	Phe	
				95					100					105		
TTA	AAC	TTA	CTG	GGA	AAT	CCT	TAC	AAA	ACC	CTA	GGG	GAA	ACA	TCT	CTT	432
Leu	Asn	Leu	Leu	Gly	Asn	Pro	Tyr	Lys	Thr	Leu	Gly	Glu	Thr	Ser	Leu	
			110					115					120			
TTT	TCT	CAT	CTC	ACA	AAA	TTG	CAA	ATC	CTG	AGA	GTG	GGA	AAT	ATG	GAC	480
Phe	Ser	His	Leu	Thr	Lys	Leu	Gln	Ile	Leu	Arg	Val	Gly	Asn	Met	Asp	
		125					130					135				

	ACC	TTC	ACT	AAG	ATT	CAA	AGA	AAA	GAT	TTT	GCT	GGA	CTT	ACC	TTC	CTT	528
	Thr	Phe	Thr	Lys	Ile	Gln	Arg	Lys	Asp	Phe	Ala	Gly	Leu	Thr	Phe	Leu	
	140					145					150						
5	GAG	GAA	CTT	GAG	ATT	GAT	GCT	TCA	GAT	CTA	CAG	AGC	TAT	GAG	CCA	AAA	576
	Glu	Glu	Leu	Glu	Ile	Asp	Ala	Ser	Asp	Leu	Gln	Ser	Tyr	Glu	Pro	Lys	
	155				160					165						170	
10	AGT	TTG	AAG	TCA	ATT	CAG	AAC	GTA	AGT	CAT	CTG	ATC	CTT	CAT	ATG	AAG	624
	Ser	Leu	Lys	Ser	Ile	Gln	Asn	Val	Ser	His	Leu	Ile	Leu	His	Met	Lys	
					175				180						185		
15	CAG	CAT	ATT	TTA	CTG	CTG	GAG	ATT	TTT	GTA	GAT	GTT	ACA	AGT	TCC	GTG	672
	Gln	His	Ile	Leu	Leu	Leu	Glu	Ile	Phe	Val	Asp	Val	Thr	Ser	Ser	Val	
				190					195					200			
20	GAA	TGT	TTG	GAA	CTG	CGA	GAT	ACT	GAT	TTG	GAC	ACT	TTC	CAT	TTT	TCA	720
	Glu	Cys	Leu	Glu	Leu	Arg	Asp	Thr	Asp	Leu	Asp	Thr	Phe	His	Phe	Ser	
			205					210					215				
25	GAA	CTA	TCC	ACT	GGT	GAA	ACA	AAT	TCA	TTG	ATT	AAA	AAG	TTT	ACA	TTT	768
	Glu	Leu	Ser	Thr	Gly	Glu	Thr	Asn	Ser	Leu	Ile	Lys	Lys	Phe	Thr	Phe	
		220					225					230					
30	AGA	AAT	GTG	AAA	ATC	ACC	GAT	GAA	AGT	TTG	TTT	CAG	GTT	ATG	AAA	CTT	816
	Arg	Asn	Val	Lys	Ile	Thr	Asp	Glu	Ser	Leu	Phe	Gln	Val	Met	Lys	Leu	
		235				240						245				250	
35	TTG	AAT	CAG	ATT	TCT	GGA	TTG	TTA	GAA	TTA	GAG	TTT	GAT	GAC	TGT	ACC	864
	Leu	Asn	Gln	Ile	Ser	Gly	Leu	Leu	Glu	Leu	Glu	Phe	Asp	Asp	Cys	Thr	
				255						260					265		
40	CTT	AAT	GGA	GTT	GGT	AAT	TTT	AGA	GCA	TCT	GAT	AAT	GAC	AGA	GTT	ATA	912
	Leu	Asn	Gly	Val	Gly	Asn	Phe	Arg	Ala	Ser	Asp	Asn	Asp	Arg	Val	Ile	
				270					275					280			
45	GAT	CCA	GGT	AAA	GTG	GAA	ACG	TTA	ACA	ATC	CGG	AGG	CTG	CAT	ATT	CCA	960
	Asp	Pro	Gly	Lys	Val	Glu	Thr	Leu	Thr	Ile	Arg	Arg	Leu	His	Ile	Pro	
			285					290					295				
50	AGG	TTT	TAC	TTA	TTT	TAT	GAT	CTG	AGC	ACT	TTA	TAT	TCA	CTT	ACA	GAA	1008
	Arg	Phe	Tyr	Leu	Phe	Tyr	Asp	Leu	Ser	Thr	Leu	Tyr	Ser	Leu	Thr	Glu	
		300					305					310					
55	AGA	GTT	AAA	AGA	ATC	ACA	GTA	GAA	AAC	AGT	AAA	GTT	TTT	CTG	GTT	CCT	1056
	Arg	Val	Lys	Arg	Ile	Thr	Val	Glu	Asn	Ser	Lys	Val	Phe	Leu	Val	Pro	
		315				320					325					330	
60	TGT	TTA	CTT	TCA	CAA	CAT	TTA	AAA	TCA	TTA	GAA	TAC	TTG	GAT	CTC	AGT	1104
	Cys	Leu	Leu	Ser	Gln	His	Leu	Lys	Ser	Leu	Glu	Tyr	Leu	Asp	Leu	Ser	
					335					340					345		
65	GAA	AAT	TTG	ATG	GTT	GAA	GAA	TAC	TTG	AAA	AAT	TCA	GCC	TGT	GAG	GAT	1152
	Glu	Asn	Leu	Met	Val	Glu	Glu	Tyr	Leu	Lys	Asn	Ser	Ala	Cys	Glu	Asp	
			350					355					360				
70	GCC	TGG	CCC	TCT	CTA	CAA	ACT	TTA	ATT	TTA	AGG	CAA	AAT	CAT	TTG	GCA	1200
	Ala	Trp	Pro	Ser	Leu	Gln	Thr	Leu	Ile	Leu	Arg	Gln	Asn	His	Leu	Ala	
			365					370					375				
75	TCA	TTG	GAA	AAA	ACC	GGA	GAG	ACT	TTG	CTC	ACT	CTG	AAA	AAC	TTG	ACT	1248

	Ser	Leu	Glu	Lys	Thr	Gly	Glu	Thr	Leu	Leu	Thr	Leu	Lys	Asn	Leu	Thr	
	380						385					390					
5	AAC	ATT	GAT	ATC	AGT	AAG	AAT	AGT	TTT	CAT	TCT	ATG	CCT	GAA	ACT	TGT	1296
	Asn	Ile	Asp	Ile	Ser	Lys	Asn	Ser	Phe	His	Ser	Met	Pro	Glu	Thr	Cys	
	395					400					405					410	
10	CAG	TGG	CCA	GAA	AAG	ATG	AAA	TAT	TTG	AAC	TTA	TCC	AGC	ACA	CGA	ATA	1344
	Gln	Trp	Pro	Glu	Lys	Met	Lys	Tyr	Leu	Asn	Leu	Ser	Ser	Thr	Arg	Ile	
					415					420					425		
15	CAC	AGT	GTA	ACA	GGC	TGC	ATT	CCC	AAG	ACA	CTG	GAA	ATT	TTA	GAT	GTT	1392
	His	Ser	Val	Thr	Gly	Cys	Ile	Pro	Lys	Thr	Leu	Glu	Ile	Leu	Asp	Val	
				430				435						440			
20	AGC	AAC	AAC	AAT	CTC	AAT	TTA	TTT	TCT	TTG	AAT	TTG	CCG	CAA	CTC	AAA	1440
	Ser	Asn	Asn	Asn	Leu	Asn	Leu	Phe	Ser	Leu	Asn	Leu	Pro	Gln	Leu	Lys	
				445				450					455				
25	GAA	CTT	TAT	ATT	TCC	AGA	AAT	AAG	TTG	ATG	ACT	CTA	CCA	GAT	GCC	TCC	1488
	Glu	Leu	Tyr	Ile	Ser	Arg	Asn	Lys	Leu	Met	Thr	Leu	Pro	Asp	Ala	Ser	
		460					465					470					
30	CTC	TTA	CCC	ATG	TTA	CTA	GTA	TTG	AAA	ATC	AGT	AGG	AAT	GCA	ATA	ACT	1536
	Leu	Leu	Pro	Met	Leu	Leu	Val	Leu	Lys	Ile	Ser	Arg	Asn	Ala	Ile	Thr	
	475					480					485					490	
35	ACG	TTT	TCT	AAG	GAG	CAA	CTT	GAC	TCA	TTT	CAC	ACA	CTG	AAG	ACT	TTG	1584
	Thr	Phe	Ser	Lys	Glu	Gln	Leu	Asp	Ser	Phe	His	Thr	Leu	Lys	Thr	Leu	
				495						500					505		
40	GAA	GCT	GGT	GGC	AAT	AAC	TTC	ATT	TGC	TCC	TGT	GAA	TTC	CTC	TCC	TTC	1632
	Glu	Ala	Gly	Gly	Asn	Asn	Phe	Ile	Cys	Ser	Cys	Glu	Phe	Leu	Ser	Phe	
				510					515					520			
45	ACT	CAG	GAG	CAG	CAA	GCA	CTG	GCC	AAA	GTC	TTG	ATT	GAT	TGG	CCA	GCA	1680
	Thr	Gln	Glu	Gln	Gln	Ala	Leu	Ala	Lys	Val	Leu	Ile	Asp	Trp	Pro	Ala	
				525				530					535				
50	AAT	TAC	CTG	TGT	GAC	TCT	CCA	TCC	CAT	GTG	CGT	GGC	CAG	CAG	GTT	CAG	1728
	Asn	Tyr	Leu	Cys	Asp	Ser	Pro	Ser	His	Val	Arg	Gly	Gln	Gln	Val	Gln	
		540					545					550					
55	GAT	GTC	CGC	CTC	TCG	GTG	TCG	GAA	TGT	CAC	AGG	ACA	GCA	CTG	GTG	TCT	1776
	Asp	Val	Arg	Leu	Ser		560				565					570	
						575											
60	GGC	ATG	TGC	TGT	GCT	CTG	TTC	CTG	CTG	ATC	CTG	CTC	ACG	GGG	GTC	CTG	1824
	Gly	Met	Cys	Cys	Ala	Leu	Phe	Leu	Leu	Ile	Leu	Leu	Thr	Gly	Val	Leu	
					575					580					585		
65	TGC	CAC	CGT	TTC	CAT	GGC	CTG	TGG	TAT	ATG	AAA	ATG	ATG	TGG	GCC	TGG	1872
	Cys	His	Arg	Phe	His	Gly	Leu	Trp	Tyr	Met	Lys	Met	Met	Trp	Ala	Trp	
				590					595					600			
70	CTC	CAG	GCC	AAA	AGG	AAG	CCC	AGG	AAA	GCT	CCC	AGC	AGG	AAC	ATC	TGC	1920
	Leu	Gln	Ala	Lys	Arg	Lys	Pro	Arg	Lys	Ala	Pro	Ser	Arg	Asn	Ile	Cys	
			605					610					615				
75	TAT	GAT	GCA	TTT	GTT	TCT	TAC	AGT	GAG	CGG	GAT	GCC	TAC	TGG	GTG	GAG	1968
	Tyr	Asp	Ala	Phe	Val	Ser	Tyr	Ser	Glu	Arg	Asp	Ala	Tyr	Trp	Val	Glu	

	45	50	55
	Leu Val 60	Leu Thr Ser Asn Gly 65	Ile Asn Thr Ile Glu 70
5	Ser 75	Ser Leu Gly Ser 80	Leu Asp 85
	Ser Asn 95	Leu Ser Trp Phe Lys 100	Pro Leu Ser Ser 105
10	Leu Asn 110	Leu Gly Asn Pro Tyr 115	Thr Leu Gly Glu 120
	Phe Ser 125	His Leu Thr Lys 130	Leu Arg Val Gly 135
15	Thr Phe 140	Thr Lys Ile Gln Arg 145	Lys Asp Phe Ala Gly 150
20	Glu Glu 155	Leu Glu Ile Asp 160	Ala Ser Asp Leu Gln 165
	Ser Leu 175	Lys Ser Ile Gln Asn Val 180	Ser His Leu Ile Leu 185
25	Gln His 190	Ile Leu Leu Glu Ile 195	Phe Val Asp Val Thr 200
	Glu Cys 205	Leu Glu Leu Arg Asp 210	Thr Asp Leu Asp Thr 215
30	Glu Leu 220	Ser Thr Gly Glu Thr 225	Asn Ser Leu Ile Lys 230
35	Arg Asn 235	Val Lys Ile Thr Asp 240	Glu Ser Leu Phe Gln 245
	Leu Asn 255	Gln Ile Ser Gly Leu Leu Glu 260	Leu Glu Phe Asp Asp 265
40	Leu Asn 270	Gly Val Gly Asn Phe Arg 275	Ala Ser Asp Asn Asp 280
	Asp Pro 285	Gly Lys Val Glu Thr 290	Ile Arg Arg Leu His 295
45	Arg Phe 300	Tyr Leu Phe Tyr Asp 305	Leu Ser Thr Leu Tyr 310
50	Arg Val 315	Lys Arg Ile Thr Val Glu 320	Asn Ser Lys Val Phe 325
	Cys Leu 335	Leu Ser Gln His Leu Lys 340	Ser Leu Tyr Leu Asp 345
55	Glu Asn 350	Leu Met Val Glu Glu Tyr 355	Lys Asn Ser Ala Cys 360
60	Ala Trp 365	Pro Ser Leu Gln Thr 370	Leu Ile Leu Arg Gln 375

Ser Leu Glu Lys Thr Gly Glu Thr Leu Leu Thr Leu Lys Asn Leu Thr
 380 385 390
 5 Asn Ile Asp Ile Ser Lys Asn Ser Phe His Ser Met Pro Glu Thr Cys
 395 400 405 410
 Gln Trp Pro Glu Lys Met Lys Tyr Leu Asn Leu Ser Ser Thr Arg Ile
 415 420 425
 10 His Ser Val Thr Gly Cys Ile Pro Lys Thr Leu Glu Ile Leu Asp Val
 430 435 440
 15 Ser Asn Asn Asn Leu Asn Leu Phe Ser Leu Asn Leu Pro Gln Leu Lys
 445 450 455
 Glu Leu Tyr Ile Ser Arg Asn Lys Leu Met Thr Leu Pro Asp Ala Ser
 460 465 470
 20 Leu Leu Pro Met Leu Leu Val Leu Lys Ile Ser Arg Asn Ala Ile Thr
 475 480 485 490
 Thr Phe Ser Lys Glu Gln Leu Asp Ser Phe His Thr Leu Lys Thr Leu
 495 500 505
 25 Glu Ala Gly Gly Asn Asn Phe Ile Cys Ser Cys Glu Phe Leu Ser Phe
 510 515 520
 Thr Gln Glu Gln Gln Ala Leu Ala Lys Val Leu Ile Asp Trp Pro Ala
 525 530 535
 30 Asn Tyr Leu Cys Asp Ser Pro Ser His Val Arg Gly Gln Gln Val Gln
 540 545 550
 35 Asp Val Arg Leu Ser Val Ser Glu Cys His Arg Thr Ala Leu Val Ser
 555 560 565 570
 Gly Met Cys Cys Ala Leu Phe Leu Leu Ile Leu Leu Thr Gly Val Leu
 575 580 585
 40 Cys His Arg Phe His Gly Leu Trp Tyr Met Lys Met Met Trp Ala Trp
 590 595 600
 Leu Gln Ala Lys Arg Lys Pro Arg Lys Ala Pro Ser Arg Asn Ile Cys
 605 610 615
 45 Tyr Asp Ala Phe Val Ser Tyr Ser Glu Arg Asp Ala Tyr Trp Val Glu
 620 625 630
 50 Asn Leu Met Val Gln Glu Leu Glu Asn Phe Asn Pro Pro Phe Lys Leu
 635 640 645 650
 Cys Leu His Lys Arg Asp Phe Ile Pro Gly Lys Trp Ile Ile Asp Asn
 655 660 665
 55 Ile Ile Asp Ser Ile Glu Lys Ser His Lys Thr Val Phe Val Leu Ser
 670 675 680
 60 Glu Asn Phe Val Lys Ser Glu Trp Cys Lys Tyr Glu Leu Asp Phe Ser
 685 690 695

His Phe Arg Leu Phe Glu Glu Asn Asn Asp Ala Ala Ile Leu Ile Leu
 700 705 710
 5 Leu Glu Pro Ile Glu Lys Lys Ala Ile Pro Gln Arg Phe Cys Lys Leu
 715 720 725 730
 Arg Lys Ile Met Asn Thr Lys Thr Tyr Leu Glu Trp Pro Met Asp Glu
 735 740 745
 10 Ala Gln Arg Glu Gly Phe Trp Val Asn Leu Arg Ala Ala Ile Lys Ser
 750 755 760

(2) INFORMATION FOR SEQ ID NO:5:

15

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2715 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: cDNA

25

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..2712

30

- (ix) FEATURE:
 (A) NAME/KEY: mat_peptide
 (B) LOCATION: 64..2712

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG AGA CAG ACT TTG CCT TGT ATC TAC TTT TGG GGG GGC CTT TTG CCC 48
 Met Arg Gln Thr Leu Pro Cys Ile Tyr Phe Trp Gly Gly Leu Leu Pro
 -21 -20 -15 -10
 40 TTT GGG ATG CTG TGT GCA TCC TCC ACC ACC AAG TGC ACT GTT AGC CAT 96
 Phe Gly Met Leu Cys Ala Ser Ser Thr Thr Lys Cys Thr Val Ser His
 -5 1 5 10
 45 GAA GTT GCT GAC TGC AGC CAC CTG AAG TTG ACT CAG GTA CCC GAT GAT 144
 Glu Val Ala Asp Cys Ser His Leu Lys Leu Thr Gln Val Pro Asp Asp
 15 20 25
 50 CTA CCC ACA AAC ATA ACA GTG TTG AAC CTT ACC CAT AAT CAA CTC AGA 192
 Leu Pro Thr Asn Ile Thr Val Leu Asn Leu Thr His Asn Gln Leu Arg
 30 35 40
 AGA TTA CCA GCC GCC AAC TTC ACA AGG TAT AGC CAG CTA ACT AGC TTG 240
 Arg Leu Pro Ala Ala Asn Phe Thr Arg Tyr Ser Gln Leu Thr Ser Leu
 45 50 55
 55 GAT GTA GGA TTT AAC ACC ATC TCA AAA CTG GAG CCA GAA TTG TGC CAG 288
 Asp Val Gly Phe Asn Thr Ile Ser Lys Leu Glu Pro Glu Leu Cys Gln
 60 65 70 75
 60 AAA CTT CCC ATG TTA AAA GTT TTG AAC CTC CAG CAC AAT GAG CTA TCT 336
 Lys Leu Pro Met Leu Lys Val Leu Asn Leu Gln His Asn Glu Leu Ser

	80								85								90								
5	CAA Gln	CTT Leu	TCT Ser	GAT Asp 95	AAA Lys	ACC Thr	TTT Phe	GCC Ala	TTC Phe 100	TGC Cys	ACG Thr	AAT Asn	TTG Leu	ACT Thr 105	GAA Glu	CTC Leu	384								
10	CAT His	CTC Leu	ATG Met 110	TCC Ser	AAC Asn	TCA Ser	ATC Ile	CAG Gln 115	AAA Lys	ATT Ile	AAA Lys	AAT Asn	AAT Asn 120	CCC Pro	TTT Phe	GTC Val	432								
15	AAG Lys	CAG Gln 125	AAG Lys	AAT Asn	TTA Leu	ATC Ile	ACA Thr 130	TTA Leu	GAT Asp	CTG Leu	TCT Ser	CAT His 135	AAT Asn	GGC Gly	TTG Leu	TCA Ser	480								
20	TCT Ser 140	ACA Thr	AAA Lys	TTA Leu	GGA Gly	ACT Thr 145	CAG Gln	GTT Val	CAG Gln	CTG Leu	GAA Glu 150	AAT Asn	CTC Leu	CAA Gln	GAG Glu 155	CTT Leu	528								
25	CTA Leu	TTA Leu	TCA Ser	AAC Asn 160	AAT Asn	AAA Lys	ATT Ile	CAA Gln	GCG Ala 165	CTA Leu	AAA Lys	AGT Ser	GAA Glu	GAA Glu	CTG Leu 170	GAT Asp	576								
30	ATC Ile	TTT Phe	GCC Ala	AAT Asn 175	TCA Ser	TCT Ser	TTA Leu	AAA Lys	AAA Lys 180	TTA Leu	GAG Glu	TTG Leu	TCA Ser	TCG Ser 185	AAT Asn	CAA Gln	624								
35	ATT Ile	AAA Lys	GAG Glu 190	TTT Phe	TCT Ser	CCA Pro	GGG Gly	TGT Cys 195	TTT Phe	CAC His	GCA Ala	ATT Ile	GGA Gly 200	AGA Arg	TTA Leu	TTT Phe	672								
40	GGC Gly 205	CTC Leu	TTT Phe	CTG Leu	AAC Asn	AAT Asn	GTC Val 210	CAG Gln	CTG Leu	GGT Gly	CCC Pro 215	AGC Ser	CTT Leu	ACA Thr	GAG Glu	AAG Lys	720								
45	CTA Leu 220	TGT Cys	TTG Leu	GAA Glu	TTA Leu	GCA Ala 225	AAC Asn	ACA Thr	AGC Ser	ATT Ile	CGG Arg 230	AAT Asn	CTG Leu	TCT Ser	CTG Leu 235	AGT Ser	768								
50	AAC Asn	AGC Ser	CAG Gln	CTG Leu 240	TCC Ser	ACC Thr	ACC Thr	AGC Ser	AAT Asn 245	ACA Thr	ACT Thr	TTC Phe	TTG Leu	GGA Gly 250	CTA Leu	AAG Lys	816								
55	TGG Trp	ACA Thr	AAT Asn	CTC Leu 255	ACT Thr	ATG Met	CTC Leu	GAT Asp	CTT Leu 260	TCC Ser	TAC Tyr	AAC Asn	AAC Asn	TTA Leu 265	AAT Asn	GTG Val	864								
60	GTT Val	GGT Gly	AAC Asn 270	GAT Asp	TCC Ser	TTT Phe	GCT Ala	TGG Trp 275	CTT Leu	CCA Pro	CAA Gln	CTA Leu	GAA Glu 280	TAT Tyr	TTC Phe	TTC Phe	912								
65	CTA Leu 285	GAG Glu	TAT Tyr	AAT Asn	AAT Asn	ATA Ile	CAG Gln 290	CAT His	TTG Leu	TTT Phe	TCT Ser	CAC His 295	TCT Ser	TTG Leu	CAC His	GGG Gly	960								
70	CTT Leu 300	TTC Phe	AAT Asn	GTG Val	AGG Arg	TAC Tyr 305	CTG Leu	AAT Asn	TTG Leu	AAA Lys	CGG Arg 310	TCT Ser	TTT Phe	ACT Thr	AAA Lys	CAA Gln 315	1008								
75	AGT Ser	ATT Ile	TCC Ser	CTT Leu 320	GCC Ala	TCA Ser	CTC Leu	CCC Pro	AAG Lys 325	ATT Ile	GAT Asp	GAT Asp	TTT Phe	TCT Ser	TTT Phe 330	CAG Gln	1056								

		TGG	CTA	AAA	TGT	TTG	GAG	CAC	CTT	AAC	ATG	GAA	GAT	AAT	GAT	ATT	CCA	1104
		Trp	Leu	Lys	Cys	Leu	Glu	His	Leu	Asn	Met	Glu	Asp	Asn	Asp	Ile	Pro	
					335					340					345			
5		GGC	ATA	AAA	AGC	AAT	ATG	TTC	ACA	GGA	TTG	ATA	AAC	CTG	AAA	TAC	TTA	1152
		Gly	Ile	Lys	Ser	Asn	Met	Phe	Thr	Gly	Leu	Ile	Asn	Leu	Lys	Tyr	Leu	
				350					355					360				
10		AGT	CTA	TCC	AAC	TCC	TTT	ACA	AGT	TTG	CGA	ACT	TTG	ACA	AAT	GAA	ACA	1200
		Ser	Leu	Ser	Asn	Ser	Phe	Thr	Ser	Leu	Arg	Thr	Leu	Thr	Asn	Glu	Thr	
			365					370					375					
15		TTT	GTA	TCA	CTT	GCT	CAT	TCT	CCC	TTA	CAC	ATA	CTC	AAC	CTA	ACC	AAG	1248
		Phe	Val	Ser	Leu	Ala	His	Ser	Pro	Leu	His	Ile	Leu	Asn	Leu	Thr	Lys	
		380					385					390					395	
20		AAT	AAA	ATC	TCA	AAA	ATA	GAG	AGT	GAT	GCT	TTC	TCT	TGG	TTG	GGC	CAC	1296
		Asn	Lys	Ile	Ser	Lys	Ile	Glu	Ser	Asp	Ala	Phe	Ser	Trp	Leu	Gly	His	
						400					405					410		
25		CTA	GAA	GTA	CTT	GAC	CTG	GGC	CTT	AAT	GAA	ATT	GGG	CAA	GAA	CTC	ACA	1344
		Leu	Glu	Val	Leu	Asp	Leu	Gly	Leu	Asn	Glu	Ile	Gly	Gln	Glu	Leu	Thr	
					415				420					425				
30		GGC	CAG	GAA	TGG	AGA	GGT	CTA	GAA	AAT	ATT	TTC	GAA	ATC	TAT	CTT	TCC	1392
		Gly	Gln	Glu	Trp	Arg	Gly	Leu	Glu	Asn	Ile	Phe	Glu	Ile	Tyr	Leu	Ser	
			430					435					440					
35		TAC	AAC	AAG	TAC	CTG	CAG	CTG	ACT	AGG	AAC	TCC	TTT	GCC	TTG	GTC	CCA	1440
		Tyr	Asn	Lys	Tyr	Leu	Gln	Leu	Thr	Arg	Asn	Ser	Phe	Ala	Leu	Val	Pro	
			445					450					455					
40		AGC	CTT	CAA	CGA	CTG	ATG	CTC	CGA	AGG	GTG	GCC	CTT	AAA	AAT	GTG	GAT	1488
		Ser	Leu	Gln	Arg	Leu	Met	Leu	Arg	Arg	Val	Ala	Leu	Lys	Asn	Val	Asp	
		460					465				470					475		
45		AGC	TCT	CCT	TCA	CCA	TTC	CAG	CCT	CTT	CGT	AAC	TTG	ACC	ATT	CTG	GAT	1536
		Ser	Ser	Pro	Ser	Pro	Phe	Gln	Pro	Leu	Arg	Asn	Leu	Thr	Ile	Leu	Asp	
					480					485						490		
50		CTA	AGC	AAC	AAC	AAC	ATA	GCC	AAC	ATA	AAT	GAT	GAC	ATG	TTG	GAG	GGT	1584
		Leu	Ser	Asn	Asn	Asn	Ile	Ala	Asn	Ile	Asn	Asp	Asp	Met	Leu	Glu	Gly	
					495				500					505				
55		CTT	GAG	AAA	CTA	GAA	ATT	CTC	GAT	TTG	CAG	CAT	AAC	AAC	TTA	GCA	CGG	1632
		Leu	Glu	Lys	Leu	Glu	Ile	Leu	Asp	Leu	Gln	His	Asn	Asn	Leu	Ala	Arg	
				510					515					520				
60		CTC	TGG	AAA	CAC	GCA	AAC	CCT	GGT	GGT	CCC	ATT	TAT	TTC	CTA	AAG	GGT	1680
		Leu	Trp	Lys	His	Ala	Asn	Pro	Gly	Gly	Pro	Ile	Tyr	Phe	Leu	Lys	Gly	
			525					530					535					
65		CTG	TCT	CAC	CTC	CAC	ATC	CTT	AAC	TTG	GAG	TCC	AAC	GGC	TTT	GAC	GAG	1728
		Leu	Ser	His	Leu	His	Ile	Leu	Asn	Leu	Glu	Ser	Asn	Gly	Phe	Asp	Glu	
		540					545					550					555	
70		ATC	CCA	GTT	GAG	GTC	TTC	AAG	GAT	TTA	TTT	GAA	CTA	AAG	ATC	ATC	GAT	1776
		Ile	Pro	Val	Glu	Val	Phe	Lys	Asp	Leu	Phe	Glu	Leu	Lys	Ile	Ile	Asp	
					560						565					570		

	TTA GGA TTG AAT AAT TTA AAC ACA CTT CCA GCA TCT GTC TTT AAT AAT	1824
	Leu Gly Leu Asn Asn Leu Asn Thr Leu Pro Ala Ser Val Phe Asn Asn	
	575 580 585	
5	CAG GTG TCT CTA AAG TCA TTG AAC CTT CAG AAG AAT CTC ATA ACA TCC	1872
	Gln Val Ser Leu Lys Ser Leu Asn Leu Gln Lys Asn Leu Ile Thr Ser	
	590 595 600	
10	GTT GAG AAG AAG GTT TTC GGG CCA GCT TTC AGG AAC CTG ACT GAG TTA	1920
	Val Glu Lys Lys Val Phe Gly Pro Ala Phe Arg Asn Leu Thr Glu Leu	
	605 610 615	
15	GAT ATG CGC TTT AAT CCC TTT GAT TGC ACG TGT GAA AGT ATT GCC TGG	1968
	Asp Met Arg Phe Asn Pro Phe Asp Cys Thr Cys Glu Ser Ile Ala Trp	
	620 625 630 635	
20	TTT GTT AAT TGG ATT AAC GAG ACC CAT ACC AAC ATC CCT GAG CTG TCA	2016
	Phe Val Asn Trp Ile Asn Glu Thr His Thr Asn Ile Pro Glu Leu Ser	
	640 645 650	
20	AGC CAC TAC CTT TGC AAC ACT CCA CCT CAC TAT CAT GGG TTC CCA GTG	2064
	Ser His Tyr Leu Cys Asn Thr Pro Pro His Tyr His Gly Phe Pro Val	
	655 660 665	
25	AGA CTT TTT GAT ACA TCA TCT TGC AAA GAC AGT GCC CCC TTT GAA CTC	2112
	Arg Leu Phe Asp Thr Ser Ser Cys Lys Asp Ser Ala Pro Phe Glu Leu	
	670 675 680	
30	TTT TTC ATG ATC AAT ACC AGT ATC CTG TTG ATT TTT ATC TTT ATT GTA	2160
	Phe Phe Met Ile Asn Thr Ser Ile Leu Leu Ile Phe Ile Phe Ile Val	
	685 690 695	
35	CTT CTC ATC CAC TTT GAG GGC TGG AGG ATA TCT TTT TAT TGG AAT GTT	2208
	Leu Leu Ile His Phe Glu Gly Trp Arg Ile Ser Phe Tyr Trp Asn Val	
	700 705 710 715	
40	TCA GTA CAT CGA GTT CTT GGT TTC AAA GAA ATA GAC AGA CAG ACA GAA	2256
	Ser Val His Arg Val Leu Gly Phe Lys Glu Ile Asp Arg Gln Thr Glu	
	720 725 730	
40	CAG TTT GAA TAT GCA GCA TAT ATA ATT CAT GCC TAT AAA GAT AAG GAT	2304
	Gln Phe Glu Tyr Ala Ala Tyr Ile Ile His Ala Tyr Lys Asp Lys Asp	
	735 740 745	
45	TGG GTC TGG GAA CAT TTC TCT TCA ATG GAA AAG GAA GAC CAA TCT CTC	2352
	Trp Val Trp Glu His Phe Ser Ser Met Glu Lys Glu Asp Gln Ser Leu	
	750 755 760	
50	AAA TTT TGT CTG GAA GAA AGG GAC TTT GAG GCG GGT GTT TTT GAA CTA	2400
	Lys Phe Cys Leu Glu Glu Arg Asp Phe Glu Ala Gly Val Phe Glu Leu	
	765 770 775	
55	GAA GCA ATT GTT AAC AGC ATC AAA AGA AGC AGA AAA ATT ATT TTT GTT	2448
	Glu Ala Ile Val Asn Ser Ile Lys Arg Ser Arg Lys Ile Ile Phe Val	
	780 785 790 795	
60	ATA ACA CAC CAT CTA TTA AAA GAC CCA TTA TGC AAA AGA TTC AAG GTA	2496
	Ile Thr His His Leu Leu Lys Asp Pro Leu Cys Lys Arg Phe Lys Val	
	800 805 810	
60	CAT CAT GCA GTT CAA CAA GCT ATT GAA CAA AAT CTG GAT TCC ATT ATA	2544

	His	His	Ala	Val	Gln	Gln	Ala	Ile	Glu	Gln	Asn	Leu	Asp	Ser	Ile	Ile	
				815					820					825			
5	TTG	GTT	TTC	CTT	GAG	GAG	ATT	CCA	GAT	TAT	AAA	CTG	AAC	CAT	GCA	CTC	2592
	Leu	Val	Phe	Leu	Glu	Glu	Ile	Pro	Asp	Tyr	Lys	Leu	Asn	His	Ala	Leu	
				830				835					840				
10	TGT	TTG	CGA	AGA	GGA	ATG	TTT	AAA	TCT	CAC	TGC	ATC	TTG	AAC	TGG	CCA	2640
	Cys	Leu	Arg	Arg	Gly	Met	Phe	Lys	Ser	His	Cys	Ile	Leu	Asn	Trp	Pro	
				845			850					855					
15	GTT	CAG	AAA	GAA	CGG	ATA	GGT	GCC	TTT	CGT	CAT	AAA	TTG	CAA	GTA	GCA	2688
	Val	Gln	Lys	Glu	Arg	Ile	Gly	Ala	Phe	Arg	His	Lys	Leu	Gln	Val	Ala	
						865					870					875	
	CTT	GGA	TCC	AAA	AAC	TCT	GTA	CAT	TAA								2715
	Leu	Gly	Ser	Lys	Asn	Ser	Val	His									
						880											
20	(2) INFORMATION FOR SEQ ID NO:6:																
	(i) SEQUENCE CHARACTERISTICS:																
	(A) LENGTH: 904 amino acids																
25	(B) TYPE: amino acid																
	(D) TOPOLOGY: linear																
	(ii) MOLECULE TYPE: protein																
30	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:																
	Met	Arg	Gln	Thr	Leu	Pro	Cys	Ile	Tyr	Phe	Trp	Gly	Gly	Leu	Leu	Pro	
	-21	-20					-15					-10					
35	Phe	Gly	Met	Leu	Cys	Ala	Ser	Ser	Thr	Thr	Lys	Cys	Thr	Val	Ser	His	
	-5					1				5					10		
	Glu	Val	Ala	Asp	Cys	Ser	His	Leu	Lys	Leu	Thr	Gln	Val	Pro	Asp	Asp	
				15					20					25			
40	Leu	Pro	Thr	Asn	Ile	Thr	Val	Leu	Asn	Leu	Thr	His	Asn	Gln	Leu	Arg	
				30				35					40				
45	Arg	Leu	Pro	Ala	Ala	Asn	Phe	Thr	Arg	Tyr	Ser	Gln	Leu	Thr	Ser	Leu	
		45				50						55					
	Asp	Val	Gly	Phe	Asn	Thr	Ile	Ser	Lys	Leu	Glu	Pro	Glu	Leu	Cys	Gln	
		60				65					70					75	
50	Lys	Leu	Pro	Met	Leu	Lys	Val	Leu	Asn	Leu	Gln	His	Asn	Glu	Leu	Ser	
					80					85					90		
	Gln	Leu	Ser	Asp	Lys	Thr	Phe	Ala	Phe	Cys	Thr	Asn	Leu	Thr	Glu	Leu	
				95					100					105			
55	His	Leu	Met	Ser	Asn	Ser	Ile	Gln	Lys	Ile	Lys	Asn	Asn	Pro	Phe	Val	
				110					115					120			
60	Lys	Gln	Lys	Asn	Leu	Ile	Thr	Leu	Asp	Leu	Ser	His	Asn	Gly	Leu	Ser	
		125					130					135					

Ser Thr Lys Leu Gly Thr Gln Val Gln Leu Glu Asn Leu Gln Glu Leu
 140 145 150 155
 5 Leu Leu Ser Asn Asn Lys Ile Gln Ala Leu Lys Ser Glu Glu Leu Asp
 160 165 170
 Ile Phe Ala Asn Ser Ser Leu Lys Lys Leu Glu Leu Ser Ser Asn Gln
 175 180 185
 10 Ile Lys Glu Phe Ser Pro Gly Cys Phe His Ala Ile Gly Arg Leu Phe
 190 195 200
 Gly Leu Phe Leu Asn Asn Val Gln Leu Gly Pro Ser Leu Thr Glu Lys
 205 210 215
 15 Leu Cys Leu Glu Leu Ala Asn Thr Ser Ile Arg Asn Leu Ser Leu Ser
 220 225 230 235
 20 Asn Ser Gln Leu Ser Thr Thr Ser Asn Thr Thr Phe Leu Gly Leu Lys
 240 245 250
 Trp Thr Asn Leu Thr Met Leu Asp Leu Ser Tyr Asn Asn Leu Asn Val
 255 260 265
 25 Val Gly Asn Asp Ser Phe Ala Trp Leu Pro Gln Leu Glu Tyr Phe Phe
 270 275 280
 Leu Glu Tyr Asn Asn Ile Gln His Leu Phe Ser His Ser Leu His Gly
 285 290 295
 30 Leu Phe Asn Val Arg Tyr Leu Asn Leu Lys Arg Ser Phe Thr Lys Gln
 300 305 310 315
 35 Ser Ile Ser Leu Ala Ser Leu Pro Lys Ile Asp Asp Phe Ser Phe Gln
 320 325 330
 Trp Leu Lys Cys Leu Glu His Leu Asn Met Glu Asp Asn Asp Ile Pro
 335 340 345
 40 Gly Ile Lys Ser Asn Met Phe Thr Gly Leu Ile Asn Leu Lys Tyr Leu
 350 355 360
 Ser Leu Ser Asn Ser Phe Thr Ser Leu Arg Thr Leu Thr Asn Glu Thr
 365 370 375
 45 Phe Val Ser Leu Ala His Ser Pro Leu His Ile Leu Asn Leu Thr Lys
 380 385 390 395
 50 Asn Lys Ile Ser Lys Ile Glu Ser Asp Ala Phe Ser Trp Leu Gly His
 400 405 410
 Leu Glu Val Leu Asp Leu Gly Leu Asn Glu Ile Gly Gln Glu Leu Thr
 415 420 425
 55 Gly Gln Glu Trp Arg Gly Leu Glu Asn Ile Phe Glu Ile Tyr Leu Ser
 430 435 440
 Tyr Asn Lys Tyr Leu Gln Leu Thr Arg Asn Ser Phe Ala Leu Val Pro
 445 450 455
 60 Ser Leu Gln Arg Leu Met Leu Arg Arg Val Ala Leu Lys Asn Val Asp

	460		465		470		475
	Ser Ser Pro Ser	Pro Phe Gln Pro Leu Arg Asn Leu Thr Ile Leu Asp					
		480		485		490	
5	Leu Ser Asn Asn Asn Ile Ala Asn Ile Asn Asp Asp Met Leu Glu Gly		500		505		
		495					
10	Leu Glu Lys Leu Glu Ile Leu Asp Leu Gln His Asn Asn Leu Ala Arg		515		520		
		510					
	Leu Trp Lys His Ala Asn Pro Gly Gly Pro Ile Tyr Phe Leu Lys Gly		530		535		
		525					
15	Leu Ser His Leu His Ile Leu Asn Leu Glu Ser Asn Gly Phe Asp Glu		545		550		555
	Ile Pro Val Glu Val Phe Lys Asp Leu Phe Glu Leu Lys Ile Ile Asp		560		565		570
20							
	Leu Gly Leu Asn Asn Leu Asn Thr Leu Pro Ala Ser Val Phe Asn Asn		575		580		585
25	Gln Val Ser Leu Lys Ser Leu Asn Leu Gln Lys Asn Leu Ile Thr Ser		590		595		600
	Val Glu Lys Lys Val Phe Gly Pro Ala Phe Arg Asn Leu Thr Glu Leu		605		610		615
30	Asp Met Arg Phe Asn Pro Phe Asp Cys Thr Cys Glu Ser Ile Ala Trp		620		625		630
	Phe Val Asn Trp Ile Asn Glu Thr His Thr Asn Ile Pro Glu Leu Ser		640		645		650
35							
	Ser His Tyr Leu Cys Asn Thr Pro Pro His Tyr His Gly Phe Pro Val		655		660		665
40	Arg Leu Phe Asp Thr Ser Ser Cys Lys Asp Ser Ala Pro Phe Glu Leu		670		675		680
	Phe Phe Met Ile Asn Thr Ser Ile Leu Leu Ile Phe Ile Phe Ile Val		685		690		695
45	Leu Leu Ile His Phe Glu Gly Trp Arg Ile Ser Phe Tyr Trp Asn Val		700		705		710
	Ser Val His Arg Val Leu Gly Phe Lys Glu Ile Asp Arg Gln Thr Glu		720		725		730
50							
	Gln Phe Glu Tyr Ala Ala Tyr Ile Ile His Ala Tyr Lys Asp Lys Asp		735		740		745
55	Trp Val Trp Glu His Phe Ser Ser Met Glu Lys Glu Asp Gln Ser Leu		750		755		760
	Lys Phe Cys Leu Glu Glu Arg Asp Phe Glu Ala Gly Val Phe Glu Leu		765		770		775
60	Glu Ala Ile Val Asn Ser Ile Lys Arg Ser Arg Lys Ile Ile Phe Val		780		785		790
							795

Ile Thr His His Leu Leu Lys Asp Pro Leu Cys Lys Arg Phe Lys Val
800 805 810

5 His His Ala Val Gln Gln Ala Ile Glu Gln Asn Leu Asp Ser Ile Ile
815 820 825

Leu Val Phe Leu Glu Glu Ile Pro Asp Tyr Lys Leu Asn His Ala Leu
830 835 840

10 Cys Leu Arg Arg Gly Met Phe Lys Ser His Cys Ile Leu Asn Trp Pro
845 850 855

Val Gln Lys Glu Arg Ile Gly Ala Phe Arg His Lys Leu Gln Val Ala
15 860 865 870 875

Leu Gly Ser Lys Asn Ser Val His
880

20 (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2400 base pairs
(B) TYPE: nucleic acid
25 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..2397

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATG GAG CTG AAT TTC TAC AAA ATC CCC GAC AAC CTC CCC TTC TCA ACC 48
Met Glu Leu Asn Phe Tyr Lys Ile Pro Asp Asn Leu Pro Phe Ser Thr
40 1 5 10 15

AAG AAC CTG GAC CTG AGC TTT AAT CCC CTG AGG CAT TTA GGC AGC TAT 96
Lys Asn Leu Asp Leu Ser Phe Asn Pro Leu Arg His Leu Gly Ser Tyr
20 25 30

45 AGC TTC TTC AGT TTC CCA GAA CTG CAG GTG CTG GAT TTA TCC AGG TGT 144
Ser Phe Phe Ser Phe Pro Glu Leu Gln Val Leu Asp Leu Ser Arg Cys
35 40 45

50 GAA ATC CAG ACA ATT GAA GAT GGG GCA TAT CAG AGC CTA AGC CAC CTC 192
Glu Ile Gln Thr Ile Glu Asp Gly Ala Tyr Gln Ser Leu Ser His Leu
50 55 60

55 TCT ACC TTA ATA TTG ACA GGA AAC CCC ATC CAG AGT TTA GCC CTG GGA 240
Ser Thr Leu Ile Leu Thr Gly Asn Pro Ile Gln Ser Leu Ala Leu Gly
65 70 75 80

GCC TTT TCT GGA CTA TCA AGT TTA CAG AAG CTG GTG GCT GTG GAG ACA 288
Ala Phe Ser Gly Leu Ser Ser Leu Gln Lys Leu Val Ala Val Glu Thr
60 85 90 95

		AAT	CTA	GCA	TCT	CTA	GAG	AAC	TTC	CCC	ATT	GGA	CAT	CTC	AAA	ACT	TTG	336
		Asn	Leu	Ala	Ser	Leu	Glu	Asn	Phe	Pro	Ile	Gly	His	Leu	Lys	Thr	Leu	
					100					105					110			
5		AAA	GAA	CTT	AAT	GTG	GCT	CAC	AAT	CTT	ATC	CAA	TCT	TTC	AAA	TTA	CCT	384
		Lys	Glu	Leu	Asn	Val	Ala	His	Asn	Leu	Ile	Gln	Ser	Phe	Lys	Leu	Pro	
				115					120					125				
10		GAG	TAT	TTT	TCT	AAT	CTG	ACC	AAT	CTA	GAG	CAC	TTG	GAC	CTT	TCC	AGC	432
		Glu	Tyr	Phe	Ser	Asn	Leu	Thr	Asn	Leu	Glu	His	Leu	Asp	Leu	Ser	Ser	
			130					135					140					
15		AAC	AAG	ATT	CAA	AGT	ATT	TAT	TGC	ACA	GAC	TTG	CGG	GTT	CTA	CAT	CAA	480
		Asn	Lys	Ile	Gln	Ser	Ile	Tyr	Cys	Thr	Asp	Leu	Arg	Val	Leu	His	Gln	
		145					150					155					160	
20		ATG	CCC	CTA	CTC	AAT	CTC	TCT	TTA	GAC	CTG	TCC	CTG	AAC	CCT	ATG	AAC	528
		Met	Pro	Leu	Leu	Asn	Leu	Ser	Leu	Asp	Leu	Ser	Leu	Asn	Pro	Met	Asn	
						165					170					175		
20		TTT	ATC	CAA	CCA	GGT	GCA	TTT	AAA	GAA	ATT	AGG	CTT	CAT	AAG	CTG	ACT	576
		Phe	Ile	Gln	Pro	Gly	Ala	Phe	Lys	Glu	Ile	Arg	Leu	His	Lys	Leu	Thr	
					180					185					190			
25		TTA	AGA	AAT	AAT	TTT	GAT	AGT	TTA	AAT	GTA	ATG	AAA	ACT	TGT	ATT	CAA	624
		Leu	Arg	Asn	Asn	Phe	Asp	Ser	Leu	Asn	Val	Met	Lys	Thr	Cys	Ile	Gln	
				195					200					205				
30		GGT	CTG	GCT	GGT	TTA	GAA	GTC	CAT	CGT	TTG	GTT	CTG	GGA	GAA	TTT	AGA	672
		Gly	Leu	Ala	Gly	Leu	Glu	Val	His	Arg	Leu	Val	Leu	Gly	Glu	Phe	Arg	
			210					215					220					
35		AAT	GAA	GGA	AAC	TTG	GAA	AAG	TTT	GAC	AAA	TCT	GCT	CTA	GAG	GGC	CTG	720
		Asn	Glu	Gly	Asn	Leu	Glu	Lys	Phe	Asp	Lys	Ser	Ala	Leu	Glu	Gly	Leu	
		225					230					235					240	
40		TGC	AAT	TTG	ACC	ATT	GAA	GAA	TTC	CGA	TTA	GCA	TAC	TTA	GAC	TAC	TAC	768
		Cys	Asn	Leu	Thr	Ile	Glu	Glu	Phe	Arg	Leu	Ala	Tyr	Leu	Asp	Tyr	Tyr	
					245						250					255		
40		CTC	GAT	GAT	ATT	ATT	GAC	TTA	TTT	AAT	TGT	TTG	ACA	AAT	GTT	TCT	TCA	816
		Leu	Asp	Asp	Ile	Ile	Asp	Leu	Phe	Asn	Cys	Leu	Thr	Asn	Val	Ser	Ser	
					260					265					270			
45		TTT	TCC	CTG	GTG	AGT	GTG	ACT	ATT	GAA	AGG	GTA	AAA	GAC	TTT	TCT	TAT	864
		Phe	Ser	Leu	Val	Ser	Val	Thr	Ile	Glu	Arg	Val	Lys	Asp	Phe	Ser	Tyr	
				275				280						285				
50		AAT	TTC	GGA	TGG	CAA	CAT	TTA	GAA	TTA	GTT	AAC	TGT	AAA	TTT	GGA	CAG	912
		Asn	Phe	Gly	Trp	Gln	His	Leu	Glu	Leu	Val	Asn	Cys	Lys	Phe	Gly	Gln	
			290					295					300					
55		TTT	CCC	ACA	TTG	AAA	CTC	AAA	TCT	CTC	AAA	AGG	CTT	ACT	TTC	ACT	TCC	960
		Phe	Pro	Thr	Leu	Lys	Leu	Lys	Ser	Leu	Lys	Arg	Leu	Thr	Phe	Thr	Ser	
		305					310					315					320	
60		AAC	AAA	GGT	GGG	AAT	GCT	TTT	TCA	GAA	GTT	GAT	CTA	CCA	AGC	CTT	GAG	1008
		Asn	Lys	Gly	Gly	Asn	Ala	Phe	Ser	Glu	Val	Asp	Leu	Pro	Ser	Leu	Glu	
						325					330					335		
60		TTT	CTA	GAT	CTC	AGT	AGA	AAT	GGC	TTG	AGT	TTC	AAA	GGT	TGC	TGT	TCT	1056

	Phe	Leu	Asp	Leu	Ser	Arg	Asn	Gly	Leu	Ser	Phe	Lys	Gly	Cys	Cys	Ser	
				340					345					350			
5	CAA	AGT	GAT	TTT	GGG	ACA	ACC	AGC	CTA	AAG	TAT	TTA	GAT	CTG	AGC	TTC	1104
	Gln	Ser	Asp	Phe	Gly	Thr	Thr	Ser	Leu	Lys	Tyr	Leu	Asp	Leu	Ser	Phe	
			355					360					365				
10	AAT	GGT	GTT	ATT	ACC	ATG	AGT	TCA	AAC	TTC	TTG	GGC	TTA	GAA	CAA	CTA	1152
	Asn	Gly	Val	Ile	Thr	Met	Ser	Ser	Asn	Phe	Leu	Gly	Leu	Glu	Gln	Leu	
			370				375					380					
15	GAA	CAT	CTG	GAT	TTC	CAG	CAT	TCC	AAT	TTG	AAA	CAA	ATG	AGT	GAG	TTT	1200
	Glu	His	Leu	Asp	Phe	Gln	His	Ser	Asn	Leu	Lys	Gln	Met	Ser	Glu	Phe	
			385			390					395					400	
	TCA	GTA	TTC	CTA	TCA	CTC	AGA	AAC	CTC	ATT	TAC	CTT	GAC	ATT	TCT	CAT	1248
	Ser	Val	Phe	Leu	Ser	Leu	Arg	Asn	Leu	Ile	Tyr	Leu	Asp	Ile	Ser	His	
					405					410					415		
20	ACT	CAC	ACC	AGA	GTT	GCT	TTC	AAT	GGC	ATC	TTC	AAT	GGC	TTG	TCC	AGT	1296
	Thr	His	Thr	Arg	Val	Ala	Phe	Asn	Gly	Ile	Phe	Asn	Gly	Leu	Ser	Ser	
				420					425					430			
25	CTC	GAA	GTC	TTG	AAA	ATG	GCT	GGC	AAT	TCT	TTC	CAG	GAA	AAC	TTC	CTT	1344
	Leu	Glu	Val	Leu	Lys	Met	Ala	Gly	Asn	Ser	Phe	Gln	Glu	Asn	Phe	Leu	
			435					440					445				
30	CCA	GAT	ATC	TTC	ACA	GAG	CTG	AGA	AAC	TTG	ACC	TTC	CTG	GAC	CTC	TCT	1392
	Pro	Asp	Ile	Phe	Thr	Glu	Leu	Arg	Asn	Leu	Thr	Phe	Leu	Asp	Leu	Ser	
		450					455					460					
35	CAG	TGT	CAA	CTG	GAG	CAG	TTG	TCT	CCA	ACA	GCA	TTT	AAC	TCA	CTC	TCC	1440
	Gln	Cys	Gln	Leu	Glu	Gln	Leu	Ser	Pro	Thr	Ala	Phe	Asn	Ser	Leu	Ser	
						470					475					480	
	AGT	CTT	CAG	GTA	CTA	AAT	ATG	AGC	CAC	AAC	AAC	TTC	TTT	TCA	TTG	GAT	1488
	Ser	Leu	Gln	Val	Leu	Asn	Met	Ser	His	Asn	Asn	Phe	Phe	Ser	Leu	Asp	
					485					490					495		
40	ACG	TTT	CCT	TAT	AAG	TGT	CTG	AAC	TCC	CTC	CAG	GTT	CTT	GAT	TAC	AGT	1536
	Thr	Phe	Pro	Tyr	Lys	Cys	Leu	Asn	Ser	Leu	Gln	Val	Leu	Asp	Tyr	Ser	
				500					505					510			
45	CTC	AAT	CAC	ATA	ATG	ACT	TCC	AAA	AAA	CAG	GAA	CTA	CAG	CAT	TTT	CCA	1584
	Leu	Asn	His	Ile	Met	Thr	Ser	Lys	Lys	Gln	Glu	Leu	Gln	His	Phe	Pro	
			515					520					525				
50	AGT	AGT	CTA	GCT	TTC	TTA	AAT	CTT	ACT	CAG	AAT	GAC	TTT	GCT	TGT	ACT	1632
	Ser	Ser	Leu	Ala	Phe	Leu	Asn	Leu	Thr	Gln	Asn	Asp	Phe	Ala	Cys	Thr	
			530				535					540					
55	TGT	GAA	CAC	CAG	AGT	TTC	CTG	CAA	TGG	ATC	AAG	GAC	CAG	AGG	CAG	CTC	1680
	Cys	Glu	His	Gln	Ser	Phe	Leu	Gln	Trp	Ile	Lys	Asp	Gln	Arg	Gln	Leu	
		545					550				555					560	
	TTG	GTG	GAA	GTT	GAA	CGA	ATG	GAA	TGT	GCA	ACA	CCT	TCA	GAT	AAG	CAG	1728
	Leu	Val	Glu	Val	Glu	Arg	Met	Glu	Cys	Ala	Thr	Pro	Ser	Asp	Lys	Gln	
					565				570						575		
60	GGC	ATG	CCT	GTG	CTG	AGT	TTG	AAT	ATC	ACC	TGT	CAG	ATG	AAT	AAG	ACC	1776
	Gly	Met	Pro	Val	Leu	Ser	Leu	Asn	Ile	Thr	Cys	Gln	Met	Asn	Lys	Thr	

	580	585	590	
5	ATC ATT GGT GTG TCG GTC CTC AGT GTG CTT GTA GTA TCT GTT GTA GCA Ile Ile Gly Val Ser Val Leu Ser Val Leu Val Val Ser Val Val Ala 595 600 605			1824
10	GTT CTG GTC TAT AAG TTC TAT TTT CAC CTG ATG CTT CTT GCT GGC TGC Val Leu Val Tyr Lys Phe Tyr Phe His Leu Met Leu Leu Ala Gly Cys 610 615 620			1872
15	ATA AAG TAT GGT AGA GGT GAA AAC ATC TAT GAT GCC TTT GTT ATC TAC Ile Lys Tyr Gly Arg Gly Glu Asn Ile Tyr Asp Ala Phe Val Ile Tyr 625 630 635 640			1920
20	TCA AGC CAG GAT GAG GAC TGG GTA AGG AAT GAG CTA GTA AAG AAT TTA Ser Ser Gln Asp Glu Asp Trp Val Arg Asn Glu Leu Val Lys Asn Leu 645 650 655			1968
25	GAA GAA GGG GTG CCT CCA TTT CAG CTC TGC CTT CAC TAC AGA GAC TTT Glu Glu Gly Val Pro Pro Phe Gln Leu Cys Leu His Tyr Arg Asp Phe 660 665 670			2016
30	ATT CCC GGT GTG GCC ATT GCT GCC AAC ATC ATC CAT GAA GGT TTC CAT Ile Pro Gly Val Ala Ile Ala Ala Asn Ile Ile His Glu Gly Phe His 675 680 685			2064
35	AAA AGC CGA AAG GTG ATT GTT GTG GTG TCC CAG CAC TTC ATC CAG AGC Lys Ser Arg Lys Val Ile Val Val Val Ser Gln His Phe Ile Gln Ser 690 695 700			2112
40	CGC TGG TGT ATC TTT GAA TAT GAG ATT GCT CAG ACC TGG CAG TTT CTG Arg Trp Cys Ile Phe Glu Tyr Glu Ile Ala Gln Thr Trp Gln Phe Leu 705 710 715 720			2160
45	AGC AGT CGT GCT GGT ATC ATC TTC ATT GTC CTG CAG AAG GTG GAG AAG Ser Ser Arg Ala Gly Ile Ile Phe Ile Val Leu Gln Lys Val Glu Lys 725 730 735			2208
50	ACC CTG CTC AGG CAG CAG GTG GAG CTG TAC CGC CTT CTC AGC AGG AAC Thr Leu Leu Arg Gln Gln Val Glu Leu Tyr Arg Leu Leu Ser Arg Asn 740 745 750			2256
55	ACT TAC CTG GAG TGG GAG GAC AGT GTC CTG GGG CGG CAC ATC TTC TGG Thr Tyr Leu Glu Trp Glu Asp Ser Val Leu Gly Arg His Ile Phe Trp 755 760 765			2304
60	AGA CGA CTC AGA AAA GCC CTG CTG GAT GGT AAA TCA TGG AAT CCA GAA Arg Arg Leu Arg Lys Ala Leu Leu Asp Gly Lys Ser Trp Asn Pro Glu 770 775 780			2352
65	GGA ACA GTG GGT ACA GGA TGC AAT TGG CAG GAA GCA ACA TCT ATC Gly Thr Val Gly Thr Gly Cys Asn Trp Gln Glu Ala Thr Ser Ile 785 790 795			2397
70	TGA			2400

(2) INFORMATION FOR SEQ ID NO:8:

- 60 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 799 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

10	Met	Glu	Leu	Asn	Phe	Tyr	Lys	Ile	Pro	Asp	Asn	Leu	Pro	Phe	Ser	Thr	1	5	10	15
10	Lys	Asn	Leu	Asp	Leu	Ser	Phe	Asn	Pro	Leu	Arg	His	Leu	Gly	Ser	Tyr	20	25	30	
15	Ser	Phe	Phe	Ser	Phe	Pro	Glu	Leu	Gln	Val	Leu	Asp	Leu	Ser	Arg	Cys	35	40	45	
	Glu	Ile	Gln	Thr	Ile	Glu	Asp	Gly	Ala	Tyr	Gln	Ser	Leu	Ser	His	Leu	50	55	60	
20	Ser	Thr	Leu	Ile	Leu	Thr	Gly	Asn	Pro	Ile	Gln	Ser	Leu	Ala	Leu	Gly	65	70	75	80
	Ala	Phe	Ser	Gly	Leu	Ser	Ser	Leu	Gln	Lys	Leu	Val	Ala	Val	Glu	Thr	85	90	95	
25	Asn	Leu	Ala	Ser	Leu	Glu	Asn	Phe	Pro	Ile	Gly	His	Leu	Lys	Thr	Leu	100	105	110	
30	Lys	Glu	Leu	Asn	Val	Ala	His	Asn	Leu	Ile	Gln	Ser	Phe	Lys	Leu	Pro	115	120	125	
	Glu	Tyr	Phe	Ser	Asn	Leu	Thr	Asn	Leu	Glu	His	Leu	Asp	Leu	Ser	Ser	130	135	140	
35	Asn	Lys	Ile	Gln	Ser	Ile	Tyr	Cys	Thr	Asp	Leu	Arg	Val	Leu	His	Gln	145	150	155	160
40	Met	Pro	Leu	Leu	Asn	Leu	Ser	Leu	Asp	Leu	Ser	Leu	Asn	Pro	Met	Asn	165	170	175	
	Phe	Ile	Gln	Pro	Gly	Ala	Phe	Lys	Glu	Ile	Arg	Leu	His	Lys	Leu	Thr	180	185	190	
45	Leu	Arg	Asn	Asn	Phe	Asp	Ser	Leu	Asn	Val	Met	Lys	Thr	Cys	Ile	Gln	195	200	205	
	Gly	Leu	Ala	Gly	Leu	Glu	Val	His	Arg	Leu	Val	Leu	Gly	Glu	Phe	Arg	210	215	220	
50	Asn	Glu	Gly	Asn	Leu	Glu	Lys	Phe	Asp	Lys	Ser	Ala	Leu	Glu	Gly	Leu	225	230	235	240
	Cys	Asn	Leu	Thr	Ile	Glu	Glu	Phe	Arg	Leu	Ala	Tyr	Leu	Asp	Tyr	Tyr	245	250	255	
55	Leu	Asp	Asp	Ile	Ile	Asp	Leu	Phe	Asn	Cys	Leu	Thr	Asn	Val	Ser	Ser	260	265	270	
60	Phe	Ser	Leu	Val	Ser	Val	Thr	Ile	Glu	Arg	Val	Lys	Asp	Phe	Ser	Tyr	275	280	285	

	Asn	Phe	Gly	Trp	Gln	His	Leu	Glu	Leu	Val	Asn	Cys	Lys	Phe	Gly	Gln
	290						295					300				
5	Phe	Pro	Thr	Leu	Lys	Leu	Lys	Ser	Leu	Lys	Arg	Leu	Thr	Phe	Thr	Ser
	305					310					315					320
	Asn	Lys	Gly	Gly	Asn	Ala	Phe	Ser	Glu	Val	Asp	Leu	Pro	Ser	Leu	Glu
					325					330					335	
10	Phe	Leu	Asp	Leu	Ser	Arg	Asn	Gly	Leu	Ser	Phe	Lys	Gly	Cys	Cys	Ser
				340					345					350		
	Gln	Ser	Asp	Phe	Gly	Thr	Thr	Ser	Leu	Lys	Tyr	Leu	Asp	Leu	Ser	Phe
15			355					360					365			
	Asn	Gly	Val	Ile	Thr	Met	Ser	Ser	Asn	Phe	Leu	Gly	Leu	Glu	Gln	Leu
	370						375					380				
20	Glu	His	Leu	Asp	Phe	Gln	His	Ser	Asn	Leu	Lys	Gln	Met	Ser	Glu	Phe
	385					390					395					400
	Ser	Val	Phe	Leu	Ser	Leu	Arg	Asn	Leu	Ile	Tyr	Leu	Asp	Ile	Ser	His
					405					410					415	
25	Thr	His	Thr	Arg	Val	Ala	Phe	Asn	Gly	Ile	Phe	Asn	Gly	Leu	Ser	Ser
				420					425					430		
	Leu	Glu	Val	Leu	Lys	Met	Ala	Gly	Asn	Ser	Phe	Gln	Glu	Asn	Phe	Leu
30			435					440					445			
	Pro	Asp	Ile	Phe	Thr	Glu	Leu	Arg	Asn	Leu	Thr	Phe	Leu	Asp	Leu	Ser
	450						455					460				
35	Gln	Cys	Gln	Leu	Glu	Gln	Leu	Ser	Pro	Thr	Ala	Phe	Asn	Ser	Leu	Ser
	465					470					475					480
	Ser	Leu	Gln	Val	Leu	Asn	Met	Ser	His	Asn	Asn	Phe	Phe	Ser	Leu	Asp
					485					490					495	
40	Thr	Phe	Pro	Tyr	Lys	Cys	Leu	Asn	Ser	Leu	Gln	Val	Leu	Asp	Tyr	Ser
				500					505					510		
	Leu	Asn	His	Ile	Met	Thr	Ser	Lys	Lys	Gln	Glu	Leu	Gln	His	Phe	Pro
45			515					520					525			
	Ser	Ser	Leu	Ala	Phe	Leu	Asn	Leu	Thr	Gln	Asn	Asp	Phe	Ala	Cys	Thr
	530						535					540				
50	Cys	Glu	His	Gln	Ser	Phe	Leu	Gln	Trp	Ile	Lys	Asp	Gln	Arg	Gln	Leu
	545					550					555					560
	Leu	Val	Glu	Val	Glu	Arg	Met	Glu	Cys	Ala	Thr	Pro	Ser	Asp	Lys	Gln
					565				570						575	
55	Gly	Met	Pro	Val	Leu	Ser	Leu	Asn	Ile	Thr	Cys	Gln	Met	Asn	Lys	Thr
				580					585					590		
	Ile	Ile	Gly	Val	Ser	Val	Leu	Ser	Val	Leu	Val	Val	Ser	Val	Val	Ala
60			595					600					605			
	Val	Leu	Val	Tyr	Lys	Phe	Tyr	Phe	His	Leu	Met	Leu	Leu	Ala	Gly	Cys

	610		615		620	
	Ile Lys Tyr Gly Arg Gly Glu Asn Ile Tyr Asp Ala Phe Val Ile Tyr					
	625		630		635	640
5	Ser Ser Gln Asp Glu Asp Trp Val Arg Asn Glu Leu Val Lys Asn Leu					
		645		650		655
	Glu Glu Gly Val Pro Pro Phe Gln Leu Cys Leu His Tyr Arg Asp Phe					
10		660		665		670
	Ile Pro Gly Val Ala Ile Ala Ala Asn Ile Ile His Glu Gly Phe His					
		675		680		685
15	Lys Ser Arg Lys Val Ile Val Val Val Ser Gln His Phe Ile Gln Ser					
		690		695		700
	Arg Trp Cys Ile Phe Glu Tyr Glu Ile Ala Gln Thr Trp Gln Phe Leu					
	705		710		715	720
20	Ser Ser Arg Ala Gly Ile Ile Phe Ile Val Leu Gln Lys Val Glu Lys					
		725		730		735
	Thr Leu Leu Arg Gln Gln Val Glu Leu Tyr Arg Leu Leu Ser Arg Asn					
25		740		745		750
	Thr Tyr Leu Glu Trp Glu Asp Ser Val Leu Gly Arg His Ile Phe Trp					
		755		760		765
30	Arg Arg Leu Arg Lys Ala Leu Leu Asp Gly Lys Ser Trp Asn Pro Glu					
		770		775		780
	Gly Thr Val Gly Thr Gly Cys Asn Trp Gln Glu Ala Thr Ser Ile					
	785		790		795	

35

(2) INFORMATION FOR SEQ ID NO:9:

	(i) SEQUENCE CHARACTERISTICS:
	(A) LENGTH: 1275 base pairs
40	(B) TYPE: nucleic acid
	(C) STRANDEDNESS: single
	(D) TOPOLOGY: linear

45	(ii) MOLECULE TYPE: cDNA
----	--------------------------

	(ix) FEATURE:
	(A) NAME/KEY: CDS
50	(B) LOCATION: 1..1095

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

	TGT TGG GAT GTT TTT GAG GGA CTT TCT CAT CTT CAA GTT CTG TAT TTG	48
55	Cys Trp Asp Val Phe Glu Gly Leu Ser His Leu Gln Val Leu Tyr Leu	
	1 5 10 15	
	AAT CAT AAC TAT CTT AAT TCC CTT CCA CCA GGA GTA TTT AGC CAT CTG	96
60	Asn His Asn Tyr Leu Asn Ser Leu Pro Gly Val Phe Ser His Leu	
	20 25 30	

	ACT	GCA	TTA	AGG	GGA	CTA	AGC	CTC	AAC	TCC	AAC	AGG	CTG	ACA	GTT	CTT	144
	Thr	Ala	Leu	Arg	Gly	Leu	Ser	Leu	Asn	Ser	Asn	Arg	Leu	Thr	Val	Leu	
			35					40				45					
5	TCT	CAC	AAT	GAT	TTA	CCT	GCT	AAT	TTA	GAG	ATC	CTG	GAC	ATA	TCC	AGG	192
	Ser	His	Asn	Asp	Leu	Pro	Ala	Asn	Leu	Glu	Ile	Leu	Asp	Ile	Ser	Arg	
		50					55					60					
10	AAC	CAG	CTC	CTA	GCT	CCT	AAT	CCT	GAT	GTA	TTT	GTA	TCA	CTT	AGT	GTC	240
	Asn	Gln	Leu	Leu	Ala	Pro	Asn	Pro	Asp	Val	Phe	Val	Ser	Leu	Ser	Val	
		65				70					75					80	
15	TTG	GAT	ATA	ACT	CAT	AAC	AAG	TTC	ATT	TGT	GAA	TGT	GAA	CTT	AGC	ACT	288
	Leu	Asp	Ile	Thr	His	Asn	Lys	Phe	Ile	Cys	Glu	Cys	Glu	Leu	Ser	Thr	
				85						90					95		
20	TTT	ATC	AAT	TGG	CTT	AAT	CAC	ACC	AAT	GTC	ACT	ATA	GCT	GGG	CCT	CCT	336
	Phe	Ile	Asn	Trp	Leu	Asn	His	Thr	Asn	Val	Thr	Ile	Ala	Gly	Pro	Pro	
				100					105					110			
25	GCA	GAC	ATA	TAT	TGT	GTG	TAC	CCT	GAC	TCG	TTC	TCT	GGG	GTT	TCC	CTC	384
	Ala	Asp	Ile	Tyr	Cys	Val	Tyr	Pro	Asp	Ser	Phe	Ser	Gly	Val	Ser	Leu	
			115					120					125				
30	TTC	TCT	CTT	TCC	ACG	GAA	GGT	TGT	GAT	GAA	GAG	GAA	GTC	TTA	AAG	TCC	432
	Phe	Ser	Leu	Ser	Thr	Glu	Gly	Cys	Asp	Glu	Glu	Glu	Val	Leu	Lys	Ser	
		130					135					140					
35	CTA	AAG	TTC	TCC	CTT	TTC	ATT	GTA	TGC	ACT	GTC	ACT	CTG	ACT	CTG	TTC	480
	Leu	Lys	Phe	Ser	Leu	Phe	Ile	Val	Cys	Thr	Val	Thr	Leu	Thr	Leu	Phe	
		145				150					155					160	
40	CTC	ATG	ACC	ATC	CTC	ACA	GTC	ACA	AAG	TTC	CGG	GGC	TTC	TGT	TTT	ATC	528
	Leu	Met	Thr	Ile	Leu	Thr	Val	Thr	Lys	Phe	Arg	Gly	Phe	Cys	Phe	Ile	
				165						170					175		
45	TGT	TAT	AAG	ACA	GCC	CAG	AGA	CTG	GTG	TTC	AAG	GAC	CAT	CCC	CAG	GGC	576
	Cys	Tyr	Lys	Thr	Ala	Gln	Arg	Leu	Val	Phe	Lys	Asp	His	Pro	Gln	Gly	
			180						185					190			
50	ACA	GAA	CCT	GAT	ATG	TAC	AAA	TAT	GAT	GCC	TAT	TTG	TGC	TTC	AGC	AGC	624
	Thr	Glu	Pro	Asp	Met	Tyr	Lys	Tyr	Asp	Ala	Tyr	Leu	Cys	Phe	Ser	Ser	
			195					200					205				
55	AAA	GAC	TTC	ACA	TGG	GTG	CAG	AAT	GCT	TTG	CTC	AAA	CAC	CTG	GAC	ACT	672
	Lys	Asp	Phe	Thr	Trp	Val	Gln	Asn	Ala	Leu	Leu	Lys	His	Leu	Asp	Thr	
		210					215					220					
60	CAA	TAC	AGT	GAC	CAA	AAC	AGA	TTC	AAC	CTG	TGC	TTT	GAA	GAA	AGA	GAC	720
	Gln	Tyr	Ser	Asp	Gln	Asn	Arg	Phe	Asn	Leu	Cys	Phe	Glu	Glu	Arg	Asp	
		225				230					235					240	
65	TTT	GTC	CCA	GGA	GAA	AAC	CGC	ATT	GCC	AAT	ATC	CAG	GAT	GCC	ATC	TGG	768
	Phe	Val	Pro	Gly	Glu	Asn	Arg	Ile	Ala	Asn	Ile	Gln	Asp	Ala	Ile	Trp	
				245						250					255		
70	AAC	AGT	AGA	AAG	ATC	GTT	TGT	CTT	GTG	AGC	AGA	CAC	TTC	CTT	AGA	GAT	816
	Asn	Ser	Arg	Lys	Ile	Val	Cys	Leu	Val	Ser	Arg	His	Phe	Leu	Arg	Asp	
			260						265					270			
75	GGC	TGG	TGC	CTT	GAA	GCC	TTC	AGT	TAT	GCC	CAG	GGC	AGG	TGC	TTA	TCT	864

	Gly Trp Cys Leu Glu Ala Phe Ser Tyr Ala Gln Gly Arg Cys Leu Ser	
	275 280 285	
5	GAC CTT AAC AGT GCT CTC ATC ATG GTG GTG GTT GGG TCC TTG TCC CAG Asp Leu Asn Ser Ala Leu Ile Met Val Val Val Gly Ser Leu Ser Gln	912
	290 295 300	
10	TAC CAG TTG ATG AAA CAT CAA TCC ATC AGA GGC TTT GTA CAG AAA CAG Tyr Gln Leu Met Lys His Gln Ser Ile Arg Gly Phe Val Gln Lys Gln	960
	305 310 315 320	
15	CAG TAT TTG AGG TGG CCT GAG GAT CTC CAG GAT GTT GGC TGG TTT CTT Gln Tyr Leu Arg Trp Pro Glu Asp Leu Gln Asp Val Gly Trp Phe Leu	1008
	325 330 335	
20	CAT AAA CTC TCT CAA CAG ATA CTA AAG AAA GAA AAG GAA AAG AAG AAA His Lys Leu Ser Gln Gln Ile Leu Lys Lys Glu Lys Glu Lys Lys Lys	1056
	340 345 350	
25	GAC AAT AAC ATT CCG TTG CAA ACT GTA GCA ACC ATC TCC TAATCAAAGG Asp Asn Asn Ile Pro Leu Gln Thr Val Ala Thr Ile Ser	1105
	355 360 365	
30	AGCAATTTCC AACTTATCTC AAGCCACAAA TAACTCTTCA CTTTGTATTT GCACCAAGTT	1165
	ATCATTTTGG GGTCTCTCTT GGAGGTTTTT TTTTCTTTT TGCTACTATG AAAACAACAT	1225
	AAATCTCTCA ATTTTCGTAT CAAAAA AAAAAA TGGCGGCCGC	1275
	(2) INFORMATION FOR SEQ ID NO:10:	
	(i) SEQUENCE CHARACTERISTICS:	
35	(A) LENGTH: 365 amino acids	
	(B) TYPE: amino acid	
	(D) TOPOLOGY: linear	
	(ii) MOLECULE TYPE: protein	
40	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:	
	Cys Trp Asp Val Phe Glu Gly Leu Ser His Leu Gln Val Leu Tyr Leu	
	1 5 10 15	
45	Asn His Asn Tyr Leu Asn Ser Leu Pro Pro Gly Val Phe Ser His Leu	
	20 25 30	
	Thr Ala Leu Arg Gly Leu Ser Leu Asn Ser Asn Arg Leu Thr Val Leu	
50	35 40 45	
	Ser His Asn Asp Leu Pro Ala Asn Leu Glu Ile Leu Asp Ile Ser Arg	
	50 55 60	
55	Asn Gln Leu Leu Ala Pro Asn Pro Asp Val Phe Val Ser Leu Ser Val	
	65 70 75 80	
	Leu Asp Ile Thr His Asn Lys Phe Ile Cys Glu Cys Glu Leu Ser Thr	
	85 90 95	
60	Phe Ile Asn Trp Leu Asn His Thr Asn Val Thr Ile Ala Gly Pro Pro	
	100 105 110	

Ala Asp Ile Tyr Cys Val Tyr Pro Asp Ser Phe Ser Gly Val Ser Leu
115 120 125

5 Phe Ser Leu Ser Thr Glu Gly Cys Asp Glu Glu Glu Val Leu Lys Ser
130 135 140

Leu Lys Phe Ser Leu Phe Ile Val Cys Thr Val Thr Leu Thr Leu Phe
145 150 155 160

10 Leu Met Thr Ile Leu Thr Val Thr Lys Phe Arg Gly Phe Cys Phe Ile
165 170 175

Cys Tyr Lys Thr Ala Gln Arg Leu Val Phe Lys Asp His Pro Gln Gly
180 185 190

15 Thr Glu Pro Asp Met Tyr Lys Tyr Asp Ala Tyr Leu Cys Phe Ser Ser
195 200 205

20 Lys Asp Phe Thr Trp Val Gln Asn Ala Leu Leu Lys His Leu Asp Thr
210 215 220

Gln Tyr Ser Asp Gln Asn Arg Phe Asn Leu Cys Phe Glu Glu Arg Asp
225 230 235 240

25 Phe Val Pro Gly Glu Asn Arg Ile Ala Asn Ile Gln Asp Ala Ile Trp
245 250 255

Asn Ser Arg Lys Ile Val Cys Leu Val Ser Arg His Phe Leu Arg Asp
260 265 270

30 Gly Trp Cys Leu Glu Ala Phe Ser Tyr Ala Gln Gly Arg Cys Leu Ser
275 280 285

35 Asp Leu Asn Ser Ala Leu Ile Met Val Val Val Gly Ser Leu Ser Gln
290 295 300

Tyr Gln Leu Met Lys His Gln Ser Ile Arg Gly Phe Val Gln Lys Gln
305 310 315 320

40 Gln Tyr Leu Arg Trp Pro Glu Asp Leu Gln Asp Val Gly Trp Phe Leu
325 330 335

His Lys Leu Ser Gln Gln Ile Leu Lys Lys Glu Lys Glu Lys Lys Lys
340 345 350

45 Asp Asn Asn Ile Pro Leu Gln Thr Val Ala Thr Ile Ser
355 360 365

50 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 3138 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..3135

(ix) FEATURE:

5 (A) NAME/KEY: mat_peptide
(B) LOCATION: 67..3135

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

10	ATG TGG ACA CTG AAG AGA CTA ATT CTT ATC CTT TTT AAC ATA ATC CTA	48
	Met Trp Thr Leu Lys Arg Leu Ile Leu Ile Leu Phe Asn Ile Ile Leu	
	-22 -20 -15 -10	
15	ATT TCC AAA CTC CTT GGG GCT AGA TGG TTT CCT AAA ACT CTG CCC TGT	96
	Ile Ser Lys Leu Leu Gly Ala Arg Trp Phe Pro Lys Thr Leu Pro Cys	
	-5 1 5 10	
20	GAT GTC ACT CTG GAT GTT CCA AAG AAC CAT GTG ATC GTG GAC TGC ACA	144
	Asp Val Thr Leu Asp Val Pro Lys Asn His Val Ile Val Asp Cys Thr	
	15 20 25	
25	GAC AAG CAT TTG ACA GAA ATT CCT GGA GGT ATT CCC ACG AAC ACC ACG	192
	Asp Lys His Leu Thr Glu Ile Pro Gly Gly Ile Pro Thr Asn Thr Thr	
	30 35 40	
30	AAC CTC ACC CTC ACC ATT AAC CAC ATA CCA GAC ATC TCC CCA GCG TCC	240
	Asn Leu Thr Leu Thr Ile Asn His Ile Pro Asp Ile Ser Pro Ala Ser	
	45 50 55	
35	TTT CAC AGA CTG GAC CAT CTG GTA GAG ATC GAT TTC AGA TGC AAC TGT	288
	Phe His Arg Leu Asp His Leu Val Glu Ile Asp Phe Arg Cys Asn Cys	
	60 65 70	
40	GTA CCT ATT CCA CTG GGG TCA AAA AAC AAC ATG TGC ATC AAG AGG CTG	336
	Val Pro Ile Pro Leu Gly Ser Lys Asn Asn Met Cys Ile Lys Arg Leu	
	75 80 85 90	
45	CAG ATT AAA CCC AGA AGC TTT AGT GGA CTC ACT TAT TTA AAA TCC CTT	384
	Gln Ile Lys Pro Arg Ser Phe Ser Gly Leu Thr Tyr Leu Lys Ser Leu	
	95 100 105	
50	TAC CTG GAT GGA AAC CAG CTA CTA GAG ATA CCG CAG GGC CTC CCG CCT	432
	Tyr Leu Asp Gly Asn Gln Leu Leu Glu Ile Pro Gln Gly Leu Pro Pro	
	110 115 120	
55	AGC TTA CAG CTT CTC AGC CTT GAG GCC AAC AAC ATC TTT TCC ATC AGA	480
	Ser Leu Gln Leu Leu Ser Leu Glu Ala Asn Asn Ile Phe Ser Ile Arg	
	125 130 135	
60	AAA GAG AAT CTA ACA GAA CTG GCC AAC ATA GAA ATA CTC TAC CTG GGC	528
	Lys Glu Asn Leu Thr Glu Leu Ala Asn Ile Glu Ile Leu Tyr Leu Gly	
	140 145 150	
65	CAA AAC TGT TAT TAT CGA AAT CCT TGT TAT GTT TCA TAT TCA ATA GAG	576
	Gln Asn Cys Tyr Tyr Arg Asn Pro Cys Tyr Val Ser Tyr Ser Ile Glu	
	155 160 165 170	
70	AAA GAT GCC TTC CTA AAC TTG ACA AAG TTA AAA GTG CTC TCC CTG AAA	624
	Lys Asp Ala Phe Leu Asn Leu Thr Lys Leu Lys Val Leu Ser Leu Lys	
	175 180 185	

5	GAT AAC AAT GTC ACA GCC GTC CCT ACT GTT TTG CCA TCT ACT TTA ACA	672
	Asp Asn Asn Val Thr Ala Val Pro Thr Val Leu Pro Ser Thr Leu Thr	
10	GAA CTA TAT CTC TAC AAC AAC ATG ATT GCA AAA ATC CAA GAA GAT GAT	720
	Glu Leu Tyr Leu Tyr Asn Asn Met Ile Ala Lys Ile Gln Glu Asp Asp	
15	TTT AAT AAC CTC AAC CAA TTA CAA ATT CTT GAC CTA AGT GGA AAT TGC	768
	Phe Asn Asn Leu Asn Gln Leu Gln Ile Leu Asp Leu Ser Gly Asn Cys	
20	CCT CGT TGT TAT AAT GCC CCA TTT CCT TGT GCG CCG TGT AAA AAT AAT	816
	Pro Arg Cys Tyr Asn Ala Pro Phe Pro Cys Ala Pro Cys Lys Asn Asn	
25	TCT CCC CTA CAG ATC CCT GTA AAT GCT TTT GAT GCG CTG ACA GAA TTA	864
	Ser Pro Leu Gln Ile Pro Val Asn Ala Phe Asp Ala Leu Thr Glu Leu	
30	AAA GTT TTA CGT CTA CAC AGT AAC TCT CTT CAG CAT GTG CCC CCA AGA	912
	Lys Val Leu Arg Leu His Ser Asn Ser Leu Gln His Val Pro Pro Arg	
35	TGG TTT AAG AAC ATC AAC AAA CTC CAG GAA CTG GAT CTG TCC CAA AAC	960
	Trp Phe Lys Asn Ile Asn Lys Leu Gln Glu Leu Asp Leu Ser Gln Asn	
40	TTC TTG GCC AAA GAA ATT GGG GAT GCT AAA TTT CTG CAT TTT CTC CCC	1008
	Phe Leu Ala Lys Glu Ile Gly Asp Ala Lys Phe Leu His Phe Leu Pro	
45	AGC CTC ATC CAA TTG GAT CTG TCT TTC AAT TTT GAA CTT CAG GTC TAT	1056
	Ser Leu Ile Gln Leu Asp Leu Ser Phe Asn Phe Glu Leu Gln Val Tyr	
50	CGT GCA TCT ATG AAT CTA TCA CAA GCA TTT TCT TCA CTG AAA AGC CTG	1104
	Arg Ala Ser Met Asn Leu Ser Gln Ala Phe Ser Ser Leu Lys Ser Leu	
55	AAA ATT CTG CGG ATC AGA GGA TAT GTC TTT AAA GAG TTG AAA AGC TTT	1152
	Lys Ile Leu Arg Ile Arg Gly Tyr Val Phe Lys Glu Leu Lys Ser Phe	
60	AAC CTC TCG CCA TTA CAT AAT CTT CAA AAT CTT GAA GTT CTT GAT CTT	1200
	Asn Leu Ser Pro Leu His Asn Leu Gln Asn Leu Glu Val Leu Asp Leu	
65	GGC ACT AAC TTT ATA AAA ATT GCT AAC CTC AGC ATG TTT AAA CAA TTT	1248
	Gly Thr Asn Phe Ile Lys Ile Ala Asn Leu Ser Met Phe Lys Gln Phe	
70	AAA AGA CTG AAA GTC ATA GAT CTT TCA GTG AAT AAA ATA TCA CCT TCA	1296
	Lys Arg Leu Lys Val Ile Asp Leu Ser Val Asn Lys Ile Ser Pro Ser	
75	GGA GAT TCA AGT GAA GTT GGC TTC TGC TCA AAT GCC AGA ACT TCT GTA	1344
	Gly Asp Ser Ser Glu Val Gly Phe Cys Ser Asn Ala Arg Thr Ser Val	

	GAA AGT TAT GAA CCC CAG GTC CTG GAA CAA TTA CAT TAT TTC AGA TAT	1392
	Glu Ser Tyr Glu Pro Gln Val Leu Glu Gln Leu His Tyr Phe Arg Tyr	
	430 435 440	
5	GAT AAG TAT GCA AGG AGT TGC AGA TTC AAA AAC AAA GAG GCT TCT TTC	1440
	Asp Lys Tyr Ala Arg Ser Cys Arg Phe Lys Asn Lys Glu Ala Ser Phe	
	445 450 455	
10	ATG TCT GTT AAT GAA AGC TGC TAC AAG TAT GGG CAG ACC TTG GAT CTA	1488
	Met Ser Val Asn Glu Ser Cys Tyr Lys Tyr Gly Gln Thr Leu Asp Leu	
	460 465 470	
15	AGT AAA AAT AGT ATA TTT TTT GTC AAG TCC TCT GAT TTT CAG CAT CTT	1536
	Ser Lys Asn Ser Ile Phe Phe Val Lys Ser Ser Asp Phe Gln His Leu	
	475 480 485 490	
20	TCT TTC CTC AAA TGC CTG AAT CTG TCA GGA AAT CTC ATT AGC CAA ACT	1584
	Ser Phe Leu Lys Cys Leu Asn Leu Ser Gly Asn Leu Ile Ser Gln Thr	
	495 500 505	
	CTT AAT GGC AGT GAA TTC CAA CCT TTA GCA GAG CTG AGA TAT TTG GAC	1632
	Leu Asn Gly Ser Glu Phe Gln Pro Leu Ala Glu Leu Arg Tyr Leu Asp	
	510 515 520	
25	TTC TCC AAC AAC CGG CTT GAT TTA CTC CAT TCA ACA GCA TTT GAA GAG	1680
	Phe Ser Asn Asn Arg Leu Asp Leu Leu His Ser Thr Ala Phe Glu Glu	
	525 530 535	
30	CTT CAC AAA CTG GAA GTT CTG GAT ATA AGC AGT AAT AGC CAT TAT TTT	1728
	Leu His Lys Leu Glu Val Leu Asp Ile Ser Ser Asn Ser His Tyr Phe	
	540 545 550	
35	CAA TCA GAA GGA ATT ACT CAT ATG CTA AAC TTT ACC AAG AAC CTA AAG	1776
	Gln Ser Glu Gly Ile Thr His Met Leu Asn Phe Thr Lys Asn Leu Lys	
	555 560 565 570	
40	GTT CTG CAG AAA CTG ATG ATG AAC GAC AAT GAC ATC TCT TCC TCC ACC	1824
	Val Leu Gln Lys Leu Met Met Asn Asp Asn Asp Ile Ser Ser Ser Thr	
	575 580 585	
	AGC AGG ACC ATG GAG AGT GAG TCT CTT AGA ACT CTG GAA TTC AGA GGA	1872
	Ser Arg Thr Met Glu Ser Glu Ser Leu Arg Thr Leu Glu Phe Arg Gly	
	590 595 600	
45	AAT CAC TTA GAT GTT TTA TGG AGA GAA GGT GAT AAC AGA TAC TTA CAA	1920
	Asn His Leu Asp Val Leu Trp Arg Glu Gly Asp Asn Arg Tyr Leu Gln	
	605 610 615	
50	TTA TTC AAG AAT CTG CTA AAA TTA GAG GAA TTA GAC ATC TCT AAA AAT	1968
	Leu Phe Lys Asn Leu Leu Lys Leu Glu Glu Leu Asp Ile Ser Lys Asn	
	620 625 630	
55	TCC CTA AGT TTC TTG CCT TCT GGA GTT TTT GAT GGT ATG CCT CCA AAT	2016
	Ser Leu Ser Phe Leu Pro Ser Gly Val Phe Asp Gly Met Pro Pro Asn	
	635 640 645 650	
60	CTA AAG AAT CTC TCT TTG GCC AAA AAT GGG CTC AAA TCT TTC AGT TGG	2064
	Leu Lys Asn Leu Ser Leu Ala Lys Asn Gly Leu Lys Ser Phe Ser Trp	
	655 660 665	
	AAG AAA CTC CAG TGT CTA AAG AAC CTG GAA ACT TTG GAC CTC AGC CAC	2112

		Lys	Lys	Leu	Gln	Cys	Leu	Lys	Asn	Leu	Glu	Thr	Leu	Asp	Leu	Ser	His	
					670					675					680			
5		AAC	CAA	CTG	ACC	ACT	GTC	CCT	GAG	AGA	TTA	TCC	AAC	TGT	TCC	AGA	AGC	2160
		Asn	Gln	Leu	Thr	Thr	Val	Pro	Glu	Arg	Leu	Ser	Asn	Cys	Ser	Arg	Ser	
				685					690					695				
10		CTC	AAG	AAT	CTG	ATT	CTT	AAG	AAT	AAT	CAA	ATC	AGG	AGT	CTG	ACG	AAG	2208
		Leu	Lys	Asn	Leu	Ile	Leu	Lys	Asn	Asn	Gln	Ile	Arg	Ser	Leu	Thr	Lys	
			700					705					710					
15		TAT	TTT	CTA	CAA	GAT	GCC	TTC	CAG	TTG	CGA	TAT	CTG	GAT	CTC	AGC	TCA	2256
		Tyr	Phe	Leu	Gln	Asp	Ala	Phe	Gln	Leu	Arg	Tyr	Leu	Asp	Leu	Ser	Ser	
		715					720					725					730	
		AAT	AAA	ATC	CAG	ATG	ATC	CAA	AAG	ACC	AGC	TTC	CCA	GAA	AAT	GTC	CTC	2304
		Asn	Lys	Ile	Gln	Met	Ile	Gln	Lys	Thr	Ser	Phe	Pro	Glu	Asn	Val	Leu	
						735					740					745		
20		AAC	AAT	CTG	AAG	ATG	TTG	CTT	TTG	CAT	CAT	AAT	CGG	TTT	CTG	TGC	ACC	2352
		Asn	Asn	Leu	Lys	Met	Leu	Leu	Leu	His	His	Asn	Arg	Phe	Leu	Cys	Thr	
					750					755					760			
25		TGT	GAT	GCT	GTG	TGG	TTT	GTC	TGG	TGG	GTT	AAC	CAT	ACG	GAG	GTG	ACT	2400
		Cys	Asp	Ala	Val	Trp	Phe	Val	Trp	Trp	Val	Asn	His	Thr	Glu	Val	Thr	
				765					770					775				
30		ATT	CCT	TAC	CTG	GCC	ACA	GAT	GTG	ACT	TGT	GTG	GGG	CCA	GGA	GCA	CAC	2448
		Ile	Pro	Tyr	Leu	Ala	Thr	Asp	Val	Thr	Cys	Val	Gly	Pro	Gly	Ala	His	
			780					785					790					
35		AAG	GGC	CAA	AGT	GTG	ATC	TCC	CTG	GAT	CTG	TAC	ACC	TGT	GAG	TTA	GAT	2496
		Lys	Gly	Gln	Ser	Val	Ile	Ser	Leu	Asp	Leu	Tyr	Thr	Cys	Glu	Leu	Asp	
		795					800					805					810	
		CTG	ACT	AAC	CTG	ATT	CTG	TTC	TCA	CTT	TCC	ATA	TCT	GTA	TCT	CTC	TTT	2544
		Leu	Thr	Asn	Leu	Ile	Leu	Phe	Ser	Leu	Ser	Ile	Ser	Val	Ser	Leu	Phe	
						815					820					825		
40		CTC	ATG	GTG	ATG	ATG	ACA	GCA	AGT	CAC	CTC	TAT	TTC	TGG	GAT	GTG	TGG	2592
		Leu	Met	Val	Met	Met	Thr	Ala	Ser	His	Leu	Tyr	Phe	Trp	Asp	Val	Trp	
					830					835					840			
45		TAT	ATT	TAC	CAT	TTC	TGT	AAG	GCC	AAG	ATA	AAG	GGG	TAT	CAG	CGT	CTA	2640
		Tyr	Ile	Tyr	His	Phe	Cys	Lys	Ala	Lys	Ile	Lys	Gly	Tyr	Gln	Arg	Leu	
				845					850					855				
50		ATA	TCA	CCA	GAC	TGT	TGC	TAT	GAT	GCT	TTT	ATT	GTG	TAT	GAC	ACT	AAA	2688
		Ile	Ser	Pro	Asp	Cys	Cys	Tyr	Asp	Ala	Phe	Ile	Val	Tyr	Asp	Thr	Lys	
			860					865					870					
55		GAC	CCA	GCT	GTG	ACC	GAG	TGG	GTT	TTG	GCT	GAG	CTG	GTG	GCC	AAA	CTG	2736
		Asp	Pro	Ala	Val	Thr	Glu	Trp	Val	Leu	Ala	Glu	Leu	Val	Ala	Lys	Leu	
		875					880					885					890	
		GAA	GAC	CCA	AGA	GAG	AAA	CAT	TTT	AAT	TTA	TGT	CTC	GAG	GAA	AGG	GAC	2784
		Glu	Asp	Pro	Arg	Glu	Lys	His	Phe	Asn	Leu	Cys	Leu	Glu	Glu	Arg	Asp	
						895				900						905		
60		TGG	TTA	CCA	GGG	CAG	CCA	GTT	CTG	GAA	AAC	CTT	TCC	CAG	AGC	ATA	CAG	2832
		Trp	Leu	Pro	Gly	Gln	Pro	Val	Leu	Glu	Asn	Leu	Ser	Gln	Ser	Ile	Gln	

	910	915	920	
5	CTT AGC AAA AAG ACA GTG TTT GTG ATG ACA GAC AAG TAT GCA AAG ACT Leu Ser Lys Lys Thr Val Phe Val Met Thr Asp Lys Tyr Ala Lys Thr 925 930 935	2880		
10	GAA AAT TTT AAG ATA GCA TTT TAC TTG TCC CAT CAG AGG CTC ATG GAT Glu Asn Phe Lys Ile Ala Phe Tyr Leu Ser His Gln Arg Leu Met Asp 940 945 950	2928		
15	GAA AAA GTT GAT GTG ATT ATC TTG ATA TTT CTT GAG AAG CCC TTT CAG Glu Lys Val Asp Val Ile Ile Leu Ile Phe Leu Glu Lys Pro Phe Gln 955 960 965 970	2976		
20	AAG TCC AAG TTC CTC CAG CTC CGG AAA AGG CTC TGT GGG AGT TCT GTC Lys Ser Lys Phe Leu Gln Leu Arg Lys Arg Leu Cys Gly Ser Ser Val 975 980 985	3024		
25	CTT GAG TGG CCA ACA AAC CCG CAA GCT CAC CCA TAC TTC TGG CAG TGT Leu Glu Trp Pro Thr Asn Pro Gln Ala His Pro Tyr Phe Trp Gln Cys 990 995 1000	3072		
30	CTA AAG AAC GCC CTG GCC ACA GAC AAT CAT GTG GCC TAT AGT CAG GTG Leu Lys Asn Ala Leu Ala Thr Asp Asn His Val Ala Tyr Ser Gln Val 1005 1010 1015	3120		
35	TTC AAG GAA ACG GTC TAG Phe Lys Glu Thr Val 1020	3138		

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1045 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met Trp Thr Leu Lys Arg Leu Ile Leu Ile Leu Phe Asn Ile Ile Leu -22 -20 -15 -10	
Ile Ser Lys Leu Leu Gly Ala Arg Trp Phe Pro Lys Thr Leu Pro Cys -5 1 5 10	
Asp Val Thr Leu Asp Val Pro Lys Asn His Val Ile Val Asp Cys Thr 15 20 25	
Asp Lys His Leu Thr Glu Ile Pro Gly Gly Ile Pro Thr Asn Thr Thr 30 35 40	
Asn Leu Thr Leu Thr Ile Asn His Ile Pro Asp Ile Ser Pro Ala Ser 45 50 55	
Phe His Arg Leu Asp His Leu Val Glu Ile Asp Phe Arg Cys Asn Cys 60 65 70	
Val Pro Ile Pro Leu Gly Ser Lys Asn Asn Met Cys Ile Lys Arg Leu	

*

	75		80		85		90
	Gln Ile Lys Pro Arg Ser Phe Ser Gly Leu Thr Tyr Leu Lys Ser Leu						
		95			100		105
5	Tyr Leu Asp Gly Asn Gln Leu Leu Glu Ile Pro Gln Gly Leu Pro Pro						
		110			115		120
10	Ser Leu Gln Leu Leu Ser Leu Glu Ala Asn Asn Ile Phe Ser Ile Arg						
		125			130		135
	Lys Glu Asn Leu Thr Glu Leu Ala Asn Ile Glu Ile Leu Tyr Leu Gly						
		140			145		150
15	Gln Asn Cys Tyr Tyr Arg Asn Pro Cys Tyr Val Ser Tyr Ser Ile Glu						
		155			160		165
	Lys Asp Ala Phe Leu Asn Leu Thr Lys Leu Lys Val Leu Ser Leu Lys						
		175			180		185
20	Asp Asn Asn Val Thr Ala Val Pro Thr Val Leu Pro Ser Thr Leu Thr						
		190			195		200
25	Glu Leu Tyr Leu Tyr Asn Asn Met Ile Ala Lys Ile Gln Glu Asp Asp						
		205			210		215
	Phe Asn Asn Leu Asn Gln Leu Gln Ile Leu Asp Leu Ser Gly Asn Cys						
		220			225		230
30	Pro Arg Cys Tyr Asn Ala Pro Phe Pro Cys Ala Pro Cys Lys Asn Asn						
		235			240		245
	Ser Pro Leu Gln Ile Pro Val Asn Ala Phe Asp Ala Leu Thr Glu Leu						
		255			260		265
35	Lys Val Leu Arg Leu His Ser Asn Ser Leu Gln His Val Pro Pro Arg						
		270			275		280
40	Trp Phe Lys Asn Ile Asn Lys Leu Gln Glu Leu Asp Leu Ser Gln Asn						
		285			290		295
	Phe Leu Ala Lys Glu Ile Gly Asp Ala Lys Phe Leu His Phe Leu Pro						
		300			305		310
45	Ser Leu Ile Gln Leu Asp Leu Ser Phe Asn Phe Glu Leu Gln Val Tyr						
		315			320		325
	Arg Ala Ser Met Asn Leu Ser Gln Ala Phe Ser Ser Leu Lys Ser Leu						
		335			340		345
50	Lys Ile Leu Arg Ile Arg Gly Tyr Val Phe Lys Glu Leu Lys Ser Phe						
		350			355		360
55	Asn Leu Ser Pro Leu His Asn Leu Gln Asn Leu Glu Val Leu Asp Leu						
		365			370		375
	Gly Thr Asn Phe Ile Lys Ile Ala Asn Leu Ser Met Phe Lys Gln Phe						
		380			385		390
60	Lys Arg Leu Lys Val Ile Asp Leu Ser Val Asn Lys Ile Ser Pro Ser						
		395			400		405
							410

	Gly	Asp	Ser	Ser	Glu	Val	Gly	Phe	Cys	Ser	Asn	Ala	Arg	Thr	Ser	Val	
					415					420					425		
5	Glu	Ser	Tyr	Glu	Pro	Gln	Val	Leu	Glu	Gln	Leu	His	Tyr	Phe	Arg	Tyr	
				430					435					440			
	Asp	Lys	Tyr	Ala	Arg	Ser	Cys	Arg	Phe	Lys	Asn	Lys	Glu	Ala	Ser	Phe	
			445					450					455				
10	Met	Ser	Val	Asn	Glu	Ser	Cys	Tyr	Lys	Tyr	Gly	Gln	Thr	Leu	Asp	Leu	
		460					465					470					
	Ser	Lys	Asn	Ser	Ile	Phe	Phe	Val	Lys	Ser	Ser	Asp	Phe	Gln	His	Leu	
15		475				480					485					490	
	Ser	Phe	Leu	Lys	Cys	Leu	Asn	Leu	Ser	Gly	Asn	Leu	Ile	Ser	Gln	Thr	
				495						500					505		
20	Leu	Asn	Gly	Ser	Glu	Phe	Gln	Pro	Leu	Ala	Glu	Leu	Arg	Tyr	Leu	Asp	
				510					515					520			
	Phe	Ser	Asn	Asn	Arg	Leu	Asp	Leu	Leu	His	Ser	Thr	Ala	Phe	Glu	Glu	
			525					530					535				
25	Leu	His	Lys	Leu	Glu	Val	Leu	Asp	Ile	Ser	Ser	Asn	Ser	His	Tyr	Phe	
		540					545					550					
	Gln	Ser	Glu	Gly	Ile	Thr	His	Met	Leu	Asn	Phe	Thr	Lys	Asn	Leu	Lys	
30		555				560					565					570	
	Val	Leu	Gln	Lys	Leu	Met	Met	Asn	Asp	Asn	Asp	Ile	Ser	Ser	Ser	Thr	
				575					580						585		
35	Ser	Arg	Thr	Met	Glu	Ser	Glu	Ser	Leu	Arg	Thr	Leu	Glu	Phe	Arg	Gly	
				590					595					600			
	Asn	His	Leu	Asp	Val	Leu	Trp	Arg	Glu	Gly	Asp	Asn	Arg	Tyr	Leu	Gln	
			605					610					615				
40	Leu	Phe	Lys	Asn	Leu	Leu	Lys	Leu	Glu	Glu	Leu	Asp	Ile	Ser	Lys	Asn	
		620					625					630					
	Ser	Leu	Ser	Phe	Leu	Pro	Ser	Gly	Val	Phe	Asp	Gly	Met	Pro	Pro	Asn	
45		635				640					645					650	
	Leu	Lys	Asn	Leu	Ser	Leu	Ala	Lys	Asn	Gly	Leu	Lys	Ser	Phe	Ser	Trp	
				655						660					665		
50	Lys	Lys	Leu	Gln	Cys	Leu	Lys	Asn	Leu	Glu	Thr	Leu	Asp	Leu	Ser	His	
				670					675					680			
	Asn	Gln	Leu	Thr	Thr	Val	Pro	Glu	Arg	Leu	Ser	Asn	Cys	Ser	Arg	Ser	
			685					690					695				
55	Leu	Lys	Asn	Leu	Ile	Leu	Lys	Asn	Asn	Gln	Ile	Arg	Ser	Leu	Thr	Lys	
		700					705					710					
	Tyr	Phe	Leu	Gln	Asp	Ala	Phe	Gln	Leu	Arg	Tyr	Leu	Asp	Leu	Ser	Ser	
60		715				720					725					730	

Asn Lys Ile Gln Met Ile Gln Lys Thr Ser Phe Pro Glu Asn Val Leu
 735 740 745
 5 Asn Asn Leu Lys Met Leu Leu Leu His His Asn Arg Phe Leu Cys Thr
 750 755 760
 Cys Asp Ala Val Trp Phe Val Trp Trp Val Asn His Thr Glu Val Thr
 765 770 775
 10 Ile Pro Tyr Leu Ala Thr Asp Val Thr Cys Val Gly Pro Gly Ala His
 780 785 790
 Lys Gly Gln Ser Val Ile Ser Leu Asp Leu Tyr Thr Cys Glu Leu Asp
 15 795 800 805 810
 Leu Thr Asn Leu Ile Leu Phe Ser Leu Ser Ile Ser Val Ser Leu Phe
 815 820 825
 20 Leu Met Val Met Met Thr Ala Ser His Leu Tyr Phe Trp Asp Val Trp
 830 835 840
 Tyr Ile Tyr His Phe Cys Lys Ala Lys Ile Lys Gly Tyr Gln Arg Leu
 845 850 855
 25 Ile Ser Pro Asp Cys Cys Tyr Asp Ala Phe Ile Val Tyr Asp Thr Lys
 860 865 870
 Asp Pro Ala Val Thr Glu Trp Val Leu Ala Glu Leu Val Ala Lys Leu
 30 875 880 885 890
 Glu Asp Pro Arg Glu Lys His Phe Asn Leu Cys Leu Glu Glu Arg Asp
 895 900 905
 35 Trp Leu Pro Gly Gln Pro Val Leu Glu Asn Leu Ser Gln Ser Ile Gln
 910 915 920
 Leu Ser Lys Lys Thr Val Phe Val Met Thr Asp Lys Tyr Ala Lys Thr
 925 930 935
 40 Glu Asn Phe Lys Ile Ala Phe Tyr Leu Ser His Gln Arg Leu Met Asp
 940 945 950
 Glu Lys Val Asp Val Ile Ile Leu Ile Phe Leu Glu Lys Pro Phe Gln
 45 955 960 965 970
 Lys Ser Lys Phe Leu Gln Leu Arg Lys Arg Leu Cys Gly Ser Ser Val
 975 980 985
 50 Leu Glu Trp Pro Thr Asn Pro Gln Ala His Pro Tyr Phe Trp Gln Cys
 990 995 1000
 Leu Lys Asn Ala Leu Ala Thr Asp Asn His Val Ala Tyr Ser Gln Val
 1005 1010 1015
 55 Phe Lys Glu Thr Val
 1020

(2) INFORMATION FOR SEQ ID NO:13:

60 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 180 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

10 (A) NAME/KEY: CDS
(B) LOCATION: 1..177

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

15 CTT GGA AAA CCT CTT CAG AAG TCT AAG TTT CTT CAG CTC AGG AAG AGA 48
Leu Gly Lys Pro Leu Gln Lys Ser Lys Phe Leu Gln Leu Arg Lys Arg
1 5 10 15
20 CTC TGC AGG AGC TCT GTC CTT GAG TGG CCT GCA AAT CCA CAG GCT CAC 96
Leu Cys Arg Ser Ser Val Leu Glu Trp Pro Ala Asn Pro Gln Ala His
20 25 30
25 CCA TAC TTC TGG CAG TGC CTG AAA AAT GCC CTG ACC ACA GAC AAT CAT 144
Pro Tyr Phe Trp Gln Cys Leu Lys Asn Ala Leu Thr Thr Asp Asn His
35 40 45
30 GTG GCT TAT AGT CAA ATG TTC AAG GAA ACA GTC TAG 180
Val Ala Tyr Ser Gln Met Phe Lys Glu Thr Val
50 55

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 59 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Leu Gly Lys Pro Leu Gln Lys Ser Lys Phe Leu Gln Leu Arg Lys Arg
1 5 10 15
45 Leu Cys Arg Ser Ser Val Leu Glu Trp Pro Ala Asn Pro Gln Ala His
20 25 30
50 Pro Tyr Phe Trp Gln Cys Leu Lys Asn Ala Leu Thr Thr Asp Asn His
35 40 45
Val Ala Tyr Ser Gln Met Phe Lys Glu Thr Val
50 55

55 (2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

60 (A) LENGTH: 990 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5 (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 2..988

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

	G AAT TCC AGA CTT ATA AAC TTG AAA AAT CTC TAT TTG GCC TGG AAC	46
	Asn Ser Arg Leu Ile Asn Leu Lys Asn Leu Tyr Leu Ala Trp Asn	
	1 5 10 15	
15	TGC TAT TTT AAC AAA GTT TGC GAG AAA ACT AAC ATA GAA GAT GGA GTA	94
	Cys Tyr Phe Asn Lys Val Cys Glu Lys Thr Asn Ile Glu Asp Gly Val	
	20 25 30	
20	TTT GAA ACG CTG ACA AAT TTG GAG TTG CTA TCA CTA TCT TTC AAT TCT	142
	Phe Glu Thr Leu Thr Asn Leu Glu Leu Leu Ser Leu Ser Phe Asn Ser	
	35 40 45	
25	CTT TCA CAT GTG CCA CCC AAA CTG CCA AGC TCC CTA CGC AAA CTT TTT	190
	Leu Ser His Val Pro Pro Lys Leu Pro Ser Ser Leu Arg Lys Leu Phe	
	50 55 60	
30	CTG AGC AAC ACC CAG ATC AAA TAC ATT AGT GAA GAA GAT TTC AAG GGA	238
	Leu Ser Asn Thr Gln Ile Lys Tyr Ile Ser Glu Glu Asp Phe Lys Gly	
	65 70 75	
35	TTG ATA AAT TTA ACA TTA CTA GAT TTA AGC GGG AAC TGT CCG AGG TGC	286
	Leu Ile Asn Leu Thr Leu Leu Asp Leu Ser Gly Asn Cys Pro Arg Cys	
	80 85 90 95	
40	TTC AAT GCC CCA TTT CCA TGC GTG CCT TGT GAT GGT GGT GCT TCA ATT	334
	Phe Asn Ala Pro Phe Pro Cys Val Pro Cys Asp Gly Gly Ala Ser Ile	
	100 105 110	
45	AAT ATA GAT CGT TTT GCT TTT CAA AAC TTG ACC CAA CTT CGA TAC CTA	382
	Asn Ile Asp Arg Phe Ala Phe Gln Asn Leu Thr Gln Leu Arg Tyr Leu	
	115 120 125	
50	AAC CTC TCT AGC ACT TCC CTC AGG AAG ATT AAT GCT GCC TGG TTT AAA	430
	Asn Leu Ser Ser Thr Ser Leu Arg Lys Ile Asn Ala Ala Trp Phe Lys	
	130 135 140	
55	AAT ATG CCT CAT CTG AAG GTG CTG GAT CTT GAA TTC AAC TAT TTA GTG	478
	Asn Met Pro His Leu Lys Val Leu Asp Leu Glu Phe Asn Tyr Leu Val	
	145 150 155	
60	GGA GAA ATA GCC TCT GGG GCA TTT TTA ACG ATG CTG CCC CGC TTA GAA	526
	Gly Glu Ile Ala Ser Gly Ala Phe Leu Thr Met Leu Pro Arg Leu Glu	
	160 165 170 175	
65	ATA CTT GAC TTG TCT TTT AAC TAT ATA AAG GGG AGT TAT CCA CAG CAT	574
	Ile Leu Asp Leu Ser Phe Asn Tyr Ile Lys Gly Ser Tyr Pro Gln His	
	180 185 190	
70	ATT AAT ATT TCC AGA AAC TTC TCT AAA CTT TTG TCT CTA CGG GCA TTG	622
	Ile Asn Ile Ser Arg Asn Phe Ser Lys Leu Leu Ser Leu Arg Ala Leu	

	195	200	205	
5	CAT TTA AGA GGT TAT GTG TTC CAG GAA CTC AGA GAA GAT GAT TTC CAG His Leu Arg Gly Tyr Val Phe Gln Glu Leu Arg Glu Asp Asp Phe Gln 210 215 220			670
10	CCC CTG ATG CAG CTT CCA AAC TTA TCG ACT ATC AAC TTG GGT ATT AAT Pro Leu Met Gln Leu Pro Asn Leu Ser Thr Ile Asn Leu Gly Ile Asn 225 230 235			718
15	TTT ATT AAG CAA ATC GAT TTC AAA CTT TTC CAA AAT TTC TCC AAT CTG Phe Ile Lys Gln Ile Asp Phe Lys Leu Phe Gln Asn Phe Ser Asn Leu 240 245 250 255			766
20	GAA ATT ATT TAC TTG TCA GAA AAC AGA ATA TCA CCG TTG GTA AAA GAT Glu Ile Ile Tyr Leu Ser Glu Asn Arg Ile Ser Pro Leu Val Lys Asp 260 265 270			814
25	ACC CGG CAG AGT TAT GCA AAT AGT TCC TCT TTT CAA CGT CAT ATC CGG Thr Arg Gln Ser Tyr Ala Asn Ser Ser Ser Phe Gln Arg His Ile Arg 275 280 285			862
30	AAA CGA CGC TCA ACA GAT TTT GAG TTT GAC CCA CAT TCG AAC TTT TAT Lys Arg Arg Ser Thr Asp Phe Glu Phe Asp Pro His Ser Asn Phe Tyr 290 295 300			910
35	CAT TTC ACC CGT CCT TTA ATA AAG CCA CAA TGT GCT GCT TAT GGA AAA His Phe Thr Arg Pro Leu Ile Lys Pro Gln Cys Ala Ala Tyr Gly Lys 305 310 315			958
40	GCC TTA GAT TTA AGC CTC AAC AGT ATT TTC TT Ala Leu Asp Leu Ser Leu Asn Ser Ile Phe 320 325			990

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 329 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Asn Ser Arg Leu Ile Asn Leu Lys Asn Leu Tyr Leu Ala Trp Asn Cys 1 5 10 15	
Tyr Phe Asn Lys Val Cys Glu Lys Thr Asn Ile Glu Asp Gly Val Phe 20 25 30	
Glu Thr Leu Thr Asn Leu Glu Leu Leu Ser Leu Ser Phe Asn Ser Leu 35 40 45	
Ser His Val Pro Pro Lys Leu Pro Ser Ser Leu Arg Lys Leu Phe Leu 50 55 60	
Ser Asn Thr Gln Ile Lys Tyr Ile Ser Glu Glu Asp Phe Lys Gly Leu 65 70 75 80	

```

      Ile Asn Leu Thr Leu Leu Asp Leu Ser Gly Asn Cys Pro Arg Cys Phe
                        85                      90                      95
5      Asn Ala Pro Phe Pro Cys Val Pro Cys Asp Gly Gly Ala Ser Ile Asn
      100                      105                      110
      Ile Asp Arg Phe Ala Phe Gln Asn Leu Thr Gln Leu Arg Tyr Leu Asn
      115                      120                      125
10     Leu Ser Ser Thr Ser Leu Arg Lys Ile Asn Ala Ala Trp Phe Lys Asn
      130                      135                      140
      Met Pro His Leu Lys Val Leu Asp Leu Glu Phe Asn Tyr Leu Val Gly
      145                      150                      155                      160
15     Glu Ile Ala Ser Gly Ala Phe Leu Thr Met Leu Pro Arg Leu Glu Ile
      165                      170                      175
      Leu Asp Leu Ser Phe Asn Tyr Ile Lys Gly Ser Tyr Pro Gln His Ile
20     180                      185                      190
      Asn Ile Ser Arg Asn Phe Ser Lys Leu Leu Ser Leu Arg Ala Leu His
      195                      200                      205
25     Leu Arg Gly Tyr Val Phe Gln Glu Leu Arg Glu Asp Asp Phe Gln Pro
      210                      215                      220
      Leu Met Gln Leu Pro Asn Leu Ser Thr Ile Asn Leu Gly Ile Asn Phe
30     225                      230                      235                      240
      Ile Lys Gln Ile Asp Phe Lys Leu Phe Gln Asn Phe Ser Asn Leu Glu
      245                      250                      255
      Ile Ile Tyr Leu Ser Glu Asn Arg Ile Ser Pro Leu Val Lys Asp Thr
35     260                      265                      270
      Arg Gln Ser Tyr Ala Asn Ser Ser Ser Phe Gln Arg His Ile Arg Lys
      275                      280                      285
40     Arg Arg Ser Thr Asp Phe Glu Phe Asp Pro His Ser Asn Phe Tyr His
      290                      295                      300
      Phe Thr Arg Pro Leu Ile Lys Pro Gln Cys Ala Ala Tyr Gly Lys Ala
45     305                      310                      315                      320
      Leu Asp Leu Ser Leu Asn Ser Ile Phe
      325

```

(2) INFORMATION FOR SEQ ID NO:17:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1557 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

55

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

60

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..513

(ix) FEATURE:

5 (A) NAME/KEY: misc_feature
(B) LOCATION: 278
(D) OTHER INFORMATION: /note= "nucleotide 278 designated
G, may be G or C"

(ix) FEATURE:

10 (A) NAME/KEY: misc_feature
(B) LOCATION: 445
(D) OTHER INFORMATION: /note= "nucleotide 445 designated
A, may be A or T"

(ix) FEATURE:

15 (A) NAME/KEY: misc_feature
(B) LOCATION: 572
(D) OTHER INFORMATION: /note= "nucleotides 572, 593, 600,
20 607, 617, 622, 625, 631, 640, 646, 653, 719, 775, and 861 are
designated C; each may be A, C, G, or T"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

25	CAG TCT CTT TCC ACA TCC CAA ACT TTC TAT GAT GCT TAC ATT TCT TAT	48
	Gln Ser Leu Ser Thr Ser Gln Thr Phe Tyr Asp Ala Tyr Ile Ser Tyr	
	1 5 10 15	
30	GAC ACC AAA GAT GCC TCT GTT ACT GAC TGG GTG ATA AAT GAG CTG CGC	96
	Asp Thr Lys Asp Ala Ser Val Thr Asp Trp Val Ile Asn Glu Leu Arg	
	20 25 30	
35	TAC CAC CTT GAA GAG AGC CGA GAC AAA AAC GTT CTC CTT TGT CTA GAG	144
	Tyr His Leu Glu Glu Ser Arg Asp Lys Asn Val Leu Leu Cys Leu Glu	
	35 40 45	
40	GAG AGG GAT TGG GAC CCG GGA TTG GCC ATC ATC GAC AAC CTC ATG CAG	192
	Glu Arg Asp Trp Asp Pro Gly Leu Ala Ile Ile Asp Asn Leu Met Gln	
	50 55 60	
45	AGC ATC AAC CAA AGC AAG AAA ACA GTA TTT GTT TTA ACC AAA AAA TAT	240
	Ser Ile Asn Gln Ser Lys Lys Thr Val Phe Val Leu Thr Lys Lys Tyr	
	65 70 75 80	
50	GCA AAA AGC TGG AAC TTT AAA ACA GCT TTT TAC TTG GGC TTG CAG AGG	288
	Ala Lys Ser Trp Asn Phe Lys Thr Ala Phe Tyr Leu Gly Leu Gln Arg	
	85 90 95	
55	CTA ATG GGT GAG AAC ATG GAT GTG ATT ATA TTT ATC CTG CTG GAG CCA	336
	Leu Met Gly Glu Asn Met Asp Val Ile Ile Phe Ile Leu Leu Glu Pro	
	100 105 110	
60	GTG TTA CAG CAT TCT CCG TAT TTG AGG CTA CGG CAG CGG ATC TGT AAG	384
	Val Leu Gln His Ser Pro Tyr Leu Arg Leu Arg Gln Arg Ile Cys Lys	
	115 120 125	
60	AGC TCC ATC CTC CAG TGG CCT GAC AAC CCG AAG GCA GAA AGG TTG TTT	432
	Ser Ser Ile Leu Gln Trp Pro Asp Asn Pro Lys Ala Glu Arg Leu Phe	
	130 135 140	
	TGG CAA ACT CTG AGA AAT GTG GTC TTG ACT GAA AAT GAT TCA CGG TAT	480

Trp Gln Thr Leu Arg Asn Val Val Leu Thr Glu Asn Asp Ser Arg Tyr
 145 150 155 160
 AAC AAT ATG TAT GTC GAT TCC ATT AAG CAA TAC TAACTGACGT TAAGTCATGA 533
 5 Asn Asn Met Tyr Val Asp Ser Ile Lys Gln Tyr
 165 170
 TTTCGCGCCA TAATAAAGAT GCAAAGGAAT GACATTTCCG TATTAGTTAT CTATTGCTAC 593
 10 GGTAACCAAA TTA CTCCAA AAACCTTACG TCGGTTTCAA AACAACCACA TTCTGCTGGC 653
 CCCACAGTTT TTGAGGGTCA GGAGTCCAGG CCCAGCATAA CTGGGTCTTC TGCTTCAGGG 713
 15 TGTCTCCAGA GGCTGCAATG TAGGTGTTCA CCAGAGACAT AGGCATCACT GGGGTCACAC 773
 TCCATGTGGT TGTTTTCTGG ATTCAATTCC TCCTGGGCTA TTGGCCAAAG GCTATACTCA 833
 TGTAAGCCAT GCGAGCCTAT CCCACAACGG CAGCTTGCTT CATCAGAGCT AGCAAAAAAG 893
 20 AGAGGTTGCT AGCAAGATGA AGTCACAATC TTTTGTAATC GAATCAAAAA AGTGATATCT 953
 CATCACTTTG GCCATATTCT ATTTGTTAGA AGTAAACCAC AGGTCCCACC AGCTCCATGG 1013
 25 GAGTGACCAC CTCAGTCCAG GGAAAACAGC TGAAGACCAA GATGGTGAGC TCTGATTGCT 1073
 TCAGTTGGTC ATCAACTATT TTCCCTTGAC TGCTGTCCCTG GGATGGCCGG CTATCTTGAT 1133
 GGATAGATTG TGAATATCAG GAGGCCAGGG ATCACTGTGG ACCATCTTAG CAGTTGACCT 1193
 30 AACACATCTT CTTTTCAATA TCTAAGAACT TTTGCCACTG TGAATAATGG TCCTAATATT 1253
 AAGCTGTTGT TTATATTTAT CATATATCTA TGGCTACATG GTTATATTAT GCTGTGGTTG 1313
 35 CGTTCGGTTT TATTTACAGT TGCTTTTACA AATATTTGCT GTAACATTTG ACTTCTAAGG 1373
 TTTAGATGCC ATTTAAGAAC TGAGATGGAT AGCTTTTAAA GCATCTTTTA CTTCTTACCA 1433
 TTTTTTAAAA GTATGCAGCT AAATTCGAAG CTTTGTGCT ATATTGTTAA TTGCCATTGC 1493
 40 TGTAATCTT AAAATGAATG AATAAAAATG TTTCAATTTA AAAAAAAAAA AAAAAAAAAA 1553
 AAAA 1557

45 (2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 171 amino acids

(B) TYPE: amino acid

50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

55 Gln Ser Leu Ser Thr Ser Gln Thr Phe Tyr Asp Ala Tyr Ile Ser Tyr
 1 5 10 15
 Asp Thr Lys Asp Ala Ser Val Thr Asp Trp Val Ile Asn Glu Leu Arg
 60 20 25 30

Tyr His Leu Glu Glu Ser Arg Asp Lys Asn Val Leu Leu Cys Leu Glu
 35 40 45
 5 Glu Arg Asp Trp Asp Pro Gly Leu Ala Ile Ile Asp Asn Leu Met Gln
 50 55 60
 Ser Ile Asn Gln Ser Lys Lys Thr Val Phe Val Leu Thr Lys Lys Tyr
 65 70 75 80
 10 Ala Lys Ser Trp Asn Phe Lys Thr Ala Phe Tyr Leu Gly Leu Gln Arg.
 85 90 95
 Leu Met Gly Glu Asn Met Asp Val Ile Ile Phe Ile Leu Leu Glu Pro
 100 105 110
 15 Val Leu Gln His Ser Pro Tyr Leu Arg Leu Arg Gln Arg Ile Cys Lys
 115 120 125
 Ser Ser Ile Leu Gln Trp Pro Asp Asn Pro Lys Ala Glu Arg Leu Phe
 20 130 135 140
 Trp Gln Thr Leu Arg Asn Val Val Leu Thr Glu Asn Asp Ser Arg Tyr
 145 150 155 160
 25 Asn Asn Met Tyr Val Asp Ser Ile Lys Gln Tyr
 165 170

(2) INFORMATION FOR SEQ ID NO:19:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 629 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 35 (ii) MOLECULE TYPE: cDNA
 (ix) FEATURE:
 40 (A) NAME/KEY: CDS
 (B) LOCATION: 1..486
 (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 45 (B) LOCATION: 144
 (D) OTHER INFORMATION: /note= "nucleotides 144 and 225
 designated C; may be C or T"

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

AAT GAA TTG ATC CCC AAT CTA GAG AAG GAA GAT GGT TCT ATC TTG ATT 48
 Asn Glu Leu Ile Pro Asn Leu Glu Lys Glu Asp Gly Ser Ile Leu Ile
 1 5 10 15
 55 TGC CTT TAT GAA AGC TAC TTT GAC CCT GGC AAA AGC ATT AGT GAA AAT 96
 Cys Leu Tyr Glu Ser Tyr Phe Asp Pro Gly Lys Ser Ile Ser Glu Asn
 20 25 30
 60 ATT GTA AGC TTC ATT GAG AAA AGC TAT AAG TCC ATC TTT GTT TTG TCC 144
 Ile Val Ser Phe Ile Glu Lys Ser Tyr Lys Ser Ile Phe Val Leu Ser

	35	40	45	
5	CCC AAC TTT GTC CAG AAT GAG TGG TGC CAT TAT GAA TTC TAC TTT GCC Pro Asn Phe Val Gln Asn Glu Trp Cys His Tyr Glu Phe Tyr Phe Ala	55	60	192
10	CAC CAC AAT CTC TTC CAT GAA AAT TCT GAT CAC ATA ATT CTT ATC TTA His His Asn Leu Phe His Glu Asn Ser Asp His Ile Ile Leu Ile Leu	65	70	240
	CTG GAA CCC ATT CCA TTC TAT TGC ATT CCC ACC AGG TAT CAT AAA CTG Leu Glu Pro Ile Pro Phe Tyr Cys Ile Pro Thr Arg Tyr His Lys Leu	75	80	288
15	GAA GCT CTC CTG GAA AAA AAA GCA TAC TTG GAA TGG CCC AAG GAT AGG Glu Ala Leu Leu Glu Lys Lys Ala Tyr Leu Glu Trp Pro Lys Asp Arg	85	90	336
20	CGT AAA TGT GGG CTT TTC TGG GCA AAC CTT CGA GCT GCT GTT AAT GTT Arg Lys Cys Gly Leu Phe Trp Ala Asn Leu Arg Ala Ala Val Asn Val	95	100	384
25	AAT GTA TTA GCC ACC AGA GAA ATG TAT GAA CTG CAG ACA TTC ACA GAG Asn Val Leu Ala Thr Arg Glu Met Tyr Glu Leu Gln Thr Phe Thr Glu	105	110	432
30	TTA AAT GAA GAG TCT CGA GGT TCT ACA ATC TCT CTG ATG AGA ACA GAC Leu Asn Glu Glu Ser Arg Gly Ser Thr Ile Ser Leu Met Arg Thr Asp	115	120	480
	TGT CTA TAAAATCCCA CAGTCCTTGG GAAGTTGGGG ACCACATACA CTGTTGGGAT Cys Leu	125	130	536
35	GTACATTGAT ACAACCTTTA TGATGGCAAT TTGACAATAT TTATTAAAAT AAAAAATGGT	135	140	596
	TATTCCTTC AAAAAAAAAA AAAAAAAAAA AAA	145	150	629
40	(2) INFORMATION FOR SEQ ID NO:20:			
	(i) SEQUENCE CHARACTERISTICS:			
	(A) LENGTH: 162 amino acids			
	(B) TYPE: amino acid			
45	(D) TOPOLOGY: linear			
	(ii) MOLECULE TYPE: protein			
50	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:			
	Asn Glu Leu Ile Pro Asn Leu Glu Lys Glu Asp Gly Ser Ile Leu Ile	1	5	10
55	Cys Leu Tyr Glu Ser Tyr Phe Asp Pro Gly Lys Ser Ile Ser Glu Asn	15	20	25
	Ile Val Ser Phe Ile Glu Lys Ser Tyr Lys Ser Ile Phe Val Leu Ser	25	30	35
60	Pro Asn Phe Val Gln Asn Glu Trp Cys His Tyr Glu Phe Tyr Phe Ala	35	40	45
		50	55	60

His His Asn Leu Phe His Glu Asn Ser Asp His Ile Ile Leu Ile Leu
 65 70 75 80
 5 Leu Glu Pro Ile Pro Phe Tyr Cys Ile Pro Thr Arg Tyr His Lys Leu
 85 90 95
 Glu Ala Leu Leu Glu Lys Lys Ala Tyr Leu Glu Trp Pro Lys Asp Arg
 100 105 110
 10 Arg Lys Cys Gly Leu Phe Trp Ala Asn Leu Arg Ala Ala Val Asn Val
 115 120 125
 Asn Val Leu Ala Thr Arg Glu Met Tyr Glu Leu Gln Thr Phe Thr Glu
 15 130 135 140
 Leu Asn Glu Glu Ser Arg Gly Ser Thr Ile Ser Leu Met Arg Thr Asp
 145 150 155 160
 20 Cys Leu

(2) INFORMATION FOR SEQ ID NO:21:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 427 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 35 (A) NAME/KEY: CDS
 (B) LOCATION: 1..426

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

40 AAG AAC TCC AAA GAA AAC CTC CAG TTT CAT GCT TTT ATT TCA TAT AGT 48
 Lys Asn Ser Lys Glu Asn Leu Gln Phe His Ala Phe Ile Ser Tyr Ser
 1 5 10 15
 45 GAA CAT GAT TCT GCC TGG GTG AAA AGT GAA TTG GTA CCT TAC CTA GAA 96
 Glu His Asp Ser Ala Trp Val Lys Ser Glu Leu Val Pro Tyr Leu Glu
 20 25 30
 AAA GAA GAT ATA CAG ATT TGT CTT CAT GAG AGA AAC TTT GTC CCT GGC 144
 50 Lys Glu Asp Ile Gln Ile Cys Leu His Glu Arg Asn Phe Val Pro Gly
 35 40 45
 AAG AGC ATT GTG GAA AAT ATC ATC AAC TGC ATT GAG AAG AGT TAC AAG 192
 55 Lys Ser Ile Val Glu Asn Ile Ile Asn Cys Ile Glu Lys Ser Tyr Lys
 50 55 60
 TCC ATC TTT GTT TTG TCT CCC AAC TTT GTC CAG AGT GAG TGG TGC CAT 240
 Ser Ile Phe Val Leu Ser Pro Asn Phe Val Glu Ser Glu Trp Cys His
 65 70 75 80
 60 TAC GAA CTC TAT TTT GCC CAT CAC AAT CTC TTT CAT GAA GGA TCT AAT 288

Tyr Glu Leu Tyr Phe Ala His His Asn Leu Phe His Glu Gly Ser Asn
 85 90 95
 5 AAC TTA ATC CTC ATC TTA CTG GAA CCC ATT CCA CAG AAC AGC ATT CCC 336
 Asn Leu Ile Leu Ile Leu Leu Glu Pro Ile Pro Gln Asn Ser Ile Pro
 100 105 110
 10 AAC AAG TAC CAC AAG CTG AAG GCT CTC ATG ACG CAG CGG ACT TAT TTG 384
 Asn Lys Tyr His Lys Leu Lys Ala Leu Met Thr Gln Arg Thr Tyr Leu
 115 120 125
 15 CAG TGG CCC AAG GAG AAA AGC AAA CGT GGG CTC TTT TGG GCT 426
 Gln Trp Pro Lys Glu Lys Ser Lys Arg Gly Leu Phe Trp Ala
 130 135 140
 A 427

(2) INFORMATION FOR SEQ ID NO:22:
 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 142 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 25 (ii) MOLECULE TYPE: protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

30 Lys Asn Ser Lys Glu Asn Leu Gln Phe His Ala Phe Ile Ser Tyr Ser
 1 5 10 15
 Glu His Asp Ser Ala Trp Val Lys Ser Glu Leu Val Pro Tyr Leu Glu
 20 25 30
 35 Lys Glu Asp Ile Gln Ile Cys Leu His Glu Arg Asn Phe Val Pro Gly
 35 40 45
 40 Lys Ser Ile Val Glu Asn Ile Ile Asn Cys Ile Glu Lys Ser Tyr Lys
 50 55 60
 Ser Ile Phe Val Leu Ser Pro Asn Phe Val Gln Ser Glu Trp Cys His
 65 70 75 80
 45 Tyr Glu Leu Tyr Phe Ala His His Asn Leu Phe His Glu Gly Ser Asn
 85 90 95
 Asn Leu Ile Leu Ile Leu Leu Glu Pro Ile Pro Gln Asn Ser Ile Pro
 100 105 110
 50 Asn Lys Tyr His Lys Leu Lys Ala Leu Met Thr Gln Arg Thr Tyr Leu
 115 120 125
 55 Gln Trp Pro Lys Glu Lys Ser Lys Arg Gly Leu Phe Trp Ala
 130 135 140

(2) INFORMATION FOR SEQ ID NO:23:
 60 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 662 base pairs
 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..627

10

(ix) FEATURE:

(A) NAME/KEY: misc_feature
(B) LOCATION: 54
(D) OTHER INFORMATION: /note= "nucleotides 54, 103, and
15 345 are designated A; each may be A or G"

(ix) FEATURE:

(A) NAME/KEY: misc_feature
(B) LOCATION: 313
(D) OTHER INFORMATION: /note= "nucleotide 313 designated
20 G, may be G or T"

(ix) FEATURE:

(A) NAME/KEY: misc_feature
(B) LOCATION: 316
(D) OTHER INFORMATION: /note= "nucleotides 316, 380, 407,
25 and 408 designated C; each may be A, C, G, or T"

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GCT TCC ACC TGT GCC TGG CCT GGC TTC CCT GGC GGG GGC GGC AAA GTG	48
Ala Ser Thr Cys Ala Trp Pro Gly Phe Pro Gly Gly Gly Gly Lys Val	
1 5 10 15	
35 GGC GAA ATG AGG ATG CCC TGC CCT ACG ATG CCT TCG TGG TCT TCG ACA	96
Gly Glu Met Arg Met Pro Cys Pro Thr Met Pro Ser Trp Ser Ser Thr	
20 25 30	
40 AAA CGC AGA GCG CAG TGG CAG ACT GGG TGT ACA ACG AGC TTC GGG GGC	144
Lys Arg Arg Ala Gln Trp Gln Thr Gly Cys Thr Thr Ser Phe Gly Gly	
35 40 45	
45 AGC TGG AGG AGT GCC GTG GGC GCT GGG CAC TCC GCC TGT GCC TGG AGG	192
Ser Trp Arg Ser Ala Val Gly Ala Gly His Ser Ala Cys Ala Trp Arg	
50 55 60	
50 AAC GCG ACT GGC TGC CTG GCA AAA CCC TCT TTG AGA ACC TGT GGG CCT	240
Asn Ala Thr Gly Cys Leu Ala Lys Pro Ser Leu Arg Thr Cys Gly Pro	
65 70 75 80	
55 CGG TCT ATG GCA GCC GCA AGA CGC TGT TTG TGC TGG CCC ACA CGG ACC	288
Arg Ser Met Ala Ala Ala Arg Arg Cys Leu Cys Trp Pro Thr Arg Thr	
85 90 95	
60 GGG TCA GTG GTC TCT TGC GCG CCA GTT CTC CTG CTG GCC CAG CAG CGC	336
Gly Ser Val Val Ser Cys Ala Pro Val Leu Leu Leu Ala Gln Gln Arg	
100 105 110	
60 CTG CTG GAA GAC CGC AAG GAC GTC GTG GTG CTG GTG ATC CTA ACG CCT	384
Leu Leu Glu Asp Arg Lys Asp Val Val Val Leu Val Ile Leu Thr Pro	

	115	120	125	
	GAC GGC CAA GCC TCC CGA CTA CCC GAT GCG CTG ACC AGC GCC TCT GCC			432
	Asp Gly Gln Ala Ser Arg Leu Pro Asp Ala Leu Thr Ser Ala Ser Ala			
5	130	135	140	
	GCC AGA GTG TCC TCC TCT GGC CCC ACC AGC CCA GTG GTC GCG CAG CTT			480
	Ala Arg Val Ser Ser Ser Gly Pro Thr Ser Pro Val Val Ala Gln Leu			
10	145	150	155	160
	CTG AGG CCA GCA TGC ATG GCC CTG ACC AGG GAC AAC CAC CAC TTC TAT			528
	Leu Arg Pro Ala Cys Met Ala Leu Thr Arg Asp Asn His His Phe Tyr			
	165	170	175	
15	AAC CGG AAC TTC TGC CAG GGA ACC CAC GGC CGA ATA GCC GTG AGC CGG			576
	Asn Arg Asn Phe Cys Gln Gly Thr His Gly Arg Ile Ala Val Ser Arg			
	180	185	190	
	AAT CCT GCA CGG TGC CAC CTC CAC ACA CAC CTA ACA TAT GCC TGC CTG			624
20	Asn Pro Ala Arg Cys His Leu His Thr His Leu Thr Tyr Ala Cys Leu			
	195	200	205	
	ATC TGACCAACAC ATGCTCGCCA CCCTCACCAC ACACC			662
25	Ile			

(2) INFORMATION FOR SEQ ID NO:24:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 209 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

40	Ala Ser Thr Cys Ala Trp Pro Gly Phe Pro Gly Gly Gly Gly Lys Val	
	1 5 10 15	
	Gly Glu Met Arg Met Pro Cys Pro Thr Met Pro Ser Trp Ser Ser Thr	
	20 25 30	
45	Lys Arg Arg Ala Gln Trp Gln Thr Gly Cys Thr Thr Ser Phe Gly Gly	
	35 40 45	
	Ser Trp Arg Ser Ala Val Gly Ala Gly His Ser Ala Cys Ala Trp Arg	
	50 55 60	
50	Asn Ala Thr Gly Cys Leu Ala Lys Pro Ser Leu Arg Thr Cys Gly Pro	
	65 70 75 80	
	Arg Ser Met Ala Ala Ala Arg Arg Cys Leu Cys Trp Pro Thr Arg Thr	
55	85 90 95	
	Gly Ser Val Val Ser Cys Ala Pro Val Leu Leu Leu Ala Gln Gln Arg	
	100 105 110	
60	Leu Leu Glu Asp Arg Lys Asp Val Val Val Leu Val Ile Leu Thr Pro	
	115 120 125	

Asp Gly Gln Ala Ser Arg Leu Pro Asp Ala Leu Thr Ser Ala Ser Ala
130 135 140

5 Ala Arg Val Ser Ser Ser Gly Pro Thr Ser Pro Val Val Ala Gln Leu
145 150 155 160

Leu Arg Pro Ala Cys Met Ala Leu Thr Arg Asp Asn His His Phe Tyr
165 170 175

10 Asn Arg Asn Phe Cys Gln Gly Thr His Gly Arg Ile Ala Val Ser Arg
180 185 190

Asn Pro Ala Arg Cys His Leu His Thr His Leu Thr Tyr Ala Cys Leu
15 195 200 205

Ile

20 (2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4865 base pairs
(B) TYPE: nucleic acid
25 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 107..2617

35 (ix) FEATURE:
(A) NAME/KEY: mat_peptide
(B) LOCATION: 173..2617

40 (ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 81
(D) OTHER INFORMATION: /note= "nucleotides 81, 3144, 3205,
and 3563 designated A, each may be A, C, G, or T"

45 (ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 84
(D) OTHER INFORMATION: /note= "nucleotide 84 designated C,
may be C or G"

50 (ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 739
(D) OTHER INFORMATION: /note= "nucleotide 739 designated
55 C, may be C or T"

(ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 3132
60 (D) OTHER INFORMATION: /note= "nucleotides 3132, 3532,
3538, and 3553 designated G, each may be G or T"

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 3638

5 (D) OTHER INFORMATION: /note= "nucleotide 3638 designated
A, may be A or T"

(ix) FEATURE:

(A) NAME/KEY: misc_feature

10 (B) LOCATION: 3677

(D) OTHER INFORMATION: /note= "nucleotides 3677, 3685, and
3736 designated C, each may be A or C"

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

	AAAATACTCC CTTGCCTCAA AAAGTCTCG GTCAAACGGT GATAGCAAAC CACGCATTCA	60
20	CAGGGCCACT GCTGCTCACA AAACCAGTGA GGATGATGCC AGGATG ATG TCT GCC Met Ser Ala -22 -20	115
25	TCG CGC CTG GCT GGG ACT CTG ATC CCA GCC ATG GCC TTC CTC TCC TGC Ser Arg Leu Ala Gly Thr Leu Ile Pro Ala Met Ala Phe Leu Ser Cys -15 -10 -5	163
	GTG AGA CCA GAA AGC TGG GAG CCC TGC GTG GAG GTT CCT AAT ATT ACT Val Arg Pro Glu Ser Trp Glu Pro Cys Val Glu Val Pro Asn Ile Thr 1 5 10	211
30	TAT CAA TGC ATG GAG CTG AAT TTC TAC AAA ATC CCC GAC AAC CTC CCC Tyr Gln Cys Met Glu Leu Asn Phe Tyr Lys Ile Pro Asp Asn Leu Pro 15 20 25	259
35	TTC TCA ACC AAG AAC CTG GAC CTG AGC TTT AAT CCC CTG AGG CAT TTA Phe Ser Thr Lys Asn Leu Asp Leu Ser Phe Asn Pro Leu Arg His Leu 30 35 40 45	307
40	GGC AGC TAT AGC TTC TTC AGT TTC CCA GAA CTG CAG GTG CTG GAT TTA Gly Ser Tyr Ser Phe Phe Ser Phe Pro Glu Leu Gln Val Leu Asp Leu 50 55 60	355
45	TCC AGG TGT GAA ATC CAG ACA ATT GAA GAT GGG GCA TAT CAG AGC CTA Ser Arg Cys Glu Ile Gln Thr Ile Glu Asp Gly Ala Tyr Gln Ser Leu 65 70 75	403
	AGC CAC CTC TCT ACC TTA ATA TTG ACA GGA AAC CCC ATC CAG AGT TTA Ser His Leu Ser Thr Leu Ile Leu Thr Gly Asn Pro Ile Gln Ser Leu 80 85 90	451
50	GCC CTG GGA GCC TTT TCT GGA CTA TCA AGT TTA CAG AAG CTG GTG GCT Ala Leu Gly Ala Phe Ser Gly Leu Ser Ser Leu Gln Lys Leu Val Ala 95 100 105	499
55	GTG GAG ACA AAT CTA GCA TCT CTA GAG AAC TTC CCC ATT GGA CAT CTC Val Glu Thr Asn Leu Ala Ser Leu Glu Asn Phe Pro Ile Gly His Leu 110 115 120 125	547
60	AAA ACT TTG AAA GAA CTT AAT GTG GCT CAC AAT CTT ATC CAA TCT TTC Lys Thr Leu Lys Glu Leu Asn Val Ala His Asn Leu Ile Gln Ser Phe 130 135 140	595

5	AAA TTA CCT GAG TAT TTT TCT AAT CTG ACC AAT CTA GAG CAC TTG GAC	643
	Lys Leu Pro Glu Tyr Phe Ser Asn Leu Thr Asn Leu Glu His Leu Asp 145 150 155	
10	CTT TCC AGC AAC AAG ATT CAA AGT ATT TAT TGC ACA GAC TTG CGG GTT	691
	Leu Ser Ser Asn Lys Ile Gln Ser Ile Tyr Cys Thr Asp Leu Arg Val 160 165 170	
15	CTA CAT CAA ATG CCC CTA CTC AAT CTC TCT TTA GAC CTG TCC CTG AAC	739
	Leu His Gln Met Pro Leu Leu Asn Leu Ser Leu Asp Leu Ser Leu Asn 175 180 185	
20	CCT ATG AAC TTT ATC CAA CCA GGT GCA TTT AAA GAA ATT AGG CTT CAT	787
	Pro Met Asn Phe Ile Gln Pro Gly Ala Phe Lys Glu Ile Arg Leu His 190 195 200 205	
25	AAG CTG ACT TTA AGA AAT AAT TTT GAT AGT TTA AAT GTA ATG AAA ACT	835
	Lys Leu Thr Leu Arg Asn Asn Phe Asp Ser Leu Asn Val Met Lys Thr 210 215 220	
30	TGT ATT CAA GGT CTG GCT GGT TTA GAA GTC CAT CGT TTG GTT CTG GGA	883
	Cys Ile Gln Gly Leu Ala Gly Leu Glu Val His Arg Leu Val Leu Gly 225 230 235	
35	GAA TTT AGA AAT GAA GGA AAC TTG GAA AAG TTT GAC AAA TCT GCT CTA	931
	Glu Phe Arg Asn Glu Gly Asn Leu Glu Lys Phe Asp Lys Ser Ala Leu 240 245 250	
40	GAG GGC CTG TGC AAT TTG ACC ATT GAA GAA TTC CGA TTA GCA TAC TTA	979
	Glu Gly Leu Cys Asn Leu Thr Ile Glu Glu Phe Arg Leu Ala Tyr Leu 255 260 265	
45	GAC TAC TAC CTC GAT GAT ATT ATT GAC TTA TTT AAT TGT TTG ACA AAT	1027
	Asp Tyr Tyr Leu Asp Asp Ile Ile Asp Leu Phe Asn Cys Leu Thr Asn 270 275 280 285	
50	GTT TCT TCA TTT TCC CTG GTG AGT GTG ACT ATT GAA AGG GTA AAA GAC	1075
	Val Ser Ser Phe Ser Leu Val Ser Val Thr Ile Glu Arg Val Lys Asp 290 295 300	
55	TTT TCT TAT AAT TTC GGA TGG CAA CAT TTA GAA TTA GTT AAC TGT AAA	1123
	Phe Ser Tyr Asn Phe Gly Trp Gln His Leu Glu Leu Val Asn Cys Lys 305 310 315	
60	TTT GGA CAG TTT CCC ACA TTG AAA CTC AAA TCT CTC AAA AGG CTT ACT	1171
	Phe Gly Gln Phe Pro Thr Leu Lys Leu Lys Ser Leu Lys Arg Leu Thr 320 325 330	
65	TTC ACT TCC AAC AAA GGT GGG AAT GCT TTT TCA GAA GTT GAT CTA CCA	1219
	Phe Thr Ser Asn Lys Gly Gly Asn Ala Phe Ser Glu Val Asp Leu Pro 335 340 345	
70	AGC CTT GAG TTT CTA GAT CTC AGT AGA AAT GGC TTG AGT TTC AAA GGT	1267
	Ser Leu Glu Phe Leu Asp Leu Ser Arg Asn Gly Leu Ser Phe Lys Gly 350 355 360 365	
75	TGC TGT TCT CAA AGT GAT TTT GGG ACA ACC AGC CTA AAG TAT TTA GAT	1315
	Cys Cys Ser Gln Ser Asp Phe Gly Thr Thr Ser Leu Lys Tyr Leu Asp 370 375 380	

	CTG AGC TTC AAT GGT GTT ATT ACC ATG AGT TCA AAC TTC TTG GGC TTA	1363
	Leu Ser Phe Asn Gly Val Ile Thr Met Ser Ser Asn Phe Leu Gly Leu	
	385 390 395	
5	GAA CAA CTA GAA CAT CTG GAT TTC CAG CAT TCC AAT TTG AAA CAA ATG	1411
	Glu Gln Leu Glu His Leu Asp Phe Gln His Ser Asn Leu Lys Gln Met	
	400 405 410	
10	AGT GAG TTT TCA GTA TTC CTA TCA CTC AGA AAC CTC ATT TAC CTT GAC	1459
	Ser Glu Phe Ser Val Phe Leu Ser Leu Arg Asn Leu Ile Tyr Leu Asp	
	415 420 425	
15	ATT TCT CAT ACT CAC ACC AGA GTT GCT TTC AAT GGC ATC TTC AAT GGC	1507
	Ile Ser His Thr His Thr Arg Val Ala Phe Asn Gly Ile Phe Asn Gly	
	430 435 440 445	
20	TTG TCC AGT CTC GAA GTC TTG AAA ATG GCT GGC AAT TCT TTC CAG GAA	1555
	Leu Ser Ser Leu Glu Val Leu Lys Met Ala Gly Asn Ser Phe Gln Glu	
	450 455 460	
	AAC TTC CTT CCA GAT ATC TTC ACA GAG CTG AGA AAC TTG ACC TTC CTG	1603
	Asn Phe Leu Pro Asp Ile Phe Thr Glu Leu Arg Asn Leu Thr Phe Leu	
	465 470 475	
25	GAC CTC TCT CAG TGT CAA CTG GAG CAG TTG TCT CCA ACA GCA TTT AAC	1651
	Asp Leu Ser Gln Cys Gln Leu Glu Gln Leu Ser Pro Thr Ala Phe Asn	
	480 485 490	
30	TCA CTC TCC AGT CTT CAG GTA CTA AAT ATG AGC CAC AAC AAC TTC TTT	1699
	Ser Leu Ser Ser Leu Gln Val Leu Asn Met Ser His Asn Asn Phe Phe	
	495 500 505	
35	TCA TTG GAT ACG TTT CCT TAT AAG TGT CTG AAC TCC CTC CAG GTT CTT	1747
	Ser Leu Asp Thr Phe Pro Tyr Lys Cys Leu Asn Ser Leu Gln Val Leu	
	510 515 520 525	
40	GAT TAC AGT CTC AAT CAC ATA ATG ACT TCC AAA AAA CAG GAA CTA CAG	1795
	Asp Tyr Ser Leu Asn His Ile Met Thr Ser Lys Lys Gln Glu Leu Gln	
	530 535 540	
	CAT TTT CCA AGT AGT CTA GCT TTC TTA AAT CTT ACT CAG AAT GAC TTT	1843
	His Phe Pro Ser Ser Leu Ala Phe Leu Asn Leu Thr Gln Asn Asp Phe	
	545 550 555	
45	GCT TGT ACT TGT GAA CAC CAG AGT TTC CTG CAA TGG ATC AAG GAC CAG	1891
	Ala Cys Thr Cys Glu His Gln Ser Phe Leu Gln Trp Ile Lys Asp Gln	
	560 565 570	
50	AGG CAG CTC TTG GTG GAA GTT GAA CGA ATG GAA TGT GCA ACA CCT TCA	1939
	Arg Gln Leu Leu Val Glu Val Glu Arg Met Glu Cys Ala Thr Pro Ser	
	575 580 585	
55	GAT AAG CAG GGC ATG CCT GTG CTG AGT TTG AAT ATC ACC TGT CAG ATG	1987
	Asp Lys Gln Gly Met Pro Val Leu Ser Leu Asn Ile Thr Cys Gln Met	
	590 595 600 605	
60	AAT AAG ACC ATC ATT GGT GTG TCG GTC CTC AGT GTG CTT GTA GTA TCT	2035
	Asn Lys Thr Ile Ile Gly Val Ser Val Leu Ser Val Leu Val Val Ser	
	610 615 620	
	GTT GTA GCA GTT CTG GTC TAT AAG TTC TAT TTT CAC CTG ATG CTT CTT	2083

	Val Val Ala Val Leu Val Tyr Lys Phe Tyr Phe His Leu Met Leu Leu	
	625 630 635	
5	GCT GGC TGC ATA AAG TAT GGT AGA GGT GAA AAC ATC TAT GAT GCC TTT Ala Gly Cys Ile Lys Tyr Gly Arg Gly Glu Asn Ile Tyr Asp Ala Phe	2131
	640 645 650	
10	GTT ATC TAC TCA AGC CAG GAT GAG GAC TGG GTA AGG AAT GAG CTA GTA Val Ile Tyr Ser Ser Gln Asp Glu Asp Trp Val Arg Asn Glu Leu Val	2179
	655 660 665	
15	AAG AAT TTA GAA GAA GGG GTG CCT CCA TTT CAG CTC TGC CTT CAC TAC Lys Asn Leu Glu Glu Gly Val Pro Pro Phe Gln Leu Cys Leu His Tyr	2227
	670 675 680 685	
	AGA GAC TTT ATT CCC GGT GTG GCC ATT GCT GCC AAC ATC ATC CAT GAA Arg Asp Phe Ile Pro Gly Val Ala Ile Ala Ala Asn Ile Ile His Glu	2275
	690 695 700	
20	GGT TTC CAT AAA AGC CGA AAG GTG ATT GTT GTG GTG TCC CAG CAC TTC Gly Phe His Lys Ser Arg Lys Val Ile Val Val Val Ser Gln His Phe	2323
	705 710 715	
25	ATC CAG AGC CGC TGG TGT ATC TTT GAA TAT GAG ATT GCT CAG ACC TGG Ile Gln Ser Arg Trp Cys Ile Phe Glu Tyr Glu Ile Ala Gln Thr Trp	2371
	720 725 730	
30	CAG TTT CTG AGC AGT CGT GCT GGT ATC ATC TTC ATT GTC CTG CAG AAG Gln Phe Leu Ser Ser Arg Ala Gly Ile Ile Phe Ile Val Leu Gln Lys	2419
	735 740 745	
35	GTG GAG AAG ACC CTG CTC AGG CAG CAG GTG GAG CTG TAC CGC CTT CTC Val Glu Lys Thr Leu Leu Arg Gln Gln Val Glu Leu Tyr Arg Leu Leu	2467
	750 755 760 765	
	AGC AGG AAC ACT TAC CTG GAG TGG GAG GAC AGT GTC CTG GGG CGG CAC Ser Arg Asn Thr Tyr Leu Glu Trp Glu Asp Ser Val Leu Gly Arg His	2515
	770 775 780	
40	ATC TTC TGG AGA CGA CTC AGA AAA GCC CTG CTG GAT GGT AAA TCA TGG Ile Phe Trp Arg Arg Leu Arg Lys Ala Leu Leu Asp Gly Lys Ser Trp	2563
	785 790 795	
45	AAT CCA GAA GGA ACA GTG GGT ACA GGA TGC AAT TGG CAG GAA GCA ACA Asn Pro Glu Gly Thr Val Gly Thr Gly Cys Asn Trp Gln Glu Ala Thr	2611
	800 805 810	
50	TCT ATC TGAAGAGGAA AAATAAAAAC CTCCTGAGGC ATTTCTTGCC CAGCTGGGTC Ser Ile	2667
	815	
	CAACACTTGT TCAGTTAATA AGTATTAAAT GCTGCCACAT GTCAGGCCTT ATGCTAAGGG	2727
55	TGAGTAATTC CATGGTGCAC TAGATATGCA GGGCTGCTAA TCTCAAGGAG CTTCCAGTGC	2787
	AGAGGGAATA AATGCTAGAC TAAAATACAG AGTCTTCCAG GTGGGCATTT CAACCAACTC	2847
	AGTCAAGGAA CCCATGACAA AGAAAGTCAT TTCAACTCTT ACCTCATCAA GTTGAATAAA	2907
60	GACAGAGAAA ACAGAAAGAG ACATTGTTCT TTTCTGAGT CTTTGAATG GAAATTGTAT	2967

	TATGTTATAG	CCATCATAAA	ACCATTTTGG	TAGTTTGTGAC	TGAACTGGGT	GTTCACTTTT	3027
	TCCTTTTTGA	TTGAATACAA	TTTAAATTCT	ACTTGATGAC	TGCAGTCGTC	AAGGGGCTCC	3087
5	TGATGCAAGA	TGCCCTTCC	ATTTTAAGTC	TGTCCTCTTA	CAGAGGTAA	AGTCTAATGG	3147
	CTAATTCCTA	AGGAAACCTG	ATTAACACAT	GCTCACAACC	ATCCTGGTCA	TTCTCGAACA	3207
10	TGTTCTATTT	TTTAACTAAT	CACCCCTGAT	ATATTTTAT	TTTTATATAT	CCAGTTTTC	3267
	TTTTTTTACG	TCTTGCCTAT	AAGCTAATAT	CATAAATAAG	GTGTGTTAAG	ACGTGCTTCA	3327
	AATATCCATA	TTAACCACTA	TTTTTCAAGG	AAGTATGGAA	AAGTACACTC	TGTCACCTTG	3387
15	TCACTCGATG	TCATTCCAAA	GTTATTGCCT	ACTAAGTAAT	GA CTGTCATG	AAAGCAGCAT	3447
	TGAAATAATT	TGTTTAAAGG	GGGCACTCTT	TTAAACGGGA	AGAAAATTC	CGCTTCCTGG	3507
20	TCTTATCATG	GACAATTTGG	GCTAGAGGCA	GGAAGGAAGT	GGGATGACCT	CAGGAAGTCA	3567
	CCTTTTCTTG	ATTCCAGAAA	CATATGGGCT	GATAAACCCG	GGGTGACCTC	ATGAAATGAG	3627
	TTGCAGCAGA	AGTTTATTTT	TTTCAGAACA	AGTGATGTTT	GATGGACCTC	TGAATCTCTT	3687
25	TAGGGAGACA	CAGATGGCTG	GGATCCCTCC	CCTGTACCCT	TCTCACTGCC	AGGAGAACTA	3747
	CGTGTGAAGG	TATTC AAGGC	AGGGAGTATA	CATTGCTGTT	TCCTGTTGGG	CAATGCTCCT	3807
30	TGACCACATT	TTGGGAAGAG	TGGATGTTAT	CATTGAGAAA	ACAATGTGTC	TGGAATTAAT	3867
	GGGGTTCTTA	TAAAGAAGGT	TCCCAGAAAA	GAATGTTTCA	TCCAGCTTCT	TCAGGAAACA	3927
	GGAACATTCA	AGGAAAAGGA	CAATCAGGAT	GTCATCAGGG	AAATGAAAAT	AAAAACCACA	3987
35	ATGAGATATC	ACCTTATACC	AGGTAGATGG	CTACTATAAA	AAAATGAAGT	GTCATCAAGG	4047
	ATATAGAGAA	ATTGGAACCC	TTCTTCACTG	CTGGAGGGAA	TGGAAAATGG	TGTAGCCGTT	4107
40	ATGAAAAACA	GTACGGAGGT	TTCTCAAAAA	TTAAAAATAG	AACTGCTATA	TGATCCAGCA	4167
	ATCTCACTTC	TGTATATATA	CCCCAAATAA	TTGAAATCAG	AATTTCAAGA	AAATATTTAC	4227
	ACTCCCATGT	TCATTGTGGC	ACTCTTCACA	ATCACTGTTT	CCAAAGTTAT	GGAAACAACC	4287
45	CAAATTTCCA	TTGGAAAATA	AATGGACAAA	GGAAATGTGC	ATATAACGTA	CAATGGGGAT	4347
	ATTATTCAGC	CTAAAAAAG	GGGGGATCCT	GTTATTTATG	ACAACATGAA	TAAACCCGGA	4407
50	GGCCATTATG	CTATGTAAAA	TGAGCAAGTA	ACAGAAAGAC	AAATACTGCC	TGATTTTCATT	4467
	TATATGAGGT	TCTAAAATAG	TCAAAC TCAT	AGAAGCAGAG	AATAGAACAG	TGGTTCCTAG	4527
	GGAAAAGGAG	GAAGGGAGAA	ATGAGGAAAT	AGGGAGTTGT	CTAATTGGTA	TAAAATTATA	4587
55	GTATGCAAGA	TGAATTAGCT	CTAAAGATCA	GCTGTATAGC	AGAGTTCGTA	TAATGAACAA	4647
	TACTGTATTA	TGCACTTAAC	ATTTTGTTAA	GAGGGTACCT	CTCATGTTAA	GTGTTCTTAC	4707
60	CATATACATA	TACACAAGGA	AGCTTTTGGA	GGTGATGGAT	ATATTTATTA	CCTTGATTGT	4767
	GGTGATGGTT	TGACAGGTAT	GTGACTATGT	CTAAACTCAT	CAAATTGTAT	ACATTAAATA	4827

TATGCAGTTT TATAATATCA AAAAAAAAAA AAAAAAAAAA

4865

5 (2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 837 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

15 Met Ser Ala Ser Arg Leu Ala Gly Thr Leu Ile Pro Ala Met Ala Phe
 -22 -20 -15 -10

20 Leu Ser Cys Val Arg Pro Glu Ser Trp Glu Pro Cys Val Glu Val Pro
 -5 1 5 10

Asn Ile Thr Tyr Gln Cys Met Glu Leu Asn Phe Tyr Lys Ile Pro Asp
 15 20 25

25 Asn Leu Pro Phe Ser Thr Lys Asn Leu Asp Leu Ser Phe Asn Pro Leu
 30 35 40

Arg His Leu Gly Ser Tyr Ser Phe Phe Ser Phe Pro Glu Leu Gln Val
 45 50 55

30 Leu Asp Leu Ser Arg Cys Glu Ile Gln Thr Ile Glu Asp Gly Ala Tyr
 60 65 70

Gln Ser Leu Ser His Leu Ser Thr Leu Ile Leu Thr Gly Asn Pro Ile
 35 75 80 85 90

Gln Ser Leu Ala Leu Gly Ala Phe Ser Gly Leu Ser Ser Leu Gln Lys
 95 100 105

40 Leu Val Ala Val Glu Thr Asn Leu Ala Ser Leu Glu Asn Phe Pro Ile
 110 115 120

Gly His Leu Lys Thr Leu Lys Glu Leu Asn Val Ala His Asn Leu Ile
 125 130 135

45 Gln Ser Phe Lys Leu Pro Glu Tyr Phe Ser Asn Leu Thr Asn Leu Glu
 140 145 150

His Leu Asp Leu Ser Ser Asn Lys Ile Gln Ser Ile Tyr Cys Thr Asp
 50 155 160 165 170

Leu Arg Val Leu His Gln Met Pro Leu Leu Asn Leu Ser Leu Asp Leu
 175 180 185

55 Ser Leu Asn Pro Met Asn Phe Ile Gln Pro Gly Ala Phe Lys Glu Ile
 190 195 200

Arg Leu His Lys Leu Thr Leu Arg Asn Asn Phe Asp Ser Leu Asn Val
 205 210 215

60 Met Lys Thr Cys Ile Gln Gly Leu Ala Gly Leu Glu Val His Arg Leu

	220		225		230
	Val Leu Gly Glu Phe Arg Asn Glu Gly Asn Leu Glu Lys Phe Asp Lys				
	235		240		245
5	Ser Ala Leu Glu Gly Leu Cys Asn Leu Thr Ile Glu Glu Phe Arg Leu				
		255		260	265
10	Ala Tyr Leu Asp Tyr Tyr Leu Asp Asp Ile Ile Asp Leu Phe Asn Cys				
		270		275	280
	Leu Thr Asn Val Ser Ser Phe Ser Leu Val Ser Val Thr Ile Glu Arg				
		285		290	295
15	Val Lys Asp Phe Ser Tyr Asn Phe Gly Trp Gln His Leu Glu Leu Val				
		300		305	310
	Asn Cys Lys Phe Gly Gln Phe Pro Thr Leu Lys Leu Lys Ser Leu Lys				
		315		320	325
20	Arg Leu Thr Phe Thr Ser Asn Lys Gly Gly Asn Ala Phe Ser Glu Val				
		335		340	345
25	Asp Leu Pro Ser Leu Glu Phe Leu Asp Leu Ser Arg Asn Gly Leu Ser				
		350		355	360
	Phe Lys Gly Cys Cys Ser Gln Ser Asp Phe Gly Thr Thr Ser Leu Lys				
		365		370	375
30	Tyr Leu Asp Leu Ser Phe Asn Gly Val Ile Thr Met Ser Ser Asn Phe				
		380		385	390
	Leu Gly Leu Glu Gln Leu Glu His Leu Asp Phe Gln His Ser Asn Leu				
		395		400	405
35	Lys Gln Met Ser Glu Phe Ser Val Phe Leu Ser Leu Arg Asn Leu Ile				
		415		420	425
40	Tyr Leu Asp Ile Ser His Thr His Thr Arg Val Ala Phe Asn Gly Ile				
		430		435	440
	Phe Asn Gly Leu Ser Ser Leu Glu Val Leu Lys Met Ala Gly Asn Ser				
		445		450	455
45	Phe Gln Glu Asn Phe Leu Pro Asp Ile Phe Thr Glu Leu Arg Asn Leu				
		460		465	470
	Thr Phe Leu Asp Leu Ser Gln Cys Gln Leu Glu Gln Leu Ser Pro Thr				
		475		480	485
50	Ala Phe Asn Ser Leu Ser Ser Leu Gln Val Leu Asn Met Ser His Asn				
		495		500	505
55	Asn Phe Phe Ser Leu Asp Thr Phe Pro Tyr Lys Cys Leu Asn Ser Leu				
		510		515	520
	Gln Val Leu Asp Tyr Ser Leu Asn His Ile Met Thr Ser Lys Lys Gln				
		525		530	535
60	Glu Leu Gln His Phe Pro Ser Ser Leu Ala Phe Leu Asn Leu Thr Gln				
		540		545	550

Asn Asp Phe Ala Cys Thr Cys Glu His Gln Ser Phe Leu Gln Trp Ile
 555 560 565 570
 5 Lys Asp Gln Arg Gln Leu Leu Val Glu Val Glu Arg Met Glu Cys Ala
 575 580 585
 Thr Pro Ser Asp Lys Gln Gly Met Pro Val Leu Ser Leu Asn Ile Thr
 590 595 600
 10 Cys Gln Met Asn Lys Thr Ile Ile Gly Val Ser Val Leu Ser Val Leu
 605 610 615
 Val Val Ser Val Val Ala Val Leu Val Tyr Lys Phe Tyr Phe His Leu
 15 620 625 630
 Met Leu Leu Ala Gly Cys Ile Lys Tyr Gly Arg Gly Glu Asn Ile Tyr
 635 640 645 650
 20 Asp Ala Phe Val Ile Tyr Ser Ser Gln Asp Glu Asp Trp Val Arg Asn
 655 660 665
 Glu Leu Val Lys Asn Leu Glu Glu Gly Val Pro Pro Phe Gln Leu Cys
 670 675 680
 25 Leu His Tyr Arg Asp Phe Ile Pro Gly Val Ala Ile Ala Ala Asn Ile
 685 690 695
 Ile His Glu Gly Phe His Lys Ser Arg Lys Val Ile Val Val Val Ser
 30 700 705 710
 Gln His Phe Ile Gln Ser Arg Trp Cys Ile Phe Glu Tyr Glu Ile Ala
 715 720 725 730
 35 Gln Thr Trp Gln Phe Leu Ser Ser Arg Ala Gly Ile Ile Phe Ile Val
 735 740 745
 Leu Gln Lys Val Glu Lys Thr Leu Leu Arg Gln Gln Val Glu Leu Tyr
 750 755 760
 40 Arg Leu Leu Ser Arg Asn Thr Tyr Leu Glu Trp Glu Asp Ser Val Leu
 765 770 775
 Gly Arg His Ile Phe Trp Arg Arg Leu Arg Lys Ala Leu Leu Asp Gly
 45 780 785 790
 Lys Ser Trp Asn Pro Glu Gly Thr Val Gly Thr Gly Cys Asn Trp Gln
 795 800 805 810
 50 Glu Ala Thr Ser Ile
 815

(2) INFORMATION FOR SEQ ID NO:27:

- 55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 300 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 60 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..300

(ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 186
 (D) OTHER INFORMATION: /note= "nucleotides 186, 196, 217, 276, and 300 designated C, each may be A, C, G, or T"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

15	TCC TAT TCT ATG GAA AAA GAT GCT TTC CTA TTT ATG AGA AAT TTG AAG	48
	Ser Tyr Ser Met Glu Lys Asp Ala Phe Leu Phe Met Arg Asn Leu Lys	
	1 5 10 15	
20	GTT CTC TCA CTA AAA GAT AAC AAT GTC ACA GCT GTC CCC ACC ACT TTG	96
	Val Leu Ser Leu Lys Asp Asn Asn Val Thr Ala Val Pro Thr Thr Leu	
	20 25 30	
25	CCA CCT AAT TTA CTA GAG CTC TAT CTT TAT AAC AAT ATC ATT AAG AAA	144
	Pro Pro Asn Leu Leu Glu Leu Tyr Leu Tyr Asn Asn Ile Ile Lys Lys	
	35 40 45	
30	ATC CAA GAA AAT GAT TTC AAT AAC CTC AAT GAG TTG CAA GTC CTT GAC	192
	Ile Gln Glu Asn Asp Phe Asn Asn Leu Asn Glu Leu Gln Val Leu Asp	
	50 55 60	
35	CTA CGT GGA AAT TGC CCT CGA TGT CAT AAT GTC CCA TAT CCG TGT ACA	240
	Leu Arg Gly Asn Cys Pro Arg Cys His Asn Val Pro Tyr Pro Cys Thr	
	65 70 75 80	
40	CCG TGT GAA AAT AAT TCC CCC TTA CAG ATC CAT GAC AAT GCT TTC AAT	288
	Pro Cys Glu Asn Asn Ser Pro Leu Gln Ile His Asp Asn Ala Phe Asn	
	85 90 95	
45	TCA TCG ACA GAC	300
	Ser Ser Thr Asp	
	100	

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 100 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

55	Ser Tyr Ser Met Glu Lys Asp Ala Phe Leu Phe Met Arg Asn Leu Lys
	1 5 10 15
60	Val Leu Ser Leu Lys Asp Asn Asn Val Thr Ala Val Pro Thr Thr Leu
	20 25 30

Pro Pro Asn Leu Leu Glu Leu Tyr Leu Tyr Asn Asn Ile Ile Lys Lys
35 40 45

Ile Gln Glu Asn Asp Phe Asn Asn Leu Asn Glu Leu Gln Val Leu Asp
5 50 55 60

Leu Arg Gly Asn Cys Pro Arg Cys His Asn Val Pro Tyr Pro Cys Thr
65 70 75 80

10 Pro Cys Glu Asn Asn Ser Pro Leu Gln Ile His Asp Asn Ala Phe Asn.
85 90 95

Ser Ser Thr Asp
100

15 (2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:
20 (A) LENGTH: 1756 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
25

(ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..1182
30

(ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1643
(D) OTHER INFORMATION: /note= "nucleotide 1643 designated
35 A, may be A or G"

(ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1664
40 (D) OTHER INFORMATION: /note= "nucleotide 1664 designated
C, may be A, C, G, or T"

(ix) FEATURE:
(A) NAME/KEY: misc_feature
45 (B) LOCATION: 1680
(D) OTHER INFORMATION: /note= "nucleotides 1680 and 1735
designated G, may be G or T"

(ix) FEATURE:
50 (A) NAME/KEY: misc_feature
(B) LOCATION: 1719
(D) OTHER INFORMATION: /note= "nucleotide 1719 designated
C, may be C or T"

(ix) FEATURE:
55 (A) NAME/KEY: misc_feature
(B) LOCATION: 1727
(D) OTHER INFORMATION: /note= "nucleotide 1727 designated
A, may be A, G, or T"
60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

5	TCT CCA GAA ATT CCC TGG AAT TCC TTG CCT CCT GAG GTT TTT GAG GGT Ser Pro Glu Ile Pro Trp Asn Ser Leu Pro Pro Glu Val Phe Glu Gly	48
10	ATG CCG CCA AAT CTA AAG AAT CTC TCC TTG GCC AAA AAT GGG CTC AAA Met Pro Pro Asn Leu Lys Asn Leu Ser Leu Ala Lys Asn Gly Leu Lys	96
15	TCT TTC TTT TGG GAC AGA CTC CAG TTA CTG AAG CAT TTG GAA ATT TTG Ser Phe Phe Trp Asp Arg Leu Gln Leu Leu Lys His Leu Glu Ile Leu	144
20	GAC CTC AGC CAT AAC CAG CTG ACA AAA GTA CCT GAG AGA TTG GCC AAC Asp Leu Ser His Asn Gln Leu Thr Lys Val Pro Glu Arg Leu Ala Asn	192
25	TGT TCC AAA AGT CTC ACA ACA CTG ATT CTT AAG CAT AAT CAA ATC AGG Cys Ser Lys Ser Leu Thr Thr Leu Ile Leu Lys His Asn Gln Ile Arg	240
30	CAA TTG ACA AAA TAT TTT CTA GAA GAT GCT TTG CAA TTG CGC TAT CTA Gln Leu Thr Lys Tyr Phe Leu Glu Asp Ala Leu Gln Leu Arg Tyr Leu	288
35	GAC ATC AGT TCA AAT AAA ATC CAG GTC ATT CAG AAG ACT AGC TTC CCA Asp Ile Ser Ser Asn Lys Ile Gln Val Ile Gln Lys Thr Ser Phe Pro	336
40	GAA AAT GTC CTC AAC AAT CTG GAG ATG TTG GTT TTA CAT CAC AAT CGC Glu Asn Val Leu Asn Asn Leu Glu Met Leu Val Leu His His Asn Arg	384
45	TTT CTT TGC AAC TGT GAT GCT GTG TGG TTT GTC TGG TGG GTT AAC CAT Phe Leu Cys Asn Cys Asp Ala Val Trp Phe Val Trp Trp Val Asn His	432
50	ACA GAT GTT ACT ATT CCA TAC CTG GCC ACT GAT GTG ACT TGT GTA GGT Thr Asp Val Thr Ile Pro Tyr Leu Ala Thr Asp Val Thr Cys Val Gly	480
55	CCA GGA GCA CAC AAA GGT CAA AGT GTC ATA TCC CTT GAT CTG TAT ACG Pro Gly Ala His Lys Gly Gln Ser Val Ile Ser Leu Asp Leu Tyr Thr	528
60	TGT GAG TTA GAT CTC ACA AAC CTG ATT CTG TTC TCA GTT TCC ATA TCA Cys Glu Leu Asp Leu Thr Asn Leu Ile Leu Phe Ser Val Ser Ile Ser	576
65	TCA GTC CTC TTT CTT ATG GTA GTT ATG ACA ACA AGT CAC CTC TTT TTC Ser Val Leu Phe Leu Met Val Val Met Thr Thr Ser His Leu Phe Phe	624
70	TGG GAT ATG TGG TAC ATT TAT TAT TTT TGG AAA GCA AAG ATA AAG GGG Trp Asp Met Trp Tyr Ile Tyr Tyr Phe Trp Lys Ala Lys Ile Lys Gly	672
75	TAT CCA GCA TCT GCA ATC CCA TGG AGT CCT TGT TAT GAT GCT TTT ATT Tyr Pro Ala Ser Ala Ile Pro Trp Ser Pro Cys Tyr Asp Ala Phe Ile	720

5	GTG TAT GAC ACT AAA AAC TCA GCT GTG ACA GAA TGG GTT TTG CAG GAG	768
	Val Tyr Asp Thr Lys Asn Ser Ala Val Thr Glu Trp Val Leu Gln Glu	
	245 250 255	
10	CTG GTG GCA AAA TTG GAA GAT CCA AGA GAA AAA CAC TTC AAT TTG TGT	816
	Leu Val Ala Lys Leu Glu Asp Pro Arg Glu Lys His Phe Asn Leu Cys	
	260 265 270	
15	CTA GAA GAA AGA GAC TGG CTA CCA GGA CAG CCA GTT CTA GAA AAC CTT	864
	Leu Glu Glu Arg Asp Trp Leu Pro Gly Gln Pro Val Leu Glu Asn Leu	
	275 280 285	
20	TCC CAG AGC ATA CAG CTC AGC AAA AAG ACA GTG TTT GTG ATG ACA CAG	912
	Ser Gln Ser Ile Gln Leu Ser Lys Lys Thr Val Phe Val Met Thr Gln	
	290 295 300	
25	AAA TAT GCT AAG ACT GAG AGT TTT AAG ATG GCA TTT TAT TTG TCT CAT	960
	Lys Tyr Ala Lys Thr Glu Ser Phe Lys Met Ala Phe Tyr Leu Ser His	
	305 310 315 320	
30	CAG AGG CTC CTG GAT GAA AAA GTG GAT GTG ATT ATC TTG ATA TTC TTG	1008
	Gln Arg Leu Leu Asp Glu Lys Val Asp Val Ile Ile Leu Ile Phe Leu	
	325 330 335	
35	GAA AGA CCT CTT CAG AAG TCT AAG TTT CTT CAG CTC AGG AAG AGA CTC	1056
	Glu Arg Pro Leu Gln Lys Ser Lys Phe Leu Gln Leu Arg Lys Arg Leu	
	340 345 350	
40	TGC AGG AGC TCT GTC CTT GAG TGG CCT GCA AAT CCA CAG GCT CAC CCA	1104
	Cys Arg Ser Ser Val Leu Glu Trp Pro Ala Asn Pro Gln Ala His Pro	
	355 360 365	
45	TAC TTC TGG CAG TGC CTG AAA AAT GCC CTG ACC ACA GAC AAT CAT GTG	1152
	Tyr Phe Trp Gln Cys Leu Lys Asn Ala Leu Thr Thr Asp Asn His Val	
	370 375 380	
50	GCT TAT AGT CAA ATG TTC AAG GAA ACA GTC TAGCTCTCTG AAGAATGTCA	1202
	Ala Tyr Ser Gln Met Phe Lys Glu Thr Val	
	385 390	
55	CCACCTAGGA CATGCCTTGG TACCTGAAGT TTTTCATAAAG GTTTCCATAA ATGAAGGTCT	1262
	GAATTTTTC TAACAGTTGT CATGGCTCAG ATTGGTGGGA AATCATCAAT ATATGGCTAA	1322
	GAAATTAAGA AGGGGAGACT GATAGAAGAT AATTTCTTTC TTCATGTGCC ATGCTCAGTT	1382
	AAATATTTTC CCTAGCTCAA ATCTGAAAAA CTGTGCCCTAG GAGACAACAC AAGGCTTTGA	1442
60	TTTATCTGCA TACAATTGAT AAGAGCCACA CATCTGCCCT GAAGAAGTAC TAGTAGTTTT	1502
	AGTAGTAGGG TAAAAATTAC ACAAGCTTTC TCTCTCTCTG ATACTGAACT GTACCAGAGT	1562
	TCAATGAAAT AAAAGCCCAG AGAACTTCTC AGTAAATGGT TTCATTATCA TGTAGTATCC	1622
	ACCATGCAAT ATGCCACAAA ACCGCTACTG GTACAGGACA GCTGGTAGCT GCTTCAAGGC	1682
	CTCTTATCAT TTTCTTGGGG CCCATGGAGG GGTCTCTCTG GAAAAAGGGA AGGTTTTTTT	1742
	TGGCCATCCA TGAA	1756

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 394 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Ser Pro Glu Ile Pro Trp Asn Ser Leu Pro Pro Glu Val Phe Glu Gly
 1 5 10 15
 Met Pro Pro Asn Leu Lys Asn Leu Ser Leu Ala Lys Asn Gly Leu Lys
 20 25 30
 Ser Phe Phe Trp Asp Arg Leu Gln Leu Leu Lys His Leu Glu Ile Leu
 35 40 45
 Asp Leu Ser His Asn Gln Leu Thr Lys Val Pro Glu Arg Leu Ala Asn
 50 55 60
 Cys Ser Lys Ser Leu Thr Thr Leu Ile Leu Lys His Asn Gln Ile Arg
 65 70 75 80
 Gln Leu Thr Lys Tyr Phe Leu Glu Asp Ala Leu Gln Leu Arg Tyr Leu
 85 90 95
 Asp Ile Ser Ser Asn Lys Ile Gln Val Ile Gln Lys Thr Ser Phe Pro
 100 105 110
 Glu Asn Val Leu Asn Asn Leu Glu Met Leu Val Leu His His Asn Arg
 115 120 125
 Phe Leu Cys Asn Cys Asp Ala Val Trp Phe Val Trp Trp Val Asn His
 130 135 140
 Thr Asp Val Thr Ile Pro Tyr Leu Ala Thr Asp Val Thr Cys Val Gly
 145 150 155 160
 Pro Gly Ala His Lys Gly Gln Ser Val Ile Ser Leu Asp Leu Tyr Thr
 165 170 175
 Cys Glu Leu Asp Leu Thr Asn Leu Ile Leu Phe Ser Val Ser Ile Ser
 180 185 190
 Ser Val Leu Phe Leu Met Val Val Met Thr Thr Ser His Leu Phe Phe
 195 200 205
 Trp Asp Met Trp Tyr Ile Tyr Tyr Phe Trp Lys Ala Lys Ile Lys Gly
 210 215 220
 Tyr Pro Ala Ser Ala Ile Pro Trp Ser Pro Cys Tyr Asp Ala Phe Ile
 225 230 235 240
 Val Tyr Asp Thr Lys Asn Ser Ala Val Thr Glu Trp Val Leu Gln Glu
 245 250 255
 Leu Val Ala Lys Leu Glu Asp Pro Arg Glu Lys His Phe Asn Leu Cys

260 265 270
 Leu Glu Glu Arg Asp Trp Leu Pro Gly Gln Pro Val Leu Glu Asn Leu
 275 280 285
 5 Ser Gln Ser Ile Gln Leu Ser Lys Lys Thr Val Phe Val Met Thr Gln
 290 295 300
 Lys Tyr Ala Lys Thr Glu Ser Phe Lys Met Ala Phe Tyr Leu Ser His
 10 305 310 315 320
 Gln Arg Leu Leu Asp Glu Lys Val Asp Val Ile Ile Leu Ile Phe Leu
 325 330 335
 15 Glu Arg Pro Leu Gln Lys Ser Lys Phe Leu Gln Leu Arg Lys Arg Leu
 340 345 350
 Cys Arg Ser Ser Val Leu Glu Trp Pro Ala Asn Pro Gln Ala His Pro
 355 360 365
 20 Tyr Phe Trp Gln Cys Leu Lys Asn Ala Leu Thr Thr Asp Asn His Val
 370 375 380
 Ala Tyr Ser Gln Met Phe Lys Glu Thr Val
 25 385 390

(2) INFORMATION FOR SEQ ID NO:31:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 999 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 35 (ii) MOLECULE TYPE: cDNA

 40 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 2..847

 45 (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 4
 (D) OTHER INFORMATION: /note= "nucleotides 4 and 23
 designated C, each may be A, C, G, or T"

 50 (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 650
 (D) OTHER INFORMATION: /note= "nucleotide 650 designated
 G, may be A or G"

 55 (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 715
 (D) OTHER INFORMATION: /note= "nucleotides 715, 825, and
 845 designated C, each may be C or T"
 60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

	C TCC GAT GCC AAG ATT CGG CAC CAG GCA TAT TCA GAG GTC ATG ATG	46
	Ser Asp Ala Lys Ile Arg His Gln Ala Tyr Ser Glu Val Met Met	
	1 5 10 15	
5	GTT GGA TGG TCA GAT TCA TAC ACC TGT GAA TAC CCT TTA AAC CTA AGG	94
	Val Gly Trp Ser Asp Ser Tyr Thr Cys Glu Tyr Pro Leu Asn Leu Arg	
	20 25 30	
10	GGA ACT AGG TTA AAA GAC GTT CAT CTC CAC GAA TTA TCT TGC AAC ACA	142
	Gly Thr Arg Leu Lys Asp Val His Leu His Glu Leu Ser Cys Asn Thr	
	35 40 45	
15	GCT CTG TTG ATT GTC ACC ATT GTG GTT ATT ATG CTA GTT CTG GGG TTG	190
	Ala Leu Leu Ile Val Thr Ile Val Val Ile Met Leu Val Leu Gly Leu	
	50 55 60	
20	GCT GTG GCC TTC TGC TGT CTC CAC TTT GAT CTG CCC TGG TAT CTC AGG	238
	Ala Val Ala Phe Cys Cys Leu His Phe Asp Leu Pro Trp Tyr Leu Arg	
	65 70 75	
25	ATG CTA GGT CAA TGC ACA CAA ACA TGG CAC AGG GTT AGG AAA ACA ACC	286
	Met Leu Gly Gln Cys Thr Gln Thr Trp His Arg Val Arg Lys Thr Thr	
	80 85 90 95	
30	CAA GAA CAA CTC AAG AGA AAT GTC CGA TTC CAC GCA TTT ATT TCA TAC	334
	Gln Glu Gln Leu Lys Arg Asn Val Arg Phe His Ala Phe Ile Ser Tyr	
	100 105 110	
35	AGT GAA CAT GAT TCT CTG TGG GTG AAG AAT GAA TTG ATC CCC AAT CTA	382
	Ser Glu His Asp Ser Leu Trp Val Lys Asn Glu Leu Ile Pro Asn Leu	
	115 120 125	
40	GAG AAG GAA GAT GGT TCT ATC TTG ATT TGC CTT TAT GAA AGC TAC TTT	430
	Glu Lys Glu Asp Gly Ser Ile Leu Ile Cys Leu Tyr Glu Ser Tyr Phe	
	130 135 140	
45	GAC CCT GGC AAA AGC ATT AGT GAA AAT ATT GTA AGC TTC ATT GAG AAA	478
	Asp Pro Gly Lys Ser Ile Ser Glu Asn Ile Val Ser Phe Ile Glu Lys	
	145 150 155	
50	AGC TAT AAG TCC ATC TTT GTT TTG TCT CCC AAC TTT GTC CAG AAT GAG	526
	Ser Tyr Lys Ser Ile Phe Val Leu Ser Pro Asn Phe Val Gln Asn Glu	
	160 165 170 175	
55	TGG TGC CAT TAT GAA TTC TAC TTT GCC CAC CAC AAT CTC TTC CAT GAA	574
	Trp Cys His Tyr Glu Phe Tyr Phe Ala His His Asn Leu Phe His Glu	
	180 185 190	
60	AAT TCT GAT CAC ATA ATT CTT ATC TTA CTG GAA CCC ATT CCA TTC TAT	622
	Asn Ser Asp His Ile Ile Leu Ile Leu Leu Glu Pro Ile Pro Phe Tyr	
	195 200 205	
65	TGC ATT CCC ACC AGG TAT CAT AAA CTG GAA GCT CTC CTG GAA AAA AAA	670
	Cys Ile Pro Thr Arg Tyr His Lys Leu Glu Ala Leu Leu Glu Lys Lys	
	210 215 220	
70	GCA TAC TTG GAA TGG CCC AAG GAT AGG CGT AAA TGT GGG CTT TTC TGG	718
	Ala Tyr Leu Glu Trp Pro Lys Asp Arg Arg Lys Cys Gly Leu Phe Trp	
	225 230 235	

GCA AAC CTT CGA GCT GCT GTT AAT GTT AAT GTA TTA GCC ACC AGA GAA 766
 Ala Asn Leu Arg Ala Ala Val Asn Val Asn Val Leu Ala Thr Arg Glu
 240 245 250 255

5 ATG TAT GAA CTG CAG ACA TTC ACA GAG TTA AAT GAA GAG TCT CGA GGT 814
 Met Tyr Glu Leu Gln Thr Phe Thr Glu Leu Asn Glu Glu Ser Arg Gly
 260 265 270

10 TCT ACA ATC TCT CTG ATG AGA ACA GAC TGT CTA TAAAATCCCA CAGTCCTTGG 867
 Ser Thr Ile Ser Leu Met Arg Thr Asp Cys Leu
 275 280

GAAGTTGGGG ACCACATACA CTGTTGGGAT GTACATTGAT ACAACCTTTA TGATGGCAAT 927

15 TTGACAATAT TTATTAAAAT AAAAAATGGT TATTCCTTC AAAAAAAAAA AAAAAAAAAA 987
 AAAAAAAAAA AA 999

20 (2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 282 amino acids

(B) TYPE: amino acid

25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

30 Ser Asp Ala Lys Ile Arg His Gln Ala Tyr Ser Glu Val Met Met Val
 1 5 10 15

35 Gly Trp Ser Asp Ser Tyr Thr Cys Glu Tyr Pro Leu Asn Leu Arg Gly
 20 25 30

Thr Arg Leu Lys Asp Val His Leu His Glu Leu Ser Cys Asn Thr Ala
 35 40 45

40 Leu Leu Ile Val Thr Ile Val Val Ile Met Leu Val Leu Gly Leu Ala
 50 55 60

Val Ala Phe Cys Cys Leu His Phe Asp Leu Pro Trp Tyr Leu Arg Met
 65 70 75 80

45 Leu Gly Gln Cys Thr Gln Thr Trp His Arg Val Arg Lys Thr Thr Gln
 85 90 95

50 Glu Gln Leu Lys Arg Asn Val Arg Phe His Ala Phe Ile Ser Tyr Ser
 100 105 110

Glu His Asp Ser Leu Trp Val Lys Asn Glu Leu Ile Pro Asn Leu Glu
 115 120 125

55 Lys Glu Asp Gly Ser Ile Leu Ile Cys Leu Tyr Glu Ser Tyr Phe Asp
 130 135 140

Pro Gly Lys Ser Ile Ser Glu Asn Ile Val Ser Phe Ile Glu Lys Ser
 145 150 155 160

60 Tyr Lys Ser Ile Phe Val Leu Ser Pro Asn Phe Val Gln Asn Glu Trp

165 170 175

Cys His Tyr Glu Phe Tyr Phe Ala His His Asn Leu Phe His Glu Asn
180 185 190

5 Ser Asp His Ile Ile Leu Ile Leu Leu Glu Pro Ile Pro Phe Tyr Cys
195 200 205

10 Ile Pro Thr Arg Tyr His Lys Leu Glu Ala Leu Leu Glu Lys Lys Ala
210 215 220

Tyr Leu Glu Trp Pro Lys Asp Arg Arg Lys Cys Gly Leu Phe Trp Ala
225 230 235 240

15 Asn Leu Arg Ala Ala Val Asn Val Asn Val Leu Ala Thr Arg Glu Met
245 250 255

Tyr Glu Leu Gln Thr Phe Thr Glu Leu Asn Glu Glu Ser Arg Gly Ser
260 265 270

20 Thr Ile Ser Leu Met Arg Thr Asp Cys Leu
275 280

(2) INFORMATION FOR SEQ ID NO:33:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1173 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

35 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..1008

(ix) FEATURE:
40 (A) NAME/KEY: misc_feature
(B) LOCATION: 854
(D) OTHER INFORMATION: /note= "nucleotide 854 designated
A, may be A or T"

45 (ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1171
(D) OTHER INFORMATION: /note= "nucleotides 1171 and 1172
designated C, each may be A, C, G, or T"

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

CTG CCT GCT GGC ACC CGG CTC CGG AGG CTG GAT GTC AGC TGC AAC AGC 48
55 Leu Pro Ala Gly Thr Arg Leu Arg Arg Leu Asp Val Ser Cys Asn Ser
1 5 10 15

ATC AGC TTC GTG GCC CCC GGC TTC TTT TCC AAG GCC AAG GAG CTG CGA 96
60 Ile Ser Phe Val Ala Pro Gly Phe Phe Ser Lys Ala Lys Glu Leu Arg
20 25 30

	GAG CTC AAC CTT AGC GCC AAC GCC CTC AAG ACA GTG GAC CAC TCC TGG	144
	Glu Leu Asn Leu Ser Ala Asn Ala Leu Lys Thr Val Asp His Ser Trp	
	35 40 45	
5	TTT GGG CCC CTG GCG AGT GCC CTG CAA ATA CTA GAT GTA AGC GCC AAC	192
	Phe Gly Pro Leu Ala Ser Ala Leu Gln Ile Leu Asp Val Ser Ala Asn	
	50 55 60	
10	CCT CTG CAC TGC GCC TGT GGG GCG GCC TTT ATG GAC TTC CTG CTG GAG	240
	Pro Leu His Cys Ala Cys Gly Ala Ala Phe Met Asp Phe Leu Leu Glu	
	65 70 75 80	
15	GTG CAG GCT GCC GTG CCC GGT CTG CCC AGC CGG GTG AAG TGT GGC AGT	288
	Val Gln Ala Ala Val Pro Gly Leu Pro Ser Arg Val Lys Cys Gly Ser	
	85 90 95	
20	CCG GGC CAG CTC CAG GGC CTC AGC ATC TTT GCA CAG GAC CTG CGC CTC	336
	Pro Gly Gln Leu Gln Gly Leu Ser Ile Phe Ala Gln Asp Leu Arg Leu	
	100 105 110	
20	TGC CTG GAT GAG GCC CTC TCC TGG GAC TGT TTC GCC CTC TCG CTG CTG	384
	Cys Leu Asp Glu Ala Leu Ser Trp Asp Cys Phe Ala Leu Ser Leu Leu	
	115 120 125	
25	GCT GTG GCT CTG GGC CTG GGT GTG CCC ATG CTG CAT CAC CTC TGT GGC	432
	Ala Val Ala Leu Gly Leu Gly Val Pro Met Leu His His Leu Cys Gly	
	130 135 140	
30	TGG GAC CTC TGG TAC TGC TTC CAC CTG TGC CTG GCC TGG CTT CCC TGG	480
	Trp Asp Leu Trp Tyr Cys Phe His Leu Cys Leu Ala Trp Leu Pro Trp	
	145 150 155 160	
35	CGG GGG CGG CAA AGT GGG CGA GAT GAG GAT GCC CTG CCC TAC GAT GCC	528
	Arg Gly Arg Gln Ser Gly Arg Asp Glu Asp Ala Leu Pro Tyr Asp Ala	
	165 170 175	
40	TTC GTG GTC TTC GAC AAA ACG CAG AGC GCA GTG GCA GAC TGG GTG TAC	576
	Phe Val Val Phe Asp Lys Thr Gln Ser Ala Val Ala Asp Trp Val Tyr	
	180 185 190	
40	AAC GAG CTT CGG GGG CAG CTG GAG GAG TGC CGT GGG CGC TGG GCA CTC	624
	Asn Glu Leu Arg Gly Gln Leu Glu Glu Cys Arg Gly Arg Trp Ala Leu	
	195 200 205	
45	CGC CTG TGC CTG GAG GAA CGC GAC TGG CTG CCT GGC AAA ACC CTC TTT	672
	Arg Leu Cys Leu Glu Glu Arg Asp Trp Leu Pro Gly Lys Thr Leu Phe	
	210 215 220	
50	GAG AAC CTG TGG GCC TCG GTC TAT GGC AGC CGC AAG ACG CTG TTT GTG	720
	Glu Asn Leu Trp Ala Ser Val Tyr Gly Ser Arg Lys Thr Leu Phe Val	
	225 230 235 240	
55	CTG GCC CAC ACG GAC CGG GTC AGT GGT CTC TTG CGC GCC AGC TTC CTG	768
	Leu Ala His Thr Asp Arg Val Ser Gly Leu Leu Arg Ala Ser Phe Leu	
	245 250 255	
60	CTG GCC CAG CAG CGC CTG CTG GAG GAC CGC AAG GAC GTC GTG GTG CTG	816
	Leu Ala Gln Gln Arg Leu Leu Glu Asp Arg Lys Asp Val Val Val Leu	
	260 265 270	
60	GTG ATC CTG AGC CCT GAC GGC CGC CGC TCC CGC TAC GAG CGG CTG CGC	864

Val Ile Leu Ser Pro Asp Gly Arg Arg Ser Arg Tyr Glu Arg Leu Arg
 275 280 285

5 CAG CGC CTC TGC CGC CAG AGT GTC CTC CTC TGG CCC CAC CAG CCC AGT 912
 Gln Arg Leu Cys Arg Gln Ser Val Leu Leu Trp Pro His Gln Pro Ser
 290 295 300

10 GGT CAG CGC AGC TTC TGG GCC CAG CTG GGC ATG GCC CTG ACC AGG GAC 960
 Gly Gln Arg Ser Phe Trp Ala Gln Leu Gly Met Ala Leu Thr Arg Asp
 305 310 315 320

AAC CAC CAC TTC TAT AAC CGG AAC TTC TGC CAG GGA CCC ACG GCC GAA 1008
 Asn His His Phe Tyr Asn Arg Asn Phe Cys Gln Gly Pro Thr Ala Glu
 325 330 335

15 TAGCCGTGAG CCGGAATCCT GCACGGTGCC ACCTCCACAC TCACCTCACC TCTGCCTGCC 1068

TGGTCTGACC CTCCCCTGCT CGCCTCCCTC ACCCCACACC TGACACAGAG CAGGCACTCA 1128

20 ATAAATGCTA CCGAAGGCTA AAAAAAAAAA AAAAAAAAAA AACCA 1173

(2) INFORMATION FOR SEQ ID NO:34:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 336 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

35 Leu Pro Ala Gly Thr Arg Leu Arg Arg Leu Asp Val Ser Cys Asn Ser
 1 5 10 15

Ile Ser Phe Val Ala Pro Gly Phe Phe Ser Lys Ala Lys Glu Leu Arg
 20 25 30

40 Glu Leu Asn Leu Ser Ala Asn Ala Leu Lys Thr Val Asp His Ser Trp
 35 40 45

Phe Gly Pro Leu Ala Ser Ala Leu Gln Ile Leu Asp Val Ser Ala Asn
 50 55 60

45 Pro Leu His Cys Ala Cys Gly Ala Ala Phe Met Asp Phe Leu Leu Glu
 65 70 75 80

50 Val Gln Ala Ala Val Pro Gly Leu Pro Ser Arg Val Lys Cys Gly Ser
 85 90 95

Pro Gly Gln Leu Gln Gly Leu Ser Ile Phe Ala Gln Asp Leu Arg Leu
 100 105 110

55 Cys Leu Asp Glu Ala Leu Ser Trp Asp Cys Phe Ala Leu Ser Leu Leu
 115 120 125

Ala Val Ala Leu Gly Leu Gly Val Pro Met Leu His His Leu Cys Gly
 130 135 140

60 Trp Asp Leu Trp Tyr Cys Phe His Leu Cys Leu Ala Trp Leu Pro Trp

145 150 155 160
 Arg Gly Arg Gln Ser Gly Arg Asp Glu Asp Ala Leu Pro Tyr Asp Ala
 165 170 175
 5 Phe Val Val Phe Asp Lys Thr Gln Ser Ala Val Ala Asp Trp Val Tyr
 180 185 190
 10 Asn Glu Leu Arg Gly Gln Leu Glu Glu Cys Arg Gly Arg Trp Ala Leu
 195 200 205
 Arg Leu Cys Leu Glu Glu Arg Asp Trp Leu Pro Gly Lys Thr Leu Phe
 210 215 220
 15 Glu Asn Leu Trp Ala Ser Val Tyr Gly Ser Arg Lys Thr Leu Phe Val
 225 230 235 240
 Leu Ala His Thr Asp Arg Val Ser Gly Leu Leu Arg Ala Ser Phe Leu
 245 250 255
 20 Leu Ala Gln Gln Arg Leu Leu Glu Asp Arg Lys Asp Val Val Val Leu
 260 265 270
 Val Ile Leu Ser Pro Asp Gly Arg Arg Ser Arg Tyr Glu Arg Leu Arg
 275 280 285
 25 Gln Arg Leu Cys Arg Gln Ser Val Leu Leu Trp Pro His Gln Pro Ser
 290 295 300
 30 Gly Gln Arg Ser Phe Trp Ala Gln Leu Gly Met Ala Leu Thr Arg Asp
 305 310 315 320
 Asn His His Phe Tyr Asn Arg Asn Phe Cys Gln Gly Pro Thr Ala Glu
 325 330 335
 35

(2) INFORMATION FOR SEQ ID NO:35:

40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 497 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: cDNA

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

TGGCCCACAC GGACCGCGTC AGTGGCCTCC TGCGCACCAG CTTCTGCTG GCTCAGCAGC 60
 GCCTGTTGGA AGACCGCAAG GACGTGGTGG TGTTGGTGAT CCTGCGTCCG GATGCCCCAC 120
 55 CGTCCCGCTA TGTGCGACTG CGCCAGCGTC TCTGCCGCCA GAGTGTGCTC TTCTGGCCCC 180
 AGCGACCCAA CGGGCAGGGG GGCTTCTGGG CCCAGCTGAG TACAGCCCTG ACTAGGGACA 240
 60 ACCGCCACTT CTATAACCAG AACTTCTGCC GGGGACCTAC AGCAGAATAG CTCAGAGCAA 300

CAGCTGGAAA	CAGCTGCATC	TTCATGTCTG	GTTCCCGAGT	TGCTCTGCCT	GCCTTGCTCT	360
GTCTTACTAC	ACCGCTATTT	GGCAAGTGC	CAATATATGC	TACCAAGCCA	CCAGGCCCAC	420
5 GGAGCAAAGG	TTGGCTGTAA	AGGGTAGTTT	TCTTCCCATG	CATCTTTCAG	GAGAGTGAAG	480
ATAGACACCA	AACCCAC					497

WHAT IS CLAIMED IS:

1. A substantially pure or recombinant DTLR2 protein or peptide which exhibits at least about 85% sequence
5 identity over a length of at least about 12 amino acids to SEQ ID NO: 4.
2. A substantially pure or recombinant DTLR3 protein or peptide which exhibits at least about 85% sequence
10 identity over a length of at least about 12 amino acids to SEQ ID NO: 6.
3. A substantially pure or recombinant DTLR4 protein or peptide which exhibits at least about 85% sequence
15 identity over a length of at least about 12 amino acids to SEQ ID NO: 26.
4. A substantially pure or recombinant DTLR5 protein or peptide which exhibits at least about 85% sequence
20 identity over a length of at least about 12 amino acids to SEQ ID NO: 10.
5. A substantially pure or recombinant DTLR6 protein or peptide which exhibits at least about 85% sequence
25 identity over a length of at least about 12 amino acids to SEQ ID NO: 12.
6. A substantially pure or recombinant DTLR7 protein or peptide which exhibits at least about 85% sequence
30 identity over a length of at least about 12 amino acids to SEQ ID NO: 16 or 18.
7. A substantially pure or recombinant DTLR8 protein or peptide which exhibits at least about 85% sequence
35 identity over a length of at least about 12 amino acids to SEQ ID NO: 32.

8. A substantially pure or recombinant DTLR9 protein or peptide which exhibits at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 22.
- 5 9. A substantially pure or recombinant DTLR10 protein or peptide which exhibits at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 34.
- 10 10. A fusion protein comprising the protein or peptide of any of claims 1-9.
- 15 11. A binding compound which specifically binds to the protein or peptide of any of claims 1-9.
12. The binding compound of claim 11 which is an antibody or antibody fragment.
- 20 13. A nucleic acid encoding the protein or peptide of any of claims 1-9.
14. An expression vector comprising the nucleic acid of claim 13.
- 25 15. A host cell comprising the vector of claim 14.
16. A process for recombinantly producing a polypeptide comprising culturing the host cell of claim 15 under conditions in which the polypeptide is expressed.
- 30

1/5


**FIG. 1**

FIG. 2A





FIG. 2B






FIG. 3

4/5




FIG. 4A




FIG. 4B



FIG. 4C




FIG. 4D




FIG. 5C




FIG. 5F




FIG. 5B




FIG. 5E




FIG. 5A




FIG. 5D

HPS Trailer Page
for
WEST

UserID: jsouaya
Printer: cm1_12e14_gbejptr

Summary

Document	Pages	Printed	Missed
WO009850547	171	171	0
Total (1)	171	171	0